

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
29 March 2001 (29.03.2001)

PCT

(10) International Publication Number  
**WO 01/21189 A1**

(51) International Patent Classification: **A61K 38/00**,  
38/04, 38/08, 38/10, 39/29, 39/295

MN 55902 (US). **KUBO, Ralph, T.** [US/US]; 6921 Pear Tree Drive, Carlsbad, CA 92009 (US). **GREY, Howard, M.** [US/US]; 1461 Caminito Batea, La Jolla, CA 92037 (US).

(21) International Application Number: PCT/US00/19774

(22) International Filing Date: 19 July 2000 (19.07.2000)

(74) Agents: **LOCKYER, Jean, M.** et al.; Townsend and Townsend and Crew LLP, Two Embarcadero Center, Eighth Floor, San Francisco, CA 94111-3834 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/357,737 19 July 1999 (19.07.1999) US

(71) Applicant (for all designated States except US): **EPIM-MUNE INC.** [US/US]; 5820 Nancy Ridge Drive, San Diego, CA 92121 (US).

(81) Designated States (national): AE, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GI, GM, KE, LS, MW, MZ, SD, SI, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

## Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: INDUCING CELLULAR IMMUNE RESPONSES TO HEPATITIS C VIRUS USING PEPTIDE AND NUCLEIC ACID COMPOSITIONS

(57) Abstract: This invention uses our knowledge of the mechanisms by which antigen is recognized by T cells to identify and prepare HCV epitopes, and to develop epitope-based vaccines directed towards HCV. More specifically, this application communicates our discovery of pharmaceutical compositions and methods of use in the prevention and treatment of HCV infection.



WO 01/21189 A1

**INDUCING CELLULAR IMMUNE RESPONSES TO HEPATITIS C VIRUS  
USING PEPTIDE AND NUCLEIC ACID COMPOSITIONS**

5

**FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT**

10        This invention was funded, in part, by the United States government under grants with the National Institutes of Health. The U.S. government has certain rights in this invention.

**INDEX**

- 15    I.     Background of the Invention  
      II.    Summary of the Invention  
      III.   Brief Description of the Figures  
      IV.   Detailed Description of the Invention  
          A.     Definitions  
20        B.     Stimulation of CTL and HTL responses  
          C.     Binding Affinity of Peptide Epitopes for HLA Molecules  
          D.     Peptide Epitope Binding Motifs and Supermotifs  
              1.     HLA-A1 supermotif  
              2.     HLA-A2 supermotif  
25        3.     HLA-A3 supermotif  
              4.     HLA-A24 supermotif  
              5.     HLA-B7 supermotif  
              6.     HLA-B27 supermotif  
              7.     HLA-B44 supermotif  
30        8.     HLA-B58 supermotif  
              9.     HLA-B62 supermotif  
              10.    HLA-A1 motif

- 11. HLA-A2.1 motif
- 12. HLA-A3 motif
- 13. HLA-A11 motif
- 14. HLA-A24 motif
- 5 15. HLA-DR-1-4-7 supermotif
- 16. HLA-DR3 motifs
- E. Enhancing Population Coverage of the Vaccine
- F. Immune Response-Stimulating Peptide Epitope Analogs
- G. Computer Screening of Protein Sequences from Disease-Related Antigens
- 10 for Supermotif- or Motif-Containing Epitopes
- H. Preparation of Peptide Epitopes
- I. Assays to Detect T-Cell Responses
- J. Use of Peptide Epitopes for Evaluating Immune Responses
- K. Vaccine Compositions
- 15 1. Minigene Vaccines
- 2. Combinations of CTL Peptides with Helper Peptides
- L. Administration of Vaccines for Therapeutic or Prophylactic Purposes
- M. Kits
- V. Examples
- 20 VI. Claims
- VII. Abstract

## I. BACKGROUND OF THE INVENTION

Hepatitis C virus (HCV) infection is a global human health problem with approximately 150,000 new reported cases each year in the U.S. alone. HCV is a single stranded RNA virus, and is the etiological agent identified in most cases of non-A, non-B post-transfusion and post-transplant hepatitis, and is a common cause of acute sporadic hepatitis (Choo *et al.*, *Science* 244:359, 1989; Kuo *et al.*, *Science* 244:362, 1989; and Alter *et al.*, in: *Current Perspective in Hepatology*, p. 83, 1989). It is estimated that more than 50% of patients infected with HCV become chronically infected and, of those, 20% develop cirrhosis of the liver within 20 years (Davis *et al.*, *New Engl. J. Med.* 321:1501,



1989; Alter *et al.*, in: *Current Perspective in Hepatology*, p. 83, 1989; Alter *et al.*, *New Engl. J. Med.* 327:1899, 1992; and Dienstag, J. L. *Gastroenterology* 85:430, 1983).

Moreover, the only therapy available for treatment of HCV infection is interferon- $\alpha$ .

Most patients are unresponsive, however, and among the responders, there is a high

5 recurrence rate within 6-12 months of cessation of treatment (Liang *et al.*, *J. Med. Virol.* 40:69, 1993). Ribavirin, a guanosine analog with a broad spectrum activity against many RNA and DNA viruses, has been shown in clinical trials to be effective against chronic HCV infection when used in combination with interferon- $\alpha$  (*see, e.g.*, Poynard *et al.*, *Lancet* 352:1426-1432, 1998; Reichard *et al.*, *Lancet* 351:83-87, 1998). However, the  
10 response rate is still well below 50%.

Virus-specific, human leukocyte antigen (HLA) class I-restricted cytotoxic T lymphocytes (CTL) are known to play a major role in the prevention and clearance of virus infections *in vivo* (Oldstone *et al.*, *Nature* 321:239, 1989; Jamieson *et al.*, *J. Virol.* 61:3930, 1987; Yap *et al.*, *Nature* 273:238, 1978; Lukacher *et al.*, *J. Exp. Med.* 160:814,  
15 1994; McMichael *et al.*, *N. Engl. J. Med.* 309:13, 1983; Sethi *et al.*, *J. Gen. Virol.* 64:443, 1983; Watari *et al.*, *J. Exp. Med.* 165:459, 1987; Yasukawa *et al.*, *J. Immunol.* 143:2051, 1989; Tigges *et al.*, *J. Virol.* 66:1622, 1993; Reddenhase *et al.*, *J. Virol.* 55:263, 1985; Quinnan *et al.*, *N. Engl. J. Med.* 307:6, 1982). HLA class I molecules are expressed on the surface of almost all nucleated cells. Following intracellular processing of antigens,  
20 epitopes from the antigens are presented as a complex with the HLA class I molecules on the surface of such cells. CTL recognize the peptide-HLA class I complex, which then results in the destruction of the cell bearing the HLA-peptide complex directly by the CTL and/or via the activation of non-destructive mechanisms *e.g.*, the production of interferon, that inhibit viral replication.

25 In view of the heterogeneous immune response observed with HCV infection, induction of a multi-specific cellular immune response directed simultaneously against multiple HCV epitopes appears to be important for the development of an efficacious vaccine against HCV. There is a need, however, to establish vaccine embodiments that elicit immune responses that correspond to responses seen in patients that clear HCV  
30 infection.

The information provided in this section is intended to disclose the presently understood state of the art as of the filing date of the present application. Information is included in this section which was generated subsequent to the priority date of this

application. Accordingly, information in this section is not intended, in any way, to delineate the priority date for the invention.

## II. SUMMARY OF THE INVENTION

5 This invention applies our knowledge of the mechanisms by which antigen is recognized by T cells, for example, to develop epitope-based vaccines directed towards HCV. More specifically, this application communicates our discovery of specific epitope pharmaceutical compositions and methods of use in the prevention and treatment of HCV infection.

10 Upon development of appropriate technology, the use of epitope-based vaccines has several advantages over current vaccines, particularly when compared to the use of whole antigens in vaccine compositions. There is evidence that the immune response to whole antigens is directed largely toward variable regions of the antigen, allowing for immune escape due to mutations. The epitopes for inclusion in an epitope-based vaccine  
15 are selected from conserved regions of viral or tumor-associated antigens, which thereby reduces the likelihood of escape mutants. Furthermore, immunosuppressive epitopes that may be present in whole antigens can be avoided with the use of epitope-based vaccines.

An additional advantage of an epitope-based vaccine approach is the ability to combine selected epitopes (CTL and HTL), and further, to modify the composition of the  
20 epitopes, achieving, for example, enhanced immunogenicity. Accordingly, the immune response can be modulated, as appropriate, for the target disease. Similar engineering of the response is not possible with traditional approaches.

Another major benefit of epitope-based immune-stimulating vaccines is their safety. The possible pathological side effects caused by infectious agents or whole  
25 protein antigens, which might have their own intrinsic biological activity, is eliminated.

An epitope-based vaccine also provides the ability to direct and focus an immune response to multiple selected antigens from the same pathogen. Thus, patient-by-patient variability in the immune response to a particular pathogen may be alleviated by inclusion of epitopes from multiple antigens from that pathogen in a vaccine composition. A  
30 "pathogen" may be an infectious agent or a tumor associated molecule.

One of the most formidable obstacles to the development of broadly efficacious epitope-based immunotherapeutics, however, has been the extreme polymorphism of HLA molecules. To date, effective non-genetically biased coverage of a population has been a task of considerable complexity; such coverage has required that epitopes be used

that are specific for HLA molecules corresponding to each individual HLA allele, therefore, impractically large numbers of epitopes would have to be used in order to cover ethnically diverse populations. Thus, there has existed a need for peptide epitopes that are bound by multiple HLA antigen molecules for use in epitope-based vaccines. The  
5 greater the number of HLA antigen molecules bound, the greater the breadth of population coverage by the vaccine.

Furthermore, as described herein in greater detail, a need has existed to modulate peptide binding properties, for example, so that peptides that are able to bind to multiple HLA antigens do so with an affinity that will stimulate an immune response.  
10 Identification of epitopes restricted by more than one HLA allele at an affinity that correlates with immunogenicity is important to provide thorough population coverage, and to allow the elicitation of responses of sufficient vigor to prevent or clear an infection in a diverse segment of the population. Such a response can also target a broad array of epitopes. The technology disclosed herein provides for such favored immune responses.

15 In a preferred embodiment, epitopes for inclusion in vaccine compositions of the invention are selected by a process whereby protein sequences of known antigens are evaluated for the presence of motif or supermotif-bearing epitopes. Peptides corresponding to a motif- or supermotif-bearing epitope are then synthesized and tested for the ability to bind to the HLA molecule that recognizes the selected motif. Those  
20 peptides that bind at an intermediate or high affinity *i.e.*, an  $IC_{50}$  (or a  $K_D$  value) of 500 nM or less for HLA class I molecules or 1000 nM or less for HLA class II molecules, are further evaluated for their ability to induce a CTL or HTL response. Immunogenic peptide epitopes are selected for inclusion in vaccine compositions.

Supermotif-bearing peptides may additionally be tested for the ability to bind to  
25 multiple alleles within the HLA supertype family. Moreover, peptide epitopes may be analogued to modify binding affinity and/or the ability to bind to multiple alleles within an HLA supertype.

The invention also includes an embodiment comprising a method for monitoring or evaluating an immune response to HCV in a patient having a known HLA-type, the  
30 method comprising incubating a T lymphocyte sample from the patient with a peptide composition comprising an HCV epitope consisting essentially of an amino acid sequence described in Tables VII to Table XX or Table XXII which binds the product of at least one HLA allele present in said patient, and detecting for the presence of a T lymphocyte

that binds to the peptide. A CTL peptide epitope may, for example, comprise a tetrameric complex.

An alternative modality for defining the peptide epitopes in accordance with the invention is to recite the physical properties, such as length; primary structure; or charge, which are correlated with binding to a particular allele-specific HLA molecule or group of allele-specific HLA molecules. A further modality for defining peptide epitopes is to recite the physical properties of an HLA binding pocket, or properties shared by several allele-specific HLA binding pockets (*e.g.* pocket configuration and charge distribution) and reciting that the peptide epitope fits and binds to said pocket or pockets.

As will be apparent from the discussion below, other methods and embodiments are also contemplated. Further, novel synthetic peptides produced by any of the methods described herein are also part of the invention.

### III. BRIEF DESCRIPTION OF THE FIGURES

Figure 1: Figure 1 provides a graph of total frequency of genotypes as a function of the number of HCV candidate epitopes bound by HLA-A and B molecules, in an average population.

Figure 2: Figure 2 illustrates the position of peptide epitopes in an experimental model minigene construct.

### IV. DETAILED DESCRIPTION OF THE INVENTION

The peptide epitopes and corresponding nucleic acid compositions of the present invention are useful for stimulating an immune response to HCV by stimulating the production of CTL or HTL responses. The peptide epitopes, which are derived directly or indirectly from native HCV amino acid sequences, are able to bind to HLA molecules and stimulate an immune response to HCV. The complete polyprotein sequence from HCV and its variants can be obtained from Genbank. Peptide epitopes and analogs thereof can also be readily determined from sequence information that may subsequently be discovered for heretofore unknown variants of HCV, as will be clear from the disclosure provided below.

The peptide epitopes of the invention have been identified in a number of ways, as will be discussed below. Also discussed in greater detail is that analog peptides have been derived and the binding activity for HLA molecules modulated by modifying specific amino acid residues to create peptide analogs exhibiting altered immunogenicity.

Further, the present invention provides compositions and combinations of compositions that enable epitope-based vaccines that are capable of interacting with HLA molecules encoded by various genetic alleles to provide broader population coverage than prior vaccines.

5

#### IV.A. Definitions

The invention can be better understood with reference to the following definitions, which are listed alphabetically:

A "computer" or "computer system" generally includes: a processor; at least one  
10 information storage/retrieval apparatus such as, for example, a hard drive, a disk drive or a tape drive; at least one input apparatus such as, for example, a keyboard, a mouse, a touch screen, or a microphone; and display structure. Additionally, the computer may include a communication channel in communication with a network. Such a computer may include more or less than what is listed above.

15 "Cross-reactive binding" indicates that a peptide is bound by more than one HLA molecule; a synonym is degenerate binding.

A "cryptic epitope" elicits a response by immunization with an isolated peptide, but the response is not cross-reactive *in vitro* when intact whole protein which comprises the epitope is used as an antigen.

20 A "dominant epitope" is an epitope that induces an immune response upon immunization with a whole native antigen (see, e.g., Sercarz, *et al.*, *Annu. Rev. Immunol.* 11:729-766, 1993). Such a response is cross-reactive *in vitro* with an isolated peptide epitope.

With regard to a particular amino acid sequence, an "epitope" is a set of amino  
25 acid residues which is involved in recognition by a particular immunoglobulin, or in the context of T cells, those residues necessary for recognition by T cell receptor proteins and/or Major Histocompatibility Complex (MHC) receptors. In an immune system setting, *in vivo* or *in vitro*, an epitope is the collective features of a molecule, such as primary, secondary and tertiary peptide structure, and charge, that together form a site  
30 recognized by an immunoglobulin, T cell receptor or HLA molecule. Throughout this disclosure epitope and peptide are often used interchangeably.

It is to be appreciated that protein or peptide molecules that comprise an epitope of the invention as well as additional amino acid(s) are still within the bounds of the invention. In certain embodiments, there is a limitation on the length of a peptide of the

invention which is not otherwise a construct. An embodiment that is length-limited occurs when the protein/peptide comprising an epitope of the invention comprises a region (i.e., a contiguous series of amino acids) having 100% identity with a native sequence. In order to avoid the definition of epitope from reading, *e.g.*, on whole natural molecules, there is a limitation on the length of any region that has 100% identity with a native peptide sequence. Thus, for a peptide comprising an epitope of the invention and a region with 100% identity with a native peptide sequence (and is not otherwise a construct), the region with 100% identity to a native sequence generally has a length of: less than or equal to 600 amino acids, often less than or equal to 500 amino acids, often less than or equal to 400 amino acids, often less than or equal to 250 amino acids, often less than or equal to 100 amino acids, often less than or equal to 85 amino acids, often less than or equal to 75 amino acids, often less than or equal to 65 amino acids, and often less than or equal to 50 amino acids. In certain embodiments, an "epitope" of the invention is comprised by a peptide having a region with less than 51 amino acids that has 100% identity to a native peptide sequence, in any increment of (49, 48, 47, 46, 45, 44, 43, 42, 41, 40, 39, 38, 37, 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5) down to 5 amino acids.

Accordingly, peptide or protein sequences longer than 600 amino acids are within the scope of the invention, so long as they do not comprise any contiguous sequence of more than 600 amino acids that have 100% identity with a native peptide sequence, if they are not otherwise a construct. For any peptide that has five contiguous residues or less that correspond to a native sequence, there is no limitation on the maximal length of that peptide in order to fall within the scope of the invention. It is presently preferred that a CTL epitope be less than 600 residues long in any increment down to eight amino acid residues.

"Human Leukocyte Antigen" or "HLA" is a human class I or class II Major Histocompatibility Complex (MHC) protein (*see, e.g.*, Stites, *et al.*, IMMUNOLOGY, 8<sup>TH</sup> ED., Lange Publishing, Los Altos, CA (1994).

An "HLA supertype or family", as used herein, describes sets of HLA molecules grouped on the basis of shared peptide-binding specificities. HLA class I molecules that share somewhat similar binding affinity for peptides bearing certain amino acid motifs are grouped into HLA supertypes. The terms HLA superfamily, HLA supertype family, HLA family, and HLA xx-like supertype molecules (where xx denotes a particular HLA type), are synonyms.

Throughout this disclosure, results are expressed in terms of "IC<sub>50</sub>'s." IC<sub>50</sub> is the concentration of peptide in a binding assay at which 50% inhibition of binding of a reference peptide is observed. Given the conditions in which the assays are run (*i.e.*, limiting HLA proteins and labeled peptide concentrations), these values approximate K<sub>D</sub> values. Assays for determining binding are described in detail, *e.g.*, in PCT publications WO 94/20127 and WO 94/03205. It should be noted that IC<sub>50</sub> values can change, often dramatically, if the assay conditions are varied, and depending on the particular reagents used (*e.g.*, HLA preparation, *etc.*). For example, excessive concentrations of HLA molecules will increase the apparent measured IC<sub>50</sub> of a given ligand.

Alternatively, binding is expressed relative to a reference peptide. Although as a particular assay becomes more, or less, sensitive, the IC<sub>50</sub>'s of the peptides tested may change somewhat, the binding relative to the reference peptide will not significantly change. For example, in an assay run under conditions such that the IC<sub>50</sub> of the reference peptide increases 10-fold, the IC<sub>50</sub> values of the test peptides will also shift approximately 10-fold. Therefore, to avoid ambiguities, the assessment of whether a peptide is a good, intermediate, weak, or negative binder is generally based on its IC<sub>50</sub>, relative to the IC<sub>50</sub> of a standard peptide.

Binding may also be determined using other assay systems including those using: live cells (*e.g.*, Ceppellini *et al.*, *Nature* 339:392, 1989; Christnick *et al.*, *Nature* 352:67, 1991; Busch *et al.*, *Int. Immunol.* 2:443, 1990; Hill *et al.*, *J. Immunol.* 147:189, 1991; del Guercio *et al.*, *J. Immunol.* 154:685, 1995), cell free systems using detergent lysates (*e.g.*, Cerundolo *et al.*, *J. Immunol.* 21:2069, 1991), immobilized purified MHC (*e.g.*, Hill *et al.*, *J. Immunol.* 152, 2890, 1994; Marshall *et al.*, *J. Immunol.* 152:4946, 1994), ELISA systems (*e.g.*, Reay *et al.*, *EMBO J.* 11:2829, 1992), surface plasmon resonance (*e.g.*, Khilko *et al.*, *J. Biol. Chem.* 268:15425, 1993); high flux soluble phase assays (Hammer *et al.*, *J. Exp. Med.* 180:2353, 1994), and measurement of class I MHC stabilization or assembly (*e.g.*, Ljunggren *et al.*, *Nature* 346:476, 1990; Schumacher *et al.*, *Cell* 62:563, 1990; Townsend *et al.*, *Cell* 62:285, 1990; Parker *et al.*, *J. Immunol.* 149:1896, 1992).

As used herein, "high affinity" with respect to HLA class I molecules is defined as binding with an IC<sub>50</sub>, or K<sub>D</sub> value, of 50 nM or less; "intermediate affinity" is binding with an IC<sub>50</sub> or K<sub>D</sub> value of between about 50 and about 500 nM. "High affinity" with respect to binding to HLA class II molecules is defined as binding with an IC<sub>50</sub> or K<sub>D</sub> value of 100 nM or less; "intermediate affinity" is binding with an IC<sub>50</sub> or K<sub>D</sub> value of between about 100 and about 1000 nM.

The terms "identical" or percent "identity," in the context of two or more peptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues that are the same, when compared and aligned for maximum correspondence over a comparison window, as measured using a sequence comparison algorithm or by manual alignment and visual inspection.

An "immunogenic peptide" or "peptide epitope" is a peptide that comprises an allele-specific motif or supermotif such that the peptide will bind an HLA molecule and induce a CTL and/or HTL response. Thus, immunogenic peptides of the invention are capable of binding to an appropriate HLA molecule and thereafter inducing an HLA-restricted cytotoxic or helper T cell response to the antigen from which the immunogenic peptide is derived.

The phrases "isolated" or "biologically pure" refer to material which is substantially or essentially free from components which normally accompany the material as it is found in its native state. Thus, isolated peptides in accordance with the invention preferably do not contain materials normally associated with the peptides in their *in situ* environment. An "isolated" epitope refers to an epitope that does not include the whole sequence of the antigen or polypeptide from which the epitope was derived. Typically the "isolated" epitope does not have attached thereto additional amino acids that result in a sequence that has 100% identity with a native sequence. The native sequence can be a sequence such as a tumor-associated antigen from which the epitope is derived.

"Major Histocompatibility Complex" or "MHC" is a cluster of genes that plays a role in control of the cellular interactions responsible for physiologic immune responses. In humans, the MHC complex is also known as the HLA complex. For a detailed description of the MHC and HLA complexes, see, Paul, FUNDAMENTAL IMMUNOLOGY, 3<sup>RD</sup> ED., Raven Press, New York, 1993.

The term "motif" refers to the pattern of residues in a peptide of defined length, usually a peptide of from about 8 to about 13 amino acids for a class I HLA motif and from about 6 to about 25 amino acids for a class II HLA motif, which is recognized by a particular HLA molecule. Peptide motifs are typically different for each protein encoded by each human HLA allele and differ in the pattern of the primary and secondary anchor residues.

A "negative binding residue" is an amino acid which, if present at certain positions (typically not primary anchor positions) in a peptide epitope, results in decreased binding affinity of the peptide for the peptide's corresponding HLA molecule.



A "non-native" sequence or "construct" refers to a sequence that is not found in nature ("non-naturally occurring"). Such sequences include, *e.g.*, peptides that are lipidated or otherwise modified and polyepitopic compositions that contain epitopes that are non contiguous in a native protein sequence.

5       The term "peptide" is used interchangeably with "oligopeptide" in the present specification to designate a series of residues, typically L-amino acids, connected one to the other, typically by peptide bonds between the  $\alpha$ -amino and carboxyl groups of adjacent amino acids. The preferred CTL-inducing peptides of the invention are 13 residues or less in length and usually consist of between about 8 and about 11 residues,  
10       preferably 9 or 10 residues. The preferred HTL-inducing oligopeptides are less than about 50 residues in length and usually consist of between about 6 and about 30 residues, more usually between about 12 and 25, and often between about 15 and 20 residues.

"Pharmaceutically acceptable" refers to a generally non-toxic, inert, and/or physiologically compatible composition.

15       A "pharmaceutical excipient" comprises a material such as an adjuvant, a carrier, pH-adjusting and buffering agents, tonicity adjusting agents, wetting agents, preservative, and the like.

A "primary anchor residue" is an amino acid at a specific position along a peptide sequence which is understood to provide a contact point between the immunogenic  
20       peptide and the HLA molecule. One to three, usually two, primary anchor residues within a peptide of defined length generally defines a "motif" for an immunogenic peptide. These residues are understood to fit in close contact with peptide binding grooves of an HLA molecule, with their side chains buried in specific pockets of the binding grooves themselves. In one embodiment, the primary anchor residues are located  
25       at position 2 (from the amino terminal position) and at the carboxyl terminal position of a 9-residue peptide epitope in accordance with the invention. The primary anchor positions for each motif and supermotif are set forth in Table 1. For example, analog peptides can be created by altering the presence or absence of particular residues in these primary anchor positions. Such analogs are used to modulate the binding affinity of a peptide  
30       comprising a particular motif or supermotif.

"Promiscuous recognition" is where a distinct peptide is recognized by the same T cell clone in the context of various HLA molecules. Promiscuous recognition or binding is synonymous with cross-reactive binding.

A "protective immune response" or "therapeutic immune response" refers to a CTL and/or an HTL response to an antigen derived from an infectious agent or a tumor antigen, which prevents or at least partially arrests disease symptoms or progression. The immune response may also include an antibody response which has been facilitated by the stimulation of helper T cells.

The term "residue" refers to an amino acid or amino acid mimetic incorporated into an oligopeptide by an amide bond or amide bond mimetic.

A "secondary anchor residue" is an amino acid at a position other than a primary anchor position in a peptide which may influence peptide binding. A secondary anchor residue occurs at a significantly higher frequency amongst bound peptides than would be expected by random distribution of amino acids at one position. The secondary anchor residues are said to occur at "secondary anchor positions." A secondary anchor residue can be identified as a residue which is present at a higher frequency among high or intermediate affinity binding peptides, or a residue otherwise associated with high or intermediate affinity binding. For example, analog peptides can be created by altering the presence or absence of particular residues in these secondary anchor positions. Such analogs are used to finely modulate the binding affinity of a peptide comprising a particular motif or supermotif.

A "subdominant epitope" is an epitope which evokes little or no response upon immunization with whole antigens which comprise the epitope, but for which a response can be obtained by immunization with an isolated peptide, and this response (unlike the case of cryptic epitopes) is detected when whole protein is used to recall the response *in vitro* or *in vivo*.

A "supermotif" is a peptide binding specificity shared by HLA molecules encoded by two or more HLA alleles. Preferably, a supermotif-bearing peptide is recognized with high or intermediate affinity (as defined herein) by two or more HLA antigens.

"Synthetic peptide" refers to a peptide that is man-made using such methods as chemical synthesis or recombinant DNA technology.

As used herein, a "vaccine" is a composition that contains one or more peptides of the invention. There are numerous embodiments of vaccines in accordance with the invention, such as by a cocktail of one or more peptides; one or more epitopes of the invention comprised by a polyepitopic peptide; or nucleic acids that encode such peptides or polypeptides, e.g., a minigene that encodes a polyepitopic peptide. The "one or more peptides" can include, e.g., at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18,

19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100 or more peptides of the invention. The peptides or polypeptides can optionally be modified, such as by lipidation, addition of targeting or other sequences. HLA class I-binding peptides of the invention can be admixed with, or  
5 linked to, HLA class II-binding peptides, to facilitate activation of both cytotoxic T lymphocytes and helper T lymphocytes. Vaccines can also comprise peptide-pulsed antigen presenting cells, e.g., dendritic cells.

The nomenclature used to describe peptide compounds follows the conventional practice wherein the amino group is presented to the left (the N-terminus) and the  
10 carboxyl group to the right (the C-terminus) of each amino acid residue. When amino acid residue positions are referred to in a peptide epitope they are numbered in an amino to carboxyl direction with position one being the position closest to the amino terminal end of the epitope, or the peptide or protein of which it may be a part. In the formulae representing selected specific embodiments of the present invention, the amino- and  
15 carboxyl-terminal groups, although not specifically shown, are in the form they would assume at physiologic pH values, unless otherwise specified. In the amino acid structure formulae, each residue is generally represented by standard three letter or single letter designations. The L-form of an amino acid residue is represented by a capital single letter or a capital first letter of a three-letter symbol, and the D-form for those amino acids  
20 having D-forms is represented by a lower case single letter or a lower case three letter symbol. Glycine has no asymmetric carbon atom and is simply referred to as "Gly" or G. Symbols for the amino acids are shown below.

Single Letter Symbol	Three Letter Symbol	Amino Acids
A	Ala	Alanine
C	Cys	Cysteine
D	Asp	Aspartic Acid
E	Glu	Glutamic Acid
F	Phe	Phenylalanine
G	Gly	Glycine
H	His	Histidine
I	Ile	Isoleucine
K	Lys	Lysine
L	Leu	Leucine
M	Met	Methionine
N	Asn	Asparagine
P	Pro	Proline
Q	Gln	Glutamine
R	Arg	Arginine
S	Ser	Serine
T	Thr	Threonine
V	Val	Valine
W	Trp	Tryptophan
Y	Tyr	Tyrosine

#### IV.B. Stimulation of CTL and HTL responses

The mechanism by which T cells recognize antigens has been delineated during the past ten years. Based on our understanding of the immune system we have developed efficacious peptide epitope vaccine compositions that can induce a therapeutic or prophylactic immune response to HCV in a broad population. For an understanding of the value and efficacy of the claimed compositions, a brief review of immunology-related technology is provided.

A complex of an HLA molecule and a peptidic antigen acts as the ligand recognized by HLA-restricted T cells (Buus, S. *et al.*, *Cell* 47:1071, 1986; Babbitt, B. P. *et al.*, *Nature* 317:359, 1985; Townsend, A. and Bodmer, H., *Annu. Rev. Immunol.* 7:601,

1989; Germain, R. N., *Annu. Rev. Immunol.* 11:403, 1993). Through the study of single amino acid substituted antigen analogs and the sequencing of endogenously bound, naturally processed peptides, critical residues that correspond to motifs required for specific binding to HLA antigen molecules have been identified and are described herein and are set forth in Tables I, II, and III (see also, e.g., Southwood, *et al.*, *J. Immunol.* 160:3363, 1998; Rammensee, *et al.*, *Immunogenetics* 41:178, 1995; Rammensee *et al.*, SYFPEITHI, access via web at : <http://134.2.96.221/scripts.hlaserver.dll/home.htm>; Sette, A. and Sidney, J. *Curr. Opin. Immunol.* 10:478, 1998; Engelhard, V. H., *Curr. Opin. Immunol.* 6:13, 1994; Sette, A. and Grey, H. M., *Curr. Opin. Immunol.* 4:79, 1992; Sinigaglia, F. and Hammer, J. *Curr. Biol.* 6:52, 1994; Ruppert *et al.*, *Cell* 74:929-937, 1993; Kondo *et al.*, *J. Immunol.* 155:4307-4312, 1995; Sidney *et al.*, *J. Immunol.* 157:3480-3490, 1996; Sidney *et al.*, *Human Immunol.* 45:79-93, 1996; Sette, A. and Sidney, J. *Immunogenetics*, in press, 1999).

Furthermore, x-ray crystallographic analysis of HLA-peptide complexes has revealed pockets within the peptide binding cleft of HLA molecules which accommodate, in an allele-specific mode, residues borne by peptide ligands; these residues in turn determine the HLA binding capacity of the peptides in which they are present. (See, e.g., Madden, D.R. *Annu. Rev. Immunol.* 13:587, 1995; Smith, *et al.*, *Immunity* 4:203, 1996; Fremont *et al.*, *Immunity* 8:305, 1998; Stern *et al.*, *Structure* 2:245, 1994; Jones, E.Y. *Curr. Opin. Immunol.* 9:75, 1997; Brown, J. H. *et al.*, *Nature* 364:33, 1993; Guo, H. C. *et al.*, *Proc. Natl. Acad. Sci. USA* 90:8053, 1993; Guo, H. C. *et al.*, *Nature* 360:364, 1992; Silver, M. L. *et al.*, *Nature* 360:367, 1992; Matsumura, M. *et al.*, *Science* 257:927, 1992; Madden *et al.*, *Cell* 70:1035, 1992; Fremont, D. H. *et al.*, *Science* 257:919, 1992; Saper, M. A., Bjorkman, P. J. and Wiley, D. C., *J. Mol. Biol.* 219:277, 1991.)

Accordingly, the definition of class I and class II allele-specific HLA binding motifs, or class I or class II supermotifs allows identification of regions within a protein that have the potential of binding particular HLA antigen(s).

The present inventors have found that the correlation of binding affinity with immunogenicity, which is disclosed herein, is an important factor to be considered when evaluating candidate peptides. Thus, by a combination of motif searches and HLA-peptide binding assays, candidates for epitope-based vaccines have been identified. After determining their binding affinity, additional confirmatory work can be performed to select, amongst these vaccine candidates, epitopes with preferred characteristics in terms of population coverage, antigenicity, and immunogenicity.

Various strategies can be utilized to evaluate immunogenicity, including:

- 1) Evaluation of primary T cell cultures from normal individuals (*see, e.g.,* Wentworth, P. A. *et al.*, *Mol. Immunol.* 32:603, 1995; Celis, E. *et al.*, *Proc. Natl. Acad. Sci. USA* 91:2105, 1994; Tsai, V. *et al.*, *J. Immunol.* 158:1796, 1997; Kawashima, I. *et al.*, *Human Immunol.* 59:1, 1998); This procedure involves the stimulation of peripheral blood lymphocytes (PBL) from normal subjects with a test peptide in the presence of antigen presenting cells *in vitro* over a period of several weeks. T cells specific for the peptide become activated during this time and are detected using, *e.g.,* a  $^{51}\text{Cr}$ -release assay involving peptide sensitized target cells.
- 2) Immunization of HLA transgenic mice (*see, e.g.,* Wentworth, P. A. *et al.*, *J. Immunol.* 26:97, 1996; Wentworth, P. A. *et al.*, *Int. Immunol.* 8:651, 1996; Alexander, J. *et al.*, *J. Immunol.* 159:4753, 1997); In this method, peptides in incomplete Freund's adjuvant are administered subcutaneously to HLA transgenic mice. Several weeks following immunization, splenocytes are removed and cultured *in vitro* in the presence of test peptide for approximately one week. Peptide-specific T cells are detected using, *e.g.,* a  $^{51}\text{Cr}$ -release assay involving peptide sensitized target cells and target cells expressing endogenously generated antigen.
- 3) Demonstration of recall T cell responses from immune individuals who have effectively been vaccinated, recovered from infection, and/or from chronically infected patients (*see, e.g.,* Rehmann, B. *et al.*, *J. Exp. Med.* 181:1047, 1995; Doolan, D. L. *et al.*, *Immunity* 7:97, 1997; Bertoni, R. *et al.*, *J. Clin. Invest.* 100:503, 1997; Threlkeld, S. C. *et al.*, *J. Immunol.* 159:1648, 1997; Diepolder, H. M. *et al.*, *J. Virol.* 71:6011, 1997). In applying this strategy, recall responses are detected by culturing PBL from subjects that have been naturally exposed to the antigen, for instance through infection, and thus have generated an immune response "naturally", or from patients who were vaccinated against the infection. PBL from subjects are cultured *in vitro* for 1-2 weeks in the presence of test peptide plus antigen presenting cells (APC) to allow activation of "memory" T cells, as compared to "naive" T cells. At the end of the culture period, T cell activity is detected using assays for T cell activity including  $^{51}\text{Cr}$  release involving peptide-sensitized targets, T cell proliferation, or lymphokine release.

The following describes the peptide epitopes and corresponding nucleic acids of the invention.

#### IV.C. Binding Affinity of Peptide Epitopes for HLA Molecules

The large degree of HLA polymorphism is an important factor to consider with the epitope-based approach to vaccine development. To address this factor, epitope selection including identification of peptides capable of binding at high or intermediate  
5 affinity to multiple HLA molecules is often utilized, most preferably these epitopes bind at high or intermediate affinity to two or more allele specific HLA molecules.

CTL-inducing peptides of interest for vaccine compositions preferably include those that have an  $IC_{50}$  or binding affinity value for class I HLA molecules of 500 nM or better (*i.e.*, the value is  $\leq 500$  nM). HTL-inducing peptides preferably include those that  
10 have an  $IC_{50}$  or binding affinity value for class II HLA molecules of 1000 nM or better, (*i.e.*, the value is  $\leq 1,000$  nM). For example, peptide binding is assessed by testing the capacity of a candidate peptide to bind to a purified HLA molecule *in vitro*. Peptides exhibiting high or intermediate affinity are then considered for further analysis. Selected peptides are tested on other members of the supertype family. In preferred embodiments,  
15 peptides that exhibit cross-reactive binding are then used in vaccines or in cellular screening analyses.

Higher HLA binding affinity is typically correlated with greater immunogenicity. Greater immunogenicity can be manifested in several different ways. Immunogenicity corresponds to whether an immune response is elicited at all, and to the vigor of any  
20 particular response, as well as to the extent of a population in which a response is elicited. For example, a peptide might elicit an immune response in a diverse array of the population, yet in no instance produce a vigorous response. In accordance with these principles, close to 90% of high binding peptides have been found to be immunogenic, as contrasted with about 50% of the peptides which bind with intermediate affinity.  
25 Moreover, higher binding affinity peptides leads to more vigorous immunogenic responses. As a result, less peptide is required to elicit a similar biological effect if a high affinity binding peptide is used. Thus, in preferred embodiments of the invention, high affinity binding epitopes are particularly useful.

The relationship between binding affinity for HLA class I molecules and  
30 immunogenicity of discrete peptide epitopes on bound antigens has been determined for the first time in the art by the present inventors. The correlation between binding affinity and immunogenicity was analyzed in two different experimental approaches (*see, e.g.*, Sette, *et al.*, *J. Immunol.* 153:5586-5592, 1994). In the first approach, the

immunogenicity of potential epitopes ranging in HLA binding affinity over a 10,000-fold range was analyzed in HLA-A\*0201 transgenic mice. In the second approach, the antigenicity of approximately 100 different hepatitis B virus (HBV)-derived potential epitopes, all carrying A\*0201 binding motifs, was assessed by using PBL from acute  
5 hepatitis patients. Pursuant to these approaches, it was determined that an affinity threshold value of approximately 500 nM (preferably 50 nM or less) determines the capacity of a peptide epitope to elicit a CTL response. These data are true for class I binding affinity measurements for naturally processed peptides and for synthesized T cell epitopes. These data also indicate the important role of determinant selection in the  
10 shaping of T cell responses (*see, e.g., Schaeffer et al. Proc. Natl. Acad. Sci. USA* 86:4649-4653, 1989).

An affinity threshold associated with immunogenicity in the context of HLA class II DR molecules has also been delineated (*see, e.g., Southwood et al. J. Immunology* 160:3363-3373, 1998). In order to define a biologically significant threshold of DR  
15 binding affinity, a database of the binding affinities of 32 DR-restricted epitopes for their restricting element (*i.e.*, the HLA molecule that binds the motif) was compiled. In approximately half of the cases (15 of 32 epitopes), DR restriction was associated with high binding affinities, *i.e.* binding affinity values of 100 nM or less. In the other half of the cases (16 of 32), DR restriction was associated with intermediate affinity (binding  
20 affinity values in the 100-1000 nM range). In only one of 32 cases was DR restriction associated with an IC<sub>50</sub> of 1000 nM or greater. Thus, 1000 nM can be defined as an affinity threshold associated with immunogenicity in the context of DR molecules.

The binding affinity of peptides for HLA molecules can be determined as described in Example 1, below.

25

#### IV.D. Peptide Epitope Binding Motifs and Supermotifs

In the past few years evidence has accumulated to demonstrate that a large fraction of HLA class I and class II molecules can be classified into a relatively few supertypes, each characterized by largely overlapping peptide binding repertoires, and  
30 consensus structures of the main peptide binding pockets.

For HLA molecule pocket analyses, the residues comprising the B and F pockets of HLA class I molecules as described in crystallographic studies were analyzed (*see, e.g., Guo, H. C. et al., Nature* 360:364, 1992; Saper, M. A., Bjorkman, P. J. and Wiley, D. C., *J. Mol. Biol.* 219:277, 1991; Madden, D. R., Garboczi, D. N. and Wiley, D. C.,



Cell 75:693, 1993; Parham, P., Adams, E. J., and Arnett, K. L., *Immunol. Rev.* 143:141, 1995). In these analyses, residues 9, 45, 63, 66, 67, 70, and 99 were considered to make up the B pocket; and the B pocket was deemed to determine the specificity for the amino acid residue in the second position of peptide ligands. Similarly, residues 77, 80, 81, and 116 were considered to determine the specificity of the F pocket; the F pocket was deemed to determine the specificity for the C-terminal residue of a peptide ligand bound by the HLA class I molecule.

Through the study of single amino acid substituted antigen analogs and the sequencing of endogenously bound, naturally processed peptides, critical residues required for allele-specific binding to HLA molecules have been identified. The presence of these residues correlates with binding affinity for HLA molecules. The identification of motifs and/or supermotifs that correlate with high and intermediate affinity binding is an important issue with respect to the identification of immunogenic peptide epitopes for the inclusion in a vaccine. Kast *et al.* (*J. Immunol.* 152:3904-3912, 1994) have shown that motif-bearing peptides account for 90% of the epitopes that bind to allele-specific HLA class I molecules. In this study all possible peptides of 9 amino acids in length and overlapping by eight amino acids (240 peptides), which cover the entire sequence of the E6 and E7 proteins of human papillomavirus type 16, were evaluated for binding to five allele-specific HLA molecules that are expressed at high frequency among different ethnic groups. This unbiased set of peptides allowed an evaluation of the predictive value of HLA class I motifs. From the set of 240 peptides, 22 peptides were identified that bound to an allele-specific HLA molecule with high or intermediate affinity. Of these 22 peptides, 20 (*i.e.* 91%) were motif-bearing. Thus, this study demonstrates the value of motifs for the identification of peptide epitopes for inclusion in a vaccine: application of motif-based identification techniques eliminates screening of 90% of the potential epitopes in a target antigen protein sequence.

Such peptide epitopes are identified in the Tables described below.

Peptides of the present invention may also comprise epitopes that bind to MHC class II DR molecules. A greater degree of heterogeneity in both size and binding frame position of the motif, relative to the N and C termini of the peptide, exists for class II peptide ligands. This increased heterogeneity of HLA class II peptide ligands is due to the structure of the binding groove of the HLA class II molecule which, unlike its class I counterpart, is open at both ends. Crystallographic analysis of HLA class II DRB\*0101-peptide complexes showed that the major energy of binding is contributed by peptide

residues complexed with complementary pockets on the DRB\*0101 molecules. An important anchor residue engages the deepest hydrophobic pocket (*see, e.g.*, Madden, D.R. *Ann. Rev. Immunol.* 13:587, 1995) and is referred to as position 1 (P1). P1 may represent the N-terminal residue of a class II binding peptide epitope, but more typically  
5 is flanked towards the N-terminus by one or more residues. Other studies have also pointed to an important role for the peptide residue in the 6<sup>th</sup> position towards the C-terminus, relative to P1, for binding to various DR molecules.

Thus, peptides of the present invention are identified by any one of several HLA-specific amino acid motifs (*see, e.g.*, Tables I-III). If the presence of the motif  
10 corresponds to the ability to bind several allele-specific HLA antigens, it is referred to as a supermotif. The HLA molecules that bind to peptides that possess a particular amino acid supermotif are collectively referred to as an HLA "supertype."

The peptide motifs and supermotifs described below, and summarized in Tables I-III, provide guidance for the identification and use of peptide epitopes in accordance with  
15 the invention.

Examples of peptide epitopes bearing a respective supermotif or motif are included in Tables as designated in the description of each motif or supermotif below. The Tables include a binding affinity ratio listing for some of the peptide epitopes. The ratio may be converted to IC<sub>50</sub> by using the following formula: IC<sub>50</sub> of the standard  
20 peptide/ratio = IC<sub>50</sub> of the test peptide (*i.e.*, the peptide epitope). The IC<sub>50</sub> values of standard peptides used to determine binding affinities for Class I peptides are shown in Table IV. The IC<sub>50</sub> values of standard peptides used to determine binding affinities for Class II peptides are shown in Table V. The peptides used as standards for the binding assays described herein are examples of standards; alternative standard peptides can also  
25 be used when performing such an analysis.

To obtain the peptide epitope sequences listed in each Table, protein sequence data from fourteen HCV isolates were evaluated for the presence of the designated supermotif or motif. The fourteen strains include HPCCGAA, HPCPLYPRE, HCV-H-CMR, HCV-J1, HPCGENANTI, HPCGENOM, HPCHUMR, HPCJCG, HPCJTA, HCV-  
30 J483, HCV-JK1, HCV-N, HPCPOLP, and HCV-J8. Peptide epitopes were additionally evaluated on the basis of their conservancy among these fourteen strains. A criterion for conservancy requires that the entire sequence of an HLA class I binding peptide be totally conserved in 79% of the sequences available for a specific protein. Similarly, a criterion for conservancy requires that the entire 9-mer core region of an HLA class II binding

peptide be totally conserved in 79% of the sequences available for a specific protein. The percent conservancy of the selected peptide epitopes is indicated on the Tables. The frequency, *i.e.* the number of strains of the fourteen strains in which the totally conserved peptide sequence was identified, is also shown. The "position" column in the Tables  
 5 designates the amino acid position of the HCV polyprotein that corresponds to the first amino acid residue of the epitope. The "number of amino acids" indicates the number of residues in the epitope sequence.

#### **HLA Class I Motifs Indicative of CTL Inducing Peptide Epitopes:**

10 The primary anchor residues of the HLA class I peptide epitope supermotifs and motifs delineated below are summarized in Table I. The HLA class I motifs set out in Table I(a) are those most particularly relevant to the invention claimed here. Primary and secondary anchor positions are summarized in Table II. Allele-specific HLA molecules that comprise HLA class I supertype families are listed in Table VI.

15

#### **IV.D.1. HLA-A1 supermotif**

The HLA-A1 supermotif is characterized by the presence in peptide ligands of a small (T or S) or hydrophobic (L, I, V, or M) primary anchor residue in position 2, and an aromatic (Y, F, or W) primary anchor residue at the C-terminal position of the epitope.  
 20 The corresponding family of HLA molecules that bind to the A1 supermotif (*i.e.*, the HLA-A1 supertype) includes at least A\*0101, A\*2601, A\*2602, A\*2501, and A\*3201 (*see, e.g.*, DiBrino, M. *et al.*, *J. Immunol.* 151:5930, 1993; DiBrino, M. *et al.*, *J. Immunol.* 152:620, 1994; Kondo, A. *et al.*, *Immunogenetics* 45:249, 1997). Other allele-specific HLA molecules predicted to be members of the A1 superfamily are shown in  
 25 Table VI. Peptides binding to each of the individual HLA proteins can be modulated by substitutions at primary and/or secondary anchor positions, preferably choosing respective residues specified for the supermotif.

Peptide epitopes that comprise the A1 supermotif are set forth in Table VII.

#### **30 IV.D.2. HLA-A2 supermotif**

Primary anchor specificities for allele-specific HLA-A2.1 molecules (Falk *et al.*, *Nature* 351:290-296, 1991; Hunt *et al.*, *Science* 255:1261-1263, 1992; Parker *et al.*, *J. Immunol.* 149:3580-3587, 1992) and cross-reactive binding within the HLA A2 family (Fruci *et al.*, *Human Immunol.* 38:187-192, 1993; Tanigaki *et al.*, *Human Immunol.*

39:155-162, 1994) have been described. The present inventors have defined additional primary anchor residues that determine cross-reactive binding to multiple allele-specific HLA A2 molecules (Ruppert *et al.*, *Cell* 74:929-937, 1993; Del Guercio *et al.*, *J. Immunol.* 154:685-693, 1995; Kast *et al.*, *J. Immunol.* 152:3904-3912, 1994). The HLA-  
5 A2 supermotif comprises peptide ligands with L, I, V, M, A, T, or Q as a primary anchor residue at position 2 and L, I, V, M, A, or T as a primary anchor residue at the C-terminal position of the epitope.

The corresponding family of HLA molecules (*i.e.*, the HLA-A2 supertype that binds these peptides) is comprised of at least: A\*0201, A\*0202, A\*0203, A\*0204,  
10 A\*0205, A\*0206, A\*0207, A\*0209, A\*0214, A\*6802, and A\*6901. Other allele-specific HLA molecules predicted to be members of the A2 superfamily are shown in Table VI. As explained in detail below, binding to each of the individual allele-specific HLA molecules can be modulated by substitutions at the primary anchor and/or secondary anchor positions, preferably choosing respective residues specified for the  
15 supermotif.

Peptide epitopes that comprise an A2 supermotif are set forth in Table VIII. The motifs comprising the primary anchor residues V, A, T, or Q at position 2 and L, I, V, A, or T at the C-terminal position are those most particularly relevant to the invention claimed herein.

20

#### IV.D.3. HLA-A3 supermotif

The HLA-A3 supermotif is characterized by the presence in peptide ligands of A, L, I, V, M, S, or, T as a primary anchor at position 2, and a positively charged residue, R or K, at the C-terminal position of the epitope (*e.g.*, in position 9 of 9-mers). Exemplary  
25 members of the corresponding family of HLA molecules (the HLA-A3 supertype) that bind the A3 supermotif include at least A\*0301, A\*1101, A\*3101, A\*3301, and A\*6801. Other allele-specific HLA molecules predicted to be members of the A3 superfamily are shown in Table VI. As explained in detail below, peptide binding to each of the individual allele-specific HLA proteins can be modulated by substitutions of amino acids  
30 at the primary and/or secondary anchor positions of the peptide, preferably choosing respective residues specified for the supermotif.

Peptide epitopes that comprise the A3 supermotif are set forth in Table IX.

#### IV.D.4. HLA-A24 supermotif

The HLA-A24 supermotif is characterized by the presence in peptide ligands of an aromatic (F, W, or Y) or hydrophobic aliphatic (L, I, V, M, or T) residue as a primary anchor in position 2, and Y, F, W, L, I, or M as primary anchor at the C-terminal position of the epitope. The corresponding family of HLA molecules that bind to the A24 supermotif (*i.e.*, the A24 supertype) includes at least A\*2402, A\*3001, and A\*2301. Other allele-specific HLA molecules predicted to be members of the A24 superfamily are shown in Table VI. Peptide binding to each of the allele-specific HLA molecules can be modulated by substitutions at primary and/or secondary anchor positions, preferably choosing respective residues specified for the supermotif.

Peptide epitopes that comprise the A24 supermotif are set forth in Table X.

#### IV.D.5. HLA-B7 supermotif

The HLA-B7 supermotif is characterized by peptides bearing proline in position 2 as a primary anchor, and a hydrophobic or aliphatic amino acid (L, I, V, M, A, F, W, or Y) as the primary anchor at the C-terminal position of the epitope. The corresponding family of HLA molecules that bind the B7 supermotif (*i.e.*, the HLA-B7 supertype) is comprised of at least twenty six HLA-B proteins including: B\*0702, B\*0703, B\*0704, B\*0705, B\*1508, B\*3501, B\*3502, B\*3503, B\*3504, B\*3505, B\*3506, B\*3507, B\*3508, B\*5101, B\*5102, B\*5103, B\*5104, B\*5105, B\*5301, B\*5401, B\*5501, B\*5502, B\*5601, B\*5602, B\*6701, and B\*7801 (*see, e.g.*, Sidney, *et al.*, *J. Immunol.* 154:247, 1995; Barber, *et al.*, *Curr. Biol.* 5:179, 1995; Hill, *et al.*, *Nature* 360:434, 1992; Rammensee, *et al.*, *Immunogenetics* 41:178, 1995). Other allele-specific HLA molecules predicted to be members of the B7 superfamily are shown in Table VI. As explained in detail below, peptide binding to each of the individual allele-specific HLA proteins can be modulated by substitutions at the primary and/or secondary anchor positions of the peptide, preferably choosing respective residues specified for the supermotif.

Peptide epitopes that comprise the B7 supermotif are set forth in Table XI.

#### IV.D.6. HLA-B27 supermotif

The HLA-B27 supermotif is characterized by the presence in peptide ligands of a positively charged (R, H, or K) residue as a primary anchor at position 2, and a hydrophobic (F, Y, L, W, M, I, A, or V) residue as a primary anchor at the C-terminal position of the epitope. Exemplary members of the corresponding family of HLA

molecules that bind to the B27 supermotif (*i.e.*, the B27 supertype) include at least B\*1401, B\*1402, B\*1509, B\*2702, B\*2703, B\*2704, B\*2705, B\*2706, B\*3801, B\*3901, B\*3902, and B\*7301. Other allele-specific HLA molecules predicted to be members of the B27 superfamily are shown in Table VI. Peptide binding to each of the  
5 allele-specific HLA molecules can be modulated by substitutions at primary and/or secondary anchor positions, preferably choosing respective residues specified for the supermotif.

Peptide epitopes that comprise the B27 supermotif are set forth in Table XII.

#### 10 IV.D.7. HLA-B44 supermotif

The HLA-B44 supermotif is characterized by the presence in peptide ligands of negatively charged (D or E) residues as a primary anchor in position 2, and hydrophobic residues (F, W, Y, L, I, M, V, or A) as a primary anchor at the C-terminal position of the epitope. Exemplary members of the corresponding family of HLA molecules that bind to  
15 the B44 supermotif (*i.e.*, the B44 supertype) include at least: B\*1801, B\*1802, B\*3701, B\*4001, B\*4002, B\*4006, B\*4402, B\*4403, and B\*4006. Peptide binding to each of the allele-specific HLA molecules can be modulated by substitutions at primary and/or secondary anchor positions; preferably choosing respective residues specified for the supermotif.

20

#### IV.D.8. HLA-B58 supermotif

The HLA-B58 supermotif is characterized by the presence in peptide ligands of a small aliphatic residue (A, S, or T) as a primary anchor residue at position 2, and an aromatic or hydrophobic residue (F, W, Y, L, I, V, M, or A) as a primary anchor residue  
25 at the C-terminal position of the epitope. Exemplary members of the corresponding family of HLA molecules that bind to the B58 supermotif (*i.e.*, the B58 supertype) include at least: B\*1516, B\*1517, B\*5701, B\*5702, and B\*5801. Other allele-specific HLA molecules predicted to be members of the B58 superfamily are shown in Table VI. Peptide binding to each of the allele-specific HLA molecules can be modulated by  
30 substitutions at primary and/or secondary anchor positions, preferably choosing respective residues specified for the supermotif.

Peptide epitopes that comprise the B58 supermotif are set forth in Table XIII.

**IV.D.9. HLA-B62 supermotif**

The HLA-B62 supermotif is characterized by the presence in peptide ligands of the polar aliphatic residue Q or a hydrophobic aliphatic residue (L, V, M, I, or P) as a primary anchor in position 2, and a hydrophobic residue (F, W, Y, M, I, V, L, or A) as a primary anchor at the C-terminal position of the epitope. Exemplary members of the corresponding family of HLA molecules that bind to the B62 supermotif (*i.e.*, the B62 supertype) include at least: B\*1501, B\*1502, B\*1513, and B5201. Other allele-specific HLA molecules predicted to be members of the B62 superfamily are shown in Table VI. Peptide binding to each of the allele-specific HLA molecules can be modulated by substitutions at primary and/or secondary anchor positions, preferably choosing respective residues specified for the supermotif.

Peptide epitopes that comprise the B62 supermotif are set forth in Table XIV.

**IV.D.10. HLA-A1 motif**

The HLA-A1 motif is characterized by the presence in peptide ligands of T, S, or M as a primary anchor residue at position 2 and the presence of Y as a primary anchor residue at the C-terminal position of the epitope. An alternative allele-specific A1 motif is characterized by a primary anchor residue at position 3 rather than position 2. This motif is characterized by the presence of D, E, A, or S as a primary anchor residue in position 3, and a Y as a primary anchor residue at the C-terminal position of the epitope. Peptide binding to HLA A1 can be modulated by substitutions at primary and/or secondary anchor positions, preferably choosing respective residues specified for the motif.

Peptide epitopes that comprise either A1 motif are set forth in Table XV. The epitopes comprising T, S, or M at position 2 and Y at the C-terminal position are also included in the listing of HLA-A1 supermotif-bearing peptide epitopes listed in Table VII.

**IV.D.11. HLA-A\*0201 motif**

An HLA-A2\*0201 motif was first determined to be characterized by the presence in peptide ligands of L or M as a primary anchor residue in position 2, and L or V as a primary anchor residue at the C-terminal position of a 9-residue peptide (Falk *et al.*, *Nature* 351:290-296, 1991). The A\*0201 motif was also determined to further comprise an I at position 2 and I or A at the C-terminal position of a nine amino acid peptide (Hunt

*et al.*, Science 255:1261-1263, March 6, 1992; Parker *et al.*, *J. Immunol.* 149:3580-3587, 1992). Subsequently, the A\*0201 allele-specific motif has been defined by the present inventors to additionally comprise V, A, T, or Q as a primary anchor residue at position 2, and M as a primary anchor residue at the C-terminal position of the epitope.

- 5 Additionally, the A\*0201 allele-specific motif has been found to comprise a T at the C-terminal position (Kast *et al.*, *J. Immunol.* 152:3904-3912, 1994). Thus, the HLA-A\*0201 motif comprises peptide ligands with L, I, V, M, A, T, or Q as primary anchor residues at position 2 and L, I, V, M, A, or T as a primary anchor residue at the C-terminal position of the epitope. The preferred and tolerated residues that characterize the
- 10 primary anchor positions of the HLA-A\*0201 motif are identical to the residues describing the A2 supermotif. (For reviews of relevant data, *see, e.g.*, Del Guercio *et al.*, *J. Immunol.* 154:685-693, 1995; Ruppert *et al.*, *Cell* 74:929-937, 1993; Sidney *et al.*, *Immunol. Today* 17:261-266, 1996; Sette and Sidney, *Curr. Opin. in Immunol.* 10:478-482, 1998). Secondary anchor residues that characterize the A\*0201 motif have
- 15 additionally been defined as disclosed herein. These are disclosed in Table II. Peptide binding to HLA-A\*0201 molecules can be modulated by substitutions at primary and/or secondary anchor positions, preferably choosing respective residues specified for the motif.

- Peptide epitopes that comprise an A\*0201 motif are set forth in Table VIII. The
- 20 A\*0201 motifs comprising the primary anchor residues V, A, T, or Q at position 2 and L, I, V, A, or T at the C-terminal position are those most particularly relevant to the invention claimed herein.

#### IV.D.12. HLA-A3 motif

- 25 The HLA-A3 motif is characterized by the presence in peptide ligands of L, M, V, I, S, A, T, F, C, G, or D as a primary anchor residue at position 2, and the presence of K, Y, R, H, F, or A as a primary anchor residue at the C-terminal position of the epitope. Peptide binding to HLA-A3 can be modulated by substitutions at primary and/or secondary anchor positions, preferably choosing respective residues specified for the
- 30 motif.

The A3 supermotif primary anchor residues comprise a subset of the A3- and A11-allele specific motif primary anchor residues. Peptide epitopes that comprise the A3 motif are set forth in Table XVI. Those peptide epitopes that also comprise the A3 supermotif are also listed in Table IX.



**IV.D.13. HLA-A11 motif**

The HLA-A11 motif is characterized by the presence in peptide ligands of V, T, M, L, I, S, A, G, N, C, D, or F as a primary anchor residue in position 2, and K, R, Y, or  
5 H as a primary anchor residue at the C-terminal position of the epitope. Peptide binding to HLA-A11 can be modulated by substitutions at primary and/or secondary anchor positions, preferably choosing respective residues specified for the motif.

Peptide epitopes that comprise the A11 motif are set forth in Table XVII; peptide epitopes comprising the A3 allele-specific motif are also present in this Table because of  
10 the overlap between the A3 and A11 motif primary anchor specificities. Further, those peptide epitopes that comprise the A3 supermotif are also listed in Table IX.

**IV.D.14. HLA-A24 motif**

The HLA-A24 motif is characterized by the presence in peptide ligands of Y, F, W, or M as a primary anchor residue in position 2, and F, L, I, or W as a primary anchor  
15 residue at the C-terminal position of the epitope. Peptide binding to HLA-A24 molecules can be modulated by substitutions at primary and/or secondary anchor positions; preferably choosing respective residues specified for the motif.

Peptide epitopes that comprise the A24 motif are set forth in Table XVIII. These  
20 epitopes are also listed in Table X, which sets forth HLA-A24-supermotif-bearing peptide epitopes, as the primary anchor residues characterizing the A24 allele-specific motif comprise a subset of the A24 supermotif primary anchor residues.

**HLA Class II Binding Motifs**

25 The primary and secondary anchor residues of the HLA class II peptide epitope supermotifs and motifs delineated below are summarized in Table III.

**IV.D.15. HLA DR-1-4-7 supermotif**

Motifs have also been identified for peptides that bind to three common HLA  
30 class II allele-specific HLA molecules: HLA DRB1\*0401, DRB1\*0101, and DRB1\*0701. Collectively, the common residues from these motifs delineate the HLA DR-1-4-7 supermotif. Peptides that bind to these DR molecules carry a supermotif characterized by a large aromatic or hydrophobic residue (Y, F, W, L, I, V, or M) as a primary anchor residue in position 1, and a small, non-charged residue (S, T, C, A, P, V,

I, L, or M) as a primary anchor residue in position 6 of a 9-mer core region. Allele-specific secondary effects and secondary anchors for each of these HLA types have also been identified. These are set forth in Table III. Peptide binding to HLA- DRB1\*0401, DRB1\*0101, and/or DRB1\*0701 can be modulated by substitutions at primary and/or secondary anchor positions, preferably choosing respective residues specified for the supermotif.

Conserved peptide epitopes *i.e.*, conserved in  $\geq 79\%$  ( $\geq 11/14$ ) of the HCV strains used for the present analysis, may be described as corresponding to epitopes containing a nine residue core comprising the DR-1-4-7 supermotif, and in which the 9 residue core is conserved in  $\geq 79\%$  (wherein position 1 of the motif is at position 1 of the nine residue core). Conserved 9-mer core regions are set forth in Table XIXa. Respective exemplary peptide epitopes of 15 amino acid residues in length, each of which comprise a conserved nine residue core, are also shown in section "a" of the Table. Cross-reactive binding data for exemplary 15-residue supermotif-bearing peptides are shown in Table XIXb.

#### IV.D.16. HLA DR3 motifs

Two alternative motifs (*i.e.*, submotifs) characterize peptide epitopes that bind to HLA-DR3 molecules. In the first motif (submotif DR3A) a large, hydrophobic residue (L, I, V, M, F, or Y) is present in anchor position 1 of a 9-mer core, and D is present as an anchor at position 4, towards the carboxyl terminus of the epitope. As in other class II motifs, core position 1 may or may not occupy the peptide N-terminal position.

The alternative DR3 submotif provides for lack of the large, hydrophobic residue at anchor position 1, and/or lack of the negatively charged or amide-like anchor residue at position 4, by the presence of a positive charge at position 6 towards the carboxyl terminus of the epitope. Thus, for the alternative allele-specific DR3 motif (submotif DR3B): L, I, V, M, F, Y, A, or Y is present at anchor position 1; D, N, Q, E, S, or T is present at anchor position 4; and K, R, or H is present at anchor position 6. Peptide binding to HLA-DR3 can be modulated by substitutions at primary and/or secondary anchor positions, preferably choosing respective residues specified for the motif.

Conserved 9-mer core regions (*i.e.*, those sequences that are conserved in at least 79% of the 14 HCV strains used for the analysis) corresponding to a nine residue sequence comprising the DR3A submotif (wherein position 1 of the motif is at position 1 of the nine residue core) are set forth in Table XXa. Respective exemplary peptide

epitopes of 15 amino acid residues in length, each of which comprise a conserved nine residue core, are also shown in Table XXa. Table XXb shows binding data of exemplary DR3 submotif A-bearing peptides.

Conserved 9-mer core regions (*i.e.*, those that are at least 79% conserved in the 14 HCV strains used for the analysis) comprising the DR3B submotif and respective exemplary 15-mer peptides comprising the DR3 submotif-B epitope are set forth in Table XXc. Table XXd shows binding data of exemplary DR3 submotif B-bearing peptides.

Each of the HLA class I or class II peptide epitopes set out in the Tables herein are deemed singly to be an inventive aspect of this application. Further, it is also an inventive aspect of this application that each peptide epitope may be used in combination with any other peptide epitope.

#### IV.E. Enhancing Population Coverage of the Vaccine

Vaccines that have broad population coverage are preferred because they are more commercially viable and generally applicable to the most people. Broad population coverage can be obtained using the peptides of the invention (and nucleic acid compositions that encode such peptides) through selecting peptide epitopes that bind to HLA alleles which, when considered in total, are present in most of the population. Table XXI lists the overall frequencies of the HLA class I supertypes in various ethnicities (Table XXIa) and the combined population coverage achieved by the A2-, A3-, and B7-supertypes (Table XXIb). The A2-, A3-, and B7 supertypes are each present on the average of over 40% in each of these five major ethnic groups. Coverage in excess of 80% is achieved with a combination of these supermotifs. These results suggest that effective and non-ethnically biased population coverage is achieved upon use of a limited number of cross-reactive peptides. Although the population coverage reached with these three main peptide specificities is high, coverage can be expanded to reach 95% population coverage and above, and more easily achieve truly multispecific responses upon use of additional supermotif or allele-specific motif bearing peptides.

The B44-, A1-, and A24-supertypes are present, on average, in a range from 25% to 40% in these major ethnic populations (Table XXIa). While less prevalent overall, the B27-, B58-, and B62 supertypes are each present with a frequency >25% in at least one major ethnic group (Table XXIa). Table XXIb summarizes the estimated prevalence of combinations of HLA supertypes that have been identified in five major ethnic groups.

The incremental coverage obtained by the inclusion of A1-, A24-, and B44-supertypes to the A2, A3, and B7 coverage, or all of the supertypes described herein, is shown.

The data presented herein, together with the previous definition of the A2-, A3-, and B7-supertypes, indicates that all antigens, with the possible exception of A29, B8, and B46, can be classified into a total of nine HLA supertypes. By including epitopes from the six most frequent supertypes, an average population coverage of 99% is obtained for five major ethnic groups..

#### IV.F. Immune Response-Stimulating Peptide Analogs

In general, CTL and HTL responses are not directed against all possible epitopes. Rather, they are restricted to a few "immunodominant" determinants (Zinkernagel, *et al.*, *Adv. Immunol.* 27:5159, 1979; Bennink, *et al.*, *J. Exp. Med.* 168:1935-1939, 1988; Rawle, *et al.*, *J. Immunol.* 146:3977-3984, 1991). It has been recognized that immunodominance (Benacerraf, *et al.*, *Science* 175:273-279, 1972) could be explained by either the ability of a given epitope to selectively bind a particular HLA protein (determinant selection theory) (Vitiello, *et al.*, *J. Immunol.* 131:1635, 1983); Rosenthal, *et al.*, *Nature* 267:156-158, 1977), or to be selectively recognized by the existing TCR (T cell receptor) specificities (repertoire theory) (Klein, J., *IMMUNOLOGY, THE SCIENCE OF SELF/NONSELF DISCRIMINATION*, John Wiley & Sons, New York, pp. 270-310, 1982). It has been demonstrated that additional factors, mostly linked to processing events, can also play a key role in dictating, beyond strict immunogenicity, which of the many potential determinants will be presented as immunodominant (Sercarz, *et al.*, *Annu. Rev. Immunol.* 11:729-766, 1993).

The concept of dominance and subdominance is relevant to immunotherapy of both infectious diseases and cancer. For example, in the course of chronic viral disease, recruitment of subdominant epitopes can be important for successful clearance of the infection, especially if dominant CTL or HTL specificities have been inactivated by functional tolerance, suppression, mutation of viruses and other mechanisms (Franco, *et al.*, *Curr. Opin. Immunol.* 7:524-531, 1995). In the case of cancer and tumor antigens, CTLs recognizing at least some of the highest binding affinity peptides might be functionally inactivated. Lower binding affinity peptides are preferentially recognized at these times, and may therefore be preferred in therapeutic or prophylactic anti-cancer vaccines.

In particular, it has been noted that a significant number of epitopes derived from known non-viral tumor associated antigens (TAA) bind HLA class I with intermediate affinity ( $IC_{50}$  in the 50-500 nM range). For example, it has been found that 8 of 15 known TAA peptides recognized by tumor infiltrating lymphocytes (TIL) or CTL bound in the 50-500 nM range. (These data are in contrast with estimates that 90% of known viral antigens were bound by HLA class I molecules with  $IC_{50}$  of 50 nM or less, while only approximately 10% bound in the 50-500 nM range (Sette, *et al.*, *J. Immunol.*, 153:558-5592, 1994). In the cancer setting this phenomenon is probably due to elimination or functional inhibition of the CTL recognizing several of the highest binding peptides, presumably because of T cell tolerization events.

Without intending to be bound by theory, it is believed that because T cells to dominant epitopes may have been clonally deleted, selecting subdominant epitopes may allow existing T cells to be recruited, which will then lead to a therapeutic or prophylactic response. However, the binding of HLA molecules to subdominant epitopes is often less vigorous than to dominant ones. Accordingly, there is a need to be able to modulate the binding affinity of particular immunogenic epitopes for one or more HLA molecules, and thereby to modulate the immune response elicited by the peptide, for example to prepare analog peptides which elicit a more vigorous response. This ability would greatly enhance the usefulness of peptide-based vaccines and therapeutic agents.

Although peptides with suitable cross-reactivity among all alleles of a superfamily are identified by the screening procedures described above, cross-reactivity is not always as complete as possible, and in certain cases procedures to increase cross-reactivity of peptides can be useful; moreover, such procedures can also be used to modify other properties of the peptides such as binding affinity or peptide stability. Having established the general rules that govern cross-reactivity of peptides for HLA alleles within a given motif or supermotif, modification (*i.e.*, analoging) of the structure of peptides of particular interest in order to achieve broader (or otherwise modified) HLA binding capacity can be performed. More specifically, peptides which exhibit the broadest cross-reactivity patterns, can be produced in accordance with the teachings herein. The present concepts related to analog generation are set forth in greater detail in co-pending U.S.S.N. 09/226,775 filed 1/6/99.

In brief, the strategy employed utilizes the motifs or supermotifs which correlate with binding to certain HLA molecules. The motifs or supermotifs are defined by having primary anchors, and in many cases secondary anchors. Analog peptides can be created

by substituting amino acid residues at primary anchor, secondary anchor, or at primary and secondary anchor positions. Generally, analogs are made for peptides that already bear a motif or supermotif. Preferred secondary anchor residues of supermotifs and motifs that have been defined for HLA class I and class II binding peptides are shown in  
5 Tables II and III, respectively.

For a number of the motifs or supermotifs in accordance with the invention, residues are defined which are deleterious to binding to allele-specific HLA molecules or members of HLA supertypes that bind the respective motif or supermotif (Tables II and III). Accordingly, removal of such residues that are detrimental to binding can be  
10 performed in accordance with the present invention. For example, in the case of the A3 supertype, when all peptides that have such deleterious residues are removed from the population of analyzed peptides, the incidence of cross-reactivity increases from 22% to 37% (*see, e.g., Sidney, J. et al., Hu. Immunol. 45:79, 1996*). Thus, one strategy to improve the cross-reactivity of peptides within a given supermotif is simply to delete one  
15 or more of the deleterious residues present within a peptide and substitute a small "neutral" residue such as Ala (that may not influence T cell recognition of the peptide). An enhanced likelihood of cross-reactivity is expected if, together with elimination of detrimental residues within a peptide, "preferred" residues associated with high affinity binding to an allele-specific HLA molecule or to multiple HLA molecules within a  
20 superfamily are inserted.

To ensure that an analog peptide, when used as a vaccine, actually elicits a CTL response to the native epitope *in vivo* (or, in the case of class II epitopes, elicits helper T cells that cross-react with the wild type peptides), the analog peptide may be used to immunize T cells *in vitro* from individuals of the appropriate HLA allele. Thereafter, the  
25 immunized cells' capacity to induce lysis of wild type peptide sensitized target cells is evaluated. It will be desirable to use as antigen presenting cells, cells that have been either infected, or transfected with the appropriate genes, or, in the case of class II epitopes only, cells that have been pulsed with whole protein antigens, to establish whether endogenously produced antigen is also recognized by the relevant T cells.

30 Another embodiment of the invention is to create analogs of weak binding peptides, to thereby ensure adequate numbers of cross-reactive cellular binders. Class I binding peptides exhibiting binding affinities of 500-5000 nM, and carrying an acceptable but suboptimal primary anchor residue at one or both positions can be "fixed" by

substituting preferred anchor residues in accordance with the respective supertype. The analog peptides can then be tested for crossbinding activity.

Another embodiment for generating effective peptide analogs involves the substitution of residues that have an adverse impact on peptide stability or solubility in, *e.g.*, a liquid environment. This substitution may occur at any position of the peptide epitope. For example, a cysteine (C) can be substituted out in favor of  $\alpha$ -amino butyric acid. Due to its chemical nature, cysteine has the propensity to form disulfide bridges and sufficiently alter the peptide structurally so as to reduce binding capacity. Substituting  $\alpha$ -amino butyric acid for C not only alleviates this problem, but actually improves binding and crossbinding capability in certain instances (*see, e.g.*, the review by Sette *et al.*, In: Persistent Viral Infections, Eds. R. Ahmed and I. Chen, John Wiley & Sons, England, 1999). Substitution of cysteine with  $\alpha$ -amino butyric acid may occur at any residue of a peptide epitope, *i.e.* at either anchor or non-anchor positions.

Representative analog peptides are set forth in Table XXII. The Table indicates the length and sequence of the analog peptide as well as the motif or supermotif, if appropriate. The information in the "Fixed Nomenclature" column indicates the residues substituted at the indicated position numbers for the respective analog.

#### IV.G. Computer Screening of Protein Sequences from Disease-Related Antigens for Supermotif- or Motif-Bearing Peptides

In order to identify supermotif- or motif-bearing epitopes in a target antigen, a native protein sequence, *e.g.*, a tumor-associated antigen, or sequences from an infectious organism, or a donor tissue for transplantation, is screened using a means for computing, such as an intellectual calculation or a computer, to determine the presence of a supermotif or motif within the sequence. The information obtained from the analysis of native peptide can be used directly to evaluate the status of the native peptide or may be utilized subsequently to generate the peptide epitope.

Computer programs that allow the rapid screening of protein sequences for the occurrence of the subject supermotifs or motifs are encompassed by the present invention; as are programs that permit the generation of analog peptides. These programs are implemented to analyze any identified amino acid sequence or operate on an unknown sequence and simultaneously determine the sequence and identify motif-bearing epitopes thereof; analogs can be simultaneously determined as well. Generally, the identified sequences will be from a pathogenic organism or a tumor-associated peptide. For

example, the target molecules considered herein include, without limitation, the core, S, E1, NS1/E2, NS2, NS3, NS4, and NS5 regions of HCV.

In cases where the sequence of multiple variants of the same target protein are available, peptides may also be selected on the basis of their conservancy. A presently preferred criterion for conservancy defines that the entire sequence of an HLA class I binding peptide or the entire 9-mer core of a class II binding peptide, be totally (*i.e.*, 100%) conserved in at least 79% of the sequences evaluated for a specific protein. This definition of conservancy has been employed herein; although, as appreciated by those in the art, lower or higher degrees of conservancy can be employed as appropriate for a given antigenic target.

It is important that the selection criteria utilized for prediction of peptide binding are as accurate as possible, to correlate most efficiently with actual binding. Prediction of peptides that bind, for example, to HLA-A\*0201, on the basis of the presence of the appropriate primary anchors, is positive at about a 30% rate (see, *e.g.*, Ruppert, J. *et al. Cell* 74:929, 1993). However, by extensively analyzing peptide-HLA binding data disclosed herein, data in related patent applications, and data in the art, the present inventors have developed a number of allele-specific polynomial algorithms that dramatically increase the predictive value over identification on the basis of the presence of primary anchor residues alone. These algorithms take into account not only the presence or absence of primary anchors, but also consider the positive or deleterious presence of secondary anchor residues (to account for the impact of different amino acids at different positions). The algorithms are essentially based on the premise that the overall affinity (or  $\Delta G$ ) of peptide-HLA interactions can be approximated as a linear polynomial function of the type:

$$\Delta G = a_{1i} \times a_{2i} \times a_{3i} \dots \times a_{ni}$$

where  $a_{ji}$  is a coefficient that represents the effect of the presence of a given amino acid ( $j$ ) at a given position ( $i$ ) along the sequence of a peptide of  $n$  amino acids. An important assumption of this method is that the effects at each position are essentially independent of each other. This assumption is justified by studies that demonstrated that peptides are bound to HLA molecules and recognized by T cells in essentially an extended conformation. Derivation of specific algorithm coefficients has been described, for example, in Gulukota, K. *et al.*, *J. Mol. Biol.* 267:1258, 1997.



Additional methods to identify preferred peptide sequences, which also make use of specific motifs, include the use of neural networks and molecular modeling programs (see, e.g., Milik *et al.*, *Nature Biotechnology* 16:753, 1998; Altuvia *et al.*, *Hum. Immunol.* 58:1, 1997; Altuvia *et al.*, *J. Mol. Biol.* 249:244, 1995; Buus, S. *Curr. Opin. Immunol.* 11:209-213, 1999; Brusic, V. *et al.*, *Bioinformatics* 14:121-130, 1998; Parker *et al.*, *J. Immunol.* 152:163, 1993; Meister *et al.*, *Vaccine* 13:581, 1995; Hammer *et al.*, *J. Exp. Med.* 180:2353, 1994; Sturniolo *et al.*, *Nature Biotechnol.* 17:555 1999).

For example, it has been shown that in sets of A\*0201 motif-bearing peptides containing at least one preferred secondary anchor residue while avoiding the presence of any deleterious secondary anchor residues, 69% of the peptides will bind A\*0201 with an IC<sub>50</sub> less than 500 nM (Ruppert, J. *et al. Cell* 74:929, 1993). These algorithms are also flexible in that cut-off scores may be adjusted to select sets of peptides with greater or lower predicted binding properties, as desired.

In utilizing computer screening to identify peptide epitopes, a protein sequence or translated sequence may be analyzed using software developed to search for motifs, for example the "FINDPATTERNS" program (Devereux, *et al. Nucl. Acids Res.* 12:387-395, 1984) or MotifSearch 1.4 software program (D. Brown, San Diego, CA) to identify potential peptide sequences containing appropriate HLA binding motifs. The identified peptides can be scored using customized polynomial algorithms to predict their capacity to bind specific HLA class I or class II alleles. As appreciated by one of ordinary skill in the art, a large array of computer programming software and hardware options are available in the relevant art which can be employed to implement the motifs of the invention in order to evaluate (e.g., without limitation, to identify epitopes, identify epitope concentration per peptide length, or to generate analogs) known or unknown peptide sequences.

In accordance with the procedures described above, HCV peptide epitopes and analogs thereof that are able to bind HLA supertype groups or allele-specific HLA molecules have been identified (Tables VII-XX; Table XXII).

#### IV.H. Preparation of Peptide Epitopes

Peptides in accordance with the invention can be prepared synthetically, by recombinant DNA technology or chemical synthesis, or from natural sources such as native tumors or pathogenic organisms. Peptide epitopes may be synthesized individually or as polyepitopic peptides. Although the peptide will preferably be substantially free of

other naturally occurring host cell proteins and fragments thereof, in some embodiments the peptides may be synthetically conjugated to native fragments or particles.

The peptides in accordance with the invention can be a variety of lengths, and either in their neutral (uncharged) forms or in forms which are salts. The peptides in  
5 accordance with the invention are either free of modifications such as glycosylation, side chain oxidation, or phosphorylation; or they contain these modifications, subject to the condition that modifications do not destroy the biological activity of the peptides as described herein.

The peptides of the invention can be prepared in a wide variety of ways. For the  
10 preferred relatively short size, the peptides can be synthesized in solution or on a solid support in accordance with conventional techniques. Various automatic synthesizers are commercially available and can be used in accordance with known protocols. (See, for example, Stewart & Young, SOLID PHASE PEPTIDE SYNTHESIS, 2D. ED., Pierce Chemical Co., 1984). Further, individual peptide epitopes can be joined using chemical ligation to  
15 produce larger peptides that are still within the bounds of the invention.

Alternatively, recombinant DNA technology can be employed wherein a nucleotide sequence which encodes an immunogenic peptide of interest is inserted into an expression vector, transformed or transfected into an appropriate host cell and cultivated under conditions suitable for expression. These procedures are generally known in the  
20 art, as described generally in Sambrook *et al.*, MOLECULAR CLONING, A LABORATORY MANUAL, Cold Spring Harbor Press, Cold Spring Harbor, New York (1989). Thus, recombinant polypeptides which comprise one or more peptide sequences of the invention can be used to present the appropriate T cell epitope.

The nucleotide coding sequence for peptide epitopes of the preferred lengths  
25 contemplated herein can be synthesized by chemical techniques, for example, the phosphotriester method of Matteucci, *et al.*, *J. Am. Chem. Soc.* 103:3185 (1981). Peptide analogs can be made simply by substituting the appropriate and desired nucleic acid base(s) for those that encode the native peptide sequence; exemplary nucleic acid substitutions are those that encode an amino acid defined by the motifs/supermotifs  
30 herein. The coding sequence can then be provided with appropriate linkers and ligated into expression vectors commonly available in the art, and the vectors used to transform suitable hosts to produce the desired fusion protein. A number of such vectors and suitable host systems are now available. For expression of the fusion proteins, the coding sequence will be provided with operably linked start and stop codons, promoter and

terminator regions and usually a replication system to provide an expression vector for expression in the desired cellular host. For example, promoter sequences compatible with bacterial hosts are provided in plasmids containing convenient restriction sites for insertion of the desired coding sequence. The resulting expression vectors are  
5 transformed into suitable bacterial hosts. Of course, yeast, insect or mammalian cell hosts may also be used, employing suitable vectors and control sequences.

It is often preferable that the peptide epitope be as small as possible while still maintaining substantially all of the immunologic activity of the native protein. When possible, it may be desirable to optimize HLA class I binding peptide epitopes of the  
10 invention to a length of about 8 to about 13 amino acid residues, preferably 9 to 10. HLA class II binding peptide epitopes may be optimized to a length of about 6 to about 30 amino acids in length, preferably to between about 13 and about 20 residues. Preferably, the peptide epitopes are commensurate in size with endogenously processed pathogen-derived peptides or tumor cell peptides that are bound to the relevant HLA molecules,  
15 however, the identification and preparation of peptides of other lengths can also be carried out using the techniques described herein.

In alternative embodiments, peptides of the invention can be linked as a polyepitopic peptide, or as a minigene that encodes a polyepitopic peptide.

In another embodiment, it is preferred to identify native peptide regions that  
20 contain a high concentration of class I and/or class II epitopes. Such a sequence is generally selected on the basis that it contains the greatest number of epitopes per amino acid length. It is to be appreciated that epitopes can be present in a frame-shifted manner, e.g. a 10 amino acid long peptide could contain two 9 amino acid long epitopes and one 10 amino acid long epitope; upon intracellular processing, each epitope can be exposed  
25 and bound by an HLA molecule upon administration of such a peptide. This larger, preferably multi-epitopic, peptide can be generated synthetically, recombinantly, or via cleavage from the native source.

#### **IV.I. Assays to Detect T-Cell Responses**

30 Once HLA binding peptides are identified, they can be tested for the ability to elicit a T-cell response. The preparation and evaluation of motif-bearing peptides are described in PCT publications WO 94/20127 and WO 94/03205. Briefly, peptides comprising epitopes from a particular antigen are synthesized and tested for their ability to bind to the appropriate HLA proteins. These assays may involve evaluating the

binding of a peptide of the invention to purified HLA class I molecules in relation to the binding of a radioiodinated reference peptide. Alternatively, cells expressing empty class I molecules (*i.e.* lacking peptide therein) may be evaluated for peptide binding by immunofluorescent staining and flow microfluorimetry. Other assays that may be used to evaluate peptide binding include peptide-dependent class I assembly assays and/or the inhibition of CTL recognition by peptide competition. Those peptides that bind to the class I molecule, typically with an affinity of 500 nM or less, are further evaluated for their ability to serve as targets for CTLs derived from infected or immunized individuals, as well as for their capacity to induce primary *in vitro* or *in vivo* CTL responses that can give rise to CTL populations capable of reacting with selected target cells associated with a disease. Corresponding assays are used for evaluation of HLA class II binding peptides. HLA class II motif-bearing peptides that are shown to bind, typically at an affinity of 1000 nM or less, are further evaluated for the ability to stimulate HTL responses.

Conventional assays utilized to detect T cell responses include proliferation assays, lymphokine secretion assays, direct cytotoxicity assays, and limiting dilution assays. For example, antigen-presenting cells that have been incubated with a peptide can be assayed for the ability to induce CTL responses in responder cell populations. Antigen-presenting cells can be normal cells such as peripheral blood mononuclear cells or dendritic cells. Alternatively, mutant non-human mammalian cell lines that are deficient in their ability to load class I molecules with internally processed peptides and that have been transfected with the appropriate human class I gene, may be used to test for the capacity of the peptide to induce *in vitro* primary CTL responses.

Peripheral blood mononuclear cells (PBMCs) may be used as the responder cell source of CTL precursors. The appropriate antigen-presenting cells are incubated with peptide, after which the peptide-loaded antigen-presenting cells are then incubated with the responder cell population under optimized culture conditions. Positive CTL activation can be determined by assaying the culture for the presence of CTLs that kill radio-labeled target cells, both specific peptide-pulsed targets as well as target cells expressing endogenously processed forms of the antigen from which the peptide sequence was derived.

More recently, a method has been devised which allows direct quantification of antigen-specific T cells by staining with Fluorescein-labelled HLA tetrameric complexes (Altman, J. D. *et al.*, *Proc. Natl. Acad. Sci. USA* 90:10330, 1993; Altman, J. D. *et al.*, *Science* 274:94, 1996). Other relatively recent technical developments include staining

for intracellular lymphokines, and interferon release assays or ELISPOT assays.

Tetramer staining, intracellular lymphokine staining and ELISPOT assays all appear to be at least 10-fold more sensitive than more conventional assays (Lalvani, A. *et al.*, *J. Exp. Med.* 186:859, 1997; Dunbar, P. R. *et al.*, *Curr. Biol.* 8:413, 1998; Murali-Krishna, K. *et al.*, *Immunity* 8:177, 1998).

HTL activation may also be assessed using such techniques known to those in the art such as T cell proliferation and secretion of lymphokines, *e.g.* IL-2 (*see, e.g.* Alexander *et al.*, *Immunity* 1:751-761, 1994).

Alternatively, immunization of HLA transgenic mice can be used to determine immunogenicity of peptide epitopes. Several transgenic mouse models including mice with human A2.1, A11 (which can additionally be used to analyze HLA-A3 epitopes), and B7 alleles have been characterized and others (*e.g.*, transgenic mice for HLA-A1 and A24) are being developed. HLA-DR1 and HLA-DR3 mouse models have also been developed. Additional transgenic mouse models with other HLA alleles may be generated as necessary. Mice may be immunized with peptides emulsified in Incomplete Freund's Adjuvant and the resulting T cells tested for their capacity to recognize peptide-pulsed target cells and target cells transfected with appropriate genes. CTL responses may be analyzed using cytotoxicity assays described above. Similarly, HTL responses may be analyzed using such assays as T cell proliferation or secretion of lymphokines.

Exemplary immunogenic peptide epitopes are set out in Table XXIII.

#### **IV.J. Use of Peptide Epitopes as Diagnostic Agents and for Evaluating Immune Responses**

In one embodiment of the invention, HLA class I and class II binding peptides as described herein can be used as reagents to evaluate an immune response. The immune response to be evaluated can be induced by using as an immunogen any agent that may result in the production of antigen-specific CTLs or HTLs that recognize and bind to the peptide epitope(s) to be employed as the reagent. The peptide reagent need not be used as the immunogen. Assay systems that can be used for such an analysis include relatively recent technical developments such as tetramers, staining for intracellular lymphokines and interferon release assays, or ELISPOT assays.

For example, a peptide of the invention may be used in a tetramer staining assay to assess peripheral blood mononuclear cells for the presence of antigen-specific CTLs following exposure to a tumor cell antigen or an immunogen. The HLA-tetrameric

complex is used to directly visualize antigen-specific CTLs (*see, e.g., Ogg et al., Science* 279:2103-2106, 1998; and Altman *et al., Science* 174:94-96, 1996) and determine the frequency of the antigen-specific CTL population in a sample of peripheral blood mononuclear cells. A tetramer reagent using a peptide of the invention may be generated  
5 as follows: A peptide that binds to an HLA molecule is refolded in the presence of the corresponding HLA heavy chain and  $\beta_2$ -microglobulin to generate a trimolecular complex. The complex is biotinylated at the carboxyl terminal end of the heavy chain at a site that was previously engineered into the protein. Tetramer formation is then induced by the addition of streptavidin. By means of fluorescently labeled streptavidin, the  
10 tetramer can be used to stain antigen-specific cells. The cells may then be identified, for example, by flow cytometry. Such an analysis may be used for diagnostic or prognostic purposes. Cells identified by the procedure can also be used for therapeutic purposes.

Peptides of the invention may also be used as reagents to evaluate immune recall responses. (*see, e.g., Bertoni et al., J. Clin. Invest.* 100:503-513, 1997 and Penna *et al., J. Exp. Med.* 174:1565-1570, 1991.) For example, patient PBMC samples from individuals  
15 with HCV infection may be analyzed for the presence of antigen-specific CTLs or HTLs using specific peptides. A blood sample containing mononuclear cells may be evaluated by cultivating the PBMCs and stimulating the cells with a peptide of the invention. After an appropriate cultivation period, the expanded cell population may be analyzed, for  
20 example, for cytotoxic activity (CTL) or for HTL activity.

The peptides may also be used as reagents to evaluate the efficacy of a vaccine. PBMCs obtained from a patient vaccinated with an immunogen may be analyzed using, for example, either of the methods described above. The patient is HLA typed, and peptide epitope reagents that recognize the allele-specific molecules present in that  
25 patient are selected for the analysis. The immunogenicity of the vaccine is indicated by the presence of epitope-specific CTLs and/or HTLs in the PBMC sample.

The peptides of the invention may also be used to make antibodies, using techniques well known in the art (*see, e.g. CURRENT PROTOCOLS IN IMMUNOLOGY*, Wiley/Greene, NY; and *Antibodies A Laboratory Manual*, Harlow and Lane, Cold Spring  
30 Harbor Laboratory Press, 1989), which may be useful as reagents to diagnose or monitor cancer. Such antibodies include those that recognize a peptide in the context of an HLA molecule, *i.e.*, antibodies that bind to a peptide-MHC complex.

#### IV.K. Vaccine Compositions

- Vaccines and methods of preparing vaccines that contain an immunogenically effective amount of one or more peptides as described herein are further embodiments of the invention. Once appropriately immunogenic epitopes have been defined, they can be sorted and delivered by various means, herein referred to as "vaccine" compositions.
- Such vaccine compositions can include, for example, lipopeptides (*e.g.*, Vitiello, A. *et al.*, *J. Clin. Invest.* 95:341, 1995), peptide compositions encapsulated in poly(DL-lactide-co-glycolide) ("PLG") microspheres (*see, e.g.*, Eldridge, *et al.*, *Molec. Immunol.* 28:287-294, 1991; Alonso *et al.*, *Vaccine* 12:299-306, 1994; Jones *et al.*, *Vaccine* 13:675-681, 1995), peptide compositions contained in immune stimulating complexes (ISCOMS) (*see, e.g.*, Takahashi *et al.*, *Nature* 344:873-875, 1990; Hu *et al.*, *Clin Exp Immunol.* 113:235-243, 1998), multiple antigen peptide systems (MAPs) (*see e.g.*, Tam, J. P., *Proc. Natl. Acad. Sci. U.S.A.* 85:5409-5413, 1988; Tam, J.P., *J. Immunol. Methods* 196:17-32, 1996), viral delivery vectors (Perkus, M. E. *et al.*, In: *Concepts in vaccine development*, Kaufmann, S. H. E., ed., p. 379, 1996; Chakrabarti, S. *et al.*, *Nature* 320:535, 1986; Hu, S. L. *et al.*, *Nature* 320:537, 1986; Kieny, M.-P. *et al.*, *AIDS Bio/Technology* 4:790, 1986; Top, F. H. *et al.*, *J. Infect. Dis.* 124:148, 1971; Chanda, P. K. *et al.*, *Virology* 175:535, 1990), particles of viral or synthetic origin (*e.g.*, Kofler, N. *et al.*, *J. Immunol. Methods.* 192:25, 1996; Eldridge, J. H. *et al.*, *Sem. Hematol.* 30:16, 1993; Faló, L. D., Jr. *et al.*, *Nature Med.* 7:649, 1995), adjuvants (Warren, H. S., Vogel, F. R., and Chedid, L. A. *Annu. Rev. Immunol.* 4:369, 1986; Gupta, R. K. *et al.*, *Vaccine* 11:293, 1993), liposomes (Reddy, R. *et al.*, *J. Immunol.* 148:1585, 1992; Rock, K. L., *Immunol. Today* 17:131, 1996), or, naked or particle absorbed cDNA (Ulmer, J. B. *et al.*, *Science* 259:1745, 1993; Robinson, H. L., Hunt, L. A., and Webster, R. G., *Vaccine* 11:957, 1993; Shiver, J. W. *et al.*, In: *Concepts in vaccine development*, Kaufmann, S. H. E., ed., p. 423, 1996; Cease, K. B., and Berzofsky, J. A., *Annu. Rev. Immunol.* 12:923, 1994 and Eldridge, J. H. *et al.*, *Sem. Hematol.* 30:16, 1993). Toxin-targeted delivery technologies, also known as receptor mediated targeting, such as those of Avant Immunotherapeutics, Inc. (Needham, Massachusetts) may also be used.
- Vaccines of the invention include nucleic acid-mediated modalities. DNA or RNA encoding one or more of the peptides of the invention can also be administered to a patient. This approach is described, for instance, in Wolff *et al.*, *Science* 247:1465 (1990) as well as U.S. Patent Nos. 5,580,859; 5,589,466; 5,804,566; 5,739,118; 5,736,524; 5,679,647; WO 98/04720; and in more detail below. Examples of DNA-based

delivery technologies include "naked DNA", facilitated (bupivacaine, polymers, peptide-mediated) delivery, cationic lipid complexes, and particle-mediated ("gene gun") or pressure-mediated delivery (*see, e.g.*, U.S. Patent No. 5,922,687).

For therapeutic or prophylactic immunization purposes, the peptides of the  
5 invention can also be expressed by viral or bacterial vectors. Examples of expression  
vectors include attenuated viral hosts, such as vaccinia or fowlpox. As an example of this  
approach, vaccinia virus is used as a vector to express nucleotide sequences that encode  
the peptides of the invention. Upon introduction into a host bearing a tumor, the  
recombinant vaccinia virus expresses the immunogenic peptide, and thereby elicits a host  
10 CTL and/or HTL response. Vaccinia vectors and methods useful in immunization  
protocols are described in, *e.g.*, U.S. Patent No. 4,722,848. Another vector is BCG  
(Bacille Calmette Guerin). BCG vectors are described in Stover *et al.*, *Nature* 351:456-  
460 (1991). A wide variety of other vectors useful for therapeutic administration or  
immunization of the peptides of the invention, *e.g.* adeno and adeno-associated virus  
15 vectors, retroviral vectors, *Salmonella typhi* vectors, detoxified anthrax toxin vectors, and  
the like, will be apparent to those skilled in the art from the description herein.

Furthermore, vaccines in accordance with the invention encompass compositions  
of one or more of the claimed peptide(s). A peptide can be present in a vaccine  
individually. Alternatively, the peptide can exist as a homopolymer comprising  
20 multiple copies of the same peptide, or as a heteropolymer of various peptides. Polymers  
have the advantage of increased immunological reaction and, where different peptide  
epitopes are used to make up the polymer, the additional ability to induce antibodies  
and/or CTLs that react with different antigenic determinants of the pathogenic organism  
or tumor-related peptide targeted for an immune response. The composition can be a  
25 naturally occurring region of an antigen or can be prepared, *e.g.*, recombinantly or by  
chemical synthesis.

Carriers that can be used with vaccines of the invention are well known in the art,  
and include, *e.g.*, thyroglobulin, albumins such as human serum albumin, tetanus toxoid,  
polyamino acids such as poly L-lysine, poly L-glutamic acid, influenza, hepatitis B virus  
30 core protein, and the like. The vaccines can contain a physiologically tolerable (*i.e.*,  
acceptable) diluent such as water, or saline, preferably phosphate buffered saline. The  
vaccines also typically include an adjuvant. Adjuvants such as incomplete Freund's  
adjuvant, aluminum phosphate, aluminum hydroxide, or alum are examples of materials  
well known in the art. Additionally, as disclosed herein, CTL responses can be primed by



conjugating peptides of the invention to lipids, such as tripalmitoyl-S-glycerylcysteinylserine (P<sub>3</sub>CSS).

Upon immunization with a peptide composition in accordance with the invention, via injection, aerosol, oral, transdermal, transmucosal, intrapleural, intrathecal, or other  
5 suitable routes, the immune system of the host responds to the vaccine by producing large amounts of CTLs and/or HTLs specific for the desired antigen. Consequently, the host becomes at least partially immune to later infection, or at least partially resistant to developing an ongoing chronic infection, or derives at least some therapeutic benefit when the antigen was tumor-associated.

10 In some embodiments it may be desirable to combine the class I peptide components with components that induce or facilitate neutralizing antibody responses to the target antigen of interest, particularly to viral envelope antigens. A preferred embodiment of such a composition comprises class I and class II epitopes in accordance with the invention. An alternative embodiment of such a composition comprises a class I  
15 and/or class II epitope in accordance with the invention, along with a PADRE™ (Epimmune, San Diego, CA) molecule (described, for example, in U.S. Patent Number 5,736,142).

A vaccine of the invention can also include antigen-presenting cells, such as dendritic cells, as a vehicle to present peptides of the invention. Vaccine compositions  
20 can be created *in vitro*, following dendritic cell mobilization and harvesting, whereby loading of dendritic cells occurs *in vitro*. For example, dendritic cells are transfected, e.g., with a minigene in accordance with the invention. The dendritic cell can then be administered to a patient to elicit immune responses *in vivo*.

Antigenic peptides are used to elicit a CTL and/or HTL response *ex vivo*, as well.  
25 The resulting CTL or HTL cells, can be used to treat tumors in patients that do not respond to other conventional forms of therapy, or will not respond to a therapeutic vaccine peptide or nucleic acid in accordance with the invention. *Ex vivo* CTL or HTL responses to a particular tumor-associated antigen are induced by incubating in tissue culture the patient's, or genetically compatible, CTL or HTL precursor cells together with  
30 a source of antigen-presenting cells (APC), such as dendritic cells, and the appropriate immunogenic peptide. After an appropriate incubation time (typically about 7-28 days), in which the precursor cells are activated and expanded into effector cells, the cells are infused back into the patient, where they will destroy (CTL) or facilitate destruction

(HTL) of their specific target cell (an infected cell or a tumor cell). Transfected dendritic cells may also be used as antigen presenting cells.

The vaccine compositions of the invention can also be used in combination with antiviral drugs such as interferon- $\alpha$ , or other treatments for viral infection.

5        Preferably, the following principles are utilized when selecting an array of epitopes for inclusion in a polyepitopic composition for use in a vaccine, or for selecting discrete epitopes to be included in a vaccine and/or to be encoded by nucleic acids such as a minigene. It is preferred that each of the following principles are balanced in order to make the selection. The multiple epitopes to be incorporated in a given vaccine  
10       composition may be, but need not be, contiguous in sequence in the native antigen from which the epitopes are derived.

       Preferably, the following principles are utilized when selecting an array of epitopes for inclusion in a polyepitopic composition for use in a vaccine, or for selecting discrete epitopes to be included in a vaccine and/or to be encoded by nucleic acids such as  
15       a minigene. Exemplary epitopes that may be utilized in a vaccine to treat or prevent HCV infection are set out in Tables XXVI-XXIX, and Table XXXII. It is preferred that each of the following principles are balanced in order to make the selection.

       1.)       Epitopes are selected which, upon administration, mimic immune responses that have been observed to be correlated with HCV clearance. For HLA Class I  
20       this includes 3-4 epitopes that come from at least one antigen of HCV. For HLA Class II a similar rationale is employed; again 3-4 epitopes are selected from at least one HCV antigen (*see e.g.*, Rosenberg *et al.*, *Science* 278:1447-1450).

       2.)       Epitopes are selected that have the requisite binding affinity established to be correlated with immunogenicity: for HLA Class I an  $IC_{50}$  of 500 nM or less, or for  
25       Class II an  $IC_{50}$  of 1000 nM or less.

       3.)       Sufficient supermotif bearing-peptides, or a sufficient array of allele-specific motif-bearing peptides, are selected to give broad population coverage. For example, it is preferable to have at least 80% population coverage. A Monte Carlo analysis, a statistical evaluation known in the art, can be employed to assess the breadth,  
30       or redundancy of, population coverage.

       4.)       When selecting epitopes from cancer-related antigens it is often preferred to select analogs because the patient may have developed tolerance to the native epitope.

When selecting epitopes for infectious disease-related antigens it is preferable to select either native or analoged epitopes.

5.) Of particular relevance are epitopes referred to as "nested epitopes."

Nested epitopes occur where at least two epitopes overlap in a given peptide sequence. A

5 nested peptide sequence can comprise both HLA class I and HLA class II epitopes.

When providing nested epitopes, it is preferable to provide a sequence that has the greatest number of epitopes per provided sequence. Preferably, one avoids providing a peptide that is any longer than the amino terminus of the amino terminal epitope and the carboxyl terminus of the carboxyl terminal epitope in the peptide. When providing a

10 longer peptide sequence, such as a sequence comprising nested epitopes, it is important to screen the sequence in order to insure that it does not have pathological or other deleterious biological properties.

6.) If a polyepitopic protein is created, or when creating a minigene, an objective is to generate the smallest peptide that encompasses the epitopes of interest.

15 This principle is similar, if not the same as that employed when selecting a peptide comprising nested epitopes. However, with an artificial polyepitopic peptide, the size minimization objective is balanced against the need to integrate any spacer sequences between epitopes in the polyepitopic protein. Spacer amino acid residues can be introduced to avoid junctional epitopes (an epitope recognized by the immune system, not

20 present in the target antigen, and only created by the man-made juxtaposition of epitopes), or to facilitate cleavage between epitopes and thereby enhance epitope presentation. Junctional epitopes are generally to be avoided because the recipient may generate an immune response to that non-native epitope. Of particular concern is a junctional epitope that is a "dominant epitope." A dominant epitope may lead to such a

25 zealous response that immune responses to other epitopes are diminished or suppressed.

Examples of polyepitopic vaccine compositions designed based on the above criteria can include epitopes from the core, S, E1, NS1/E2, NS2, NS3, NS4, and NS5 domains of the HCV polyprotein. These regions encompass the following amino acid sequences using numbering relative to the prototype HCV-1 strain (Genbank accession

30 number M62321; *see, e.g.*, US Patent Nos. 5,683,864 and 5,670,153): C domain (amino acids 1-120); S (amino acids 120-400); NS3 (amino acids 1050-1640); NS4 (amino acids 1640-2000); NS5 (amino acids 2000-3011); and envelop proteins, E1 and E2/NS1, encompassing amino acids 192-750. Amino acids 750 to 1050 are designated as domain X as applied to the present invention. As appreciated by one of ordinary skill in the art,

the designation of the amino acid range for each domain may diverge to some extent from that of HCV-1 depending on the strain of HCV. One of ordinary skill in the art, when looking at an HCV polyprotein sequence, would readily be able to determine the domain boundaries.

- 5           Specific embodiments of the polyepitopic compositions of the present invention include a pharmaceutical composition comprising a pharmaceutically acceptable carrier and combination of motif-bearing peptides that are immunologically cross-reactive with peptides of HCV-1, wherein at least one of the peptides bears a motif of Table Ia, and further wherein the combination of motif-bearing peptides consists of: a) one or more  
10 peptides comprising at least 8 amino acids from an HCV C domain; b) one or more peptides comprising at least 8 amino acids of a further domain selected from the group consisting of: an S domain, an NS3 domain, an NS4 domain, or an NS5 domain, and; c) optionally, one or more motif-bearing peptides from one or more additional HCV domains with a *proviso* that an additional domain is not a further domain listed in "b".
- 15 Preferably, such a pharmaceutical composition may additionally comprise one or more distinct HCV motif-bearing peptide(s) comprising at least 8 amino acids of an X domain or, alternatively, the composition may further comprise additional HCV motif-bearing peptide(s) that are from an envelope domain, the envelope domain peptide(s) consisting of one or more copies of a single HCV peptide comprising at least 8 amino acids of an  
20 envelope domain.

- In another embodiment, the polyepitopic pharmaceutical composition may comprise a pharmaceutically acceptable carrier and combination of motif-bearing peptides that are immunologically cross-reactive with HCV-1 peptides, the peptides from multiple domains of HCV, wherein at least one of the peptides bears a motif of Table Ia,  
25 and wherein the combination of motif-bearing peptides consists essentially of: a) one or more peptides comprising at least 8 amino acids from a C domain; and, b) one or more peptides comprising at least 8 amino acids from an S, NS3, NS4, or NS5 domain, and, one HCV peptide comprising at least 8 amino acids of an envelope domain. Such a composition may further comprise one or more HCV motif-bearing peptides comprising  
30 at least 8 amino acids of an X domain.

          Alternatively, a pharmaceutical composition of the invention may comprise: a) a pharmaceutically acceptable carrier; and, b) a combination of one or more motif-bearing peptides of at least 8 amino acids derived from one or more hepatitis C virus (HCV) domains, wherein said peptides are cross-reactive with peptides of HCV-1, with a *proviso*

that the combination does not include a peptide of at least 8 amino acids from an HCV C domain, and wherein at least one of the peptides bears a motif of Table Ia, said domains selected from the group consisting of: an S domain; an NS3 domain; an NS4 domain; an NS5 domain; and, an X domain. Such a composition may additionally comprise motif-bearing HCV envelope peptide(s) consisting of one or more copies of a single HCV peptide comprising at least 8 amino acids of an envelope domain.

Lastly, an embodiment of the invention may comprise a pharmaceutical composition comprising a pharmaceutically acceptable carrier and combination of two or more motif-bearing peptides from a single domain of an HCV-1 strain, said peptides immunologically cross-reactive with peptides of an HCV-1 antigen, wherein at least one of the peptides bears a motif of Table Ia, and the peptides are derived from HCV, and the HCV domain is selected from the group consisting of: a C domain; an S domain; an NS3 domain; an NS4 domain; an NS5 domain; an X domain; or, an envelope domain from a single HCV strain, with a *proviso* that the envelope domain is other than a variable envelope domain.

In the embodiments set forth, "peptides immunologically cross-reactive with HCV-1" refers to peptides that are bound by the same antibody; "derived from" refers to a fragment or subsequence and conservatively modified variants thereof.

#### IV.K.1. Minigene Vaccines

A number of different approaches are available which allow simultaneous delivery of multiple epitopes. Nucleic acids encoding the peptides of the invention are a particularly useful embodiment of the invention. Epitopes for inclusion in a minigene are preferably selected according to the guidelines set forth in the previous section. A preferred means of administering nucleic acids encoding the peptides of the invention uses minigene constructs encoding a peptide comprising one or multiple epitopes of the invention.

The use of multi-epitope minigenes is described below and in, *e.g.*, co-pending application U.S.S.N. 09/311,784; An, L. and Whitton, J. L., *J. Virol.* 71:2292, 1997; Thomson, S. A. *et al.*, *J. Immunol.* 157:822, 1996; Whitton, J. L. *et al.*, *J. Virol.* 67:348, 1993; Hanke, R. *et al.*, *Vaccine* 16:426, 1998. For example, a multi-epitope DNA plasmid encoding supermotif- and/or motif-bearing HCV epitopes derived from multiple regions of the HCV polyprotein sequence, the PADRE™ universal helper T cell epitope (or

multiple HTL epitopes from HCV), and an endoplasmic reticulum-translocating signal sequence can be engineered.

The immunogenicity of a multi-epitopic minigene can be tested in transgenic mice to evaluate the magnitude of CTL induction responses against the epitopes tested.

5 Further, the immunogenicity of DNA-encoded epitopes *in vivo* can be correlated with the *in vitro* responses of specific CTL lines against target cells transfected with the DNA plasmid. Thus, these experiments can show that the minigene serves to both: 1.) generate a CTL response and 2.) that the induced CTLs recognized cells expressing the encoded epitopes.

10 For example, to create a DNA sequence encoding the selected epitopes (minigene) for expression in human cells, the amino acid sequences of the epitopes may be reverse translated. A human codon usage table can be used to guide the codon choice for each amino acid. These epitope-encoding DNA sequences may be directly adjoined, so that when translated, a continuous polypeptide sequence is created. To optimize expression  
15 and/or immunogenicity, additional elements can be incorporated into the minigene design. Examples of amino acid sequences that can be reverse translated and included in the minigene sequence include: HLA class I epitopes, HLA class II epitopes, a ubiquitination signal sequence, and/or an endoplasmic reticulum targeting signal. In addition, HLA presentation of CTL and HTL epitopes may be improved by including  
20 synthetic (*e.g.* poly-alanine) or naturally-occurring flanking sequences adjacent to the CTL or HTL epitopes; these larger peptides comprising the epitope(s) are within the scope of the invention.

The minigene sequence may be converted to DNA by assembling oligonucleotides that encode the plus and minus strands of the minigene. Overlapping oligonucleotides  
25 (30-100 bases long) may be synthesized, phosphorylated, purified and annealed under appropriate conditions using well known techniques. The ends of the oligonucleotides can be joined, for example, using T4 DNA ligase. This synthetic minigene, encoding the epitope polypeptide, can then be cloned into a desired expression vector.

Standard regulatory sequences well known to those of skill in the art are  
30 preferably included in the vector to ensure expression in the target cells. Several vector elements are desirable: a promoter with a down-stream cloning site for minigene insertion; a polyadenylation signal for efficient transcription termination; an *E. coli* origin of replication; and an *E. coli* selectable marker (*e.g.* ampicillin or kanamycin resistance). Numerous promoters can be used for this purpose, *e.g.*, the human cytomegalovirus

(hCMV) promoter. See, *e.g.*, U.S. Patent Nos. 5,580,859 and 5,589,466 for other suitable promoter sequences.

Additional vector modifications may be desired to optimize minigene expression and immunogenicity. In some cases, introns are required for efficient gene expression, and one or more synthetic or naturally-occurring introns could be incorporated into the transcribed region of the minigene. The inclusion of mRNA stabilization sequences and sequences for replication in mammalian cells may also be considered for increasing minigene expression.

Once an expression vector is selected, the minigene is cloned into the polylinker region downstream of the promoter. This plasmid is transformed into an appropriate *E. coli* strain, and DNA is prepared using standard techniques. The orientation and DNA sequence of the minigene, as well as all other elements included in the vector, are confirmed using restriction mapping and DNA sequence analysis. Bacterial cells harboring the correct plasmid can be stored as a master cell bank and a working cell bank.

In addition, immunostimulatory sequences (ISSs or CpGs) appear to play a role in the immunogenicity of DNA vaccines. These sequences may be included in the vector, outside the minigene coding sequence, if desired to enhance immunogenicity.

In some embodiments, a bi-cistronic expression vector which allows production of both the minigene-encoded epitopes and a second protein (included to enhance or decrease immunogenicity) can be used. Examples of proteins or polypeptides that could beneficially enhance the immune response if co-expressed include cytokines (*e.g.*, IL-2, IL-12, GM-CSF), cytokine-inducing molecules (*e.g.*, LeIF), costimulatory molecules, or for HTL responses, pan-DR binding proteins (PADRE™, Epimmune, San Diego, CA). Helper (HTL) epitopes can be joined to intracellular targeting signals and expressed separately from expressed CTL epitopes; this allows direction of the HTL epitopes to a cell compartment different than that of the CTL epitopes. If required, this could facilitate more efficient entry of HTL epitopes into the HLA class II pathway, thereby improving HTL induction. In contrast to HTL or CTL induction, specifically decreasing the immune response by co-expression of immunosuppressive molecules (*e.g.* TGF- $\beta$ ) may be beneficial in certain diseases.

Therapeutic quantities of plasmid DNA can be produced for example, by fermentation in *E. coli*, followed by purification. Aliquots from the working cell bank are used to inoculate growth medium, and grown to saturation in shaker flasks or a bioreactor

according to well known techniques. Plasmid DNA can be purified using standard bioseparation technologies such as solid phase anion-exchange resins supplied by QIAGEN, Inc. (Valencia, California). If required, supercoiled DNA can be isolated from the open circular and linear forms using gel electrophoresis or other methods.

5 Purified plasmid DNA can be prepared for injection using a variety of formulations. The simplest of these is reconstitution of lyophilized DNA in sterile phosphate-buffer saline (PBS). This approach, known as "naked DNA," is currently being used for intramuscular (IM) administration in clinical trials. To maximize the immunotherapeutic effects of minigene DNA vaccines, an alternative method for  
10 formulating purified plasmid DNA may be desirable. A variety of methods have been described, and new techniques may become available. Cationic lipids can also be used in the formulation (see, *e.g.*, as described by WO 93/24640; Mannino & Gould-Fogerite, *BioTechniques* 6(7): 682 (1988); U.S. Pat No. 5,279,833; WO 91/06309; and Felgner, *et al.*, *Proc. Nat'l Acad. Sci. USA* 84:7413 (1987). In addition, glycolipids, fusogenic  
15 liposomes, peptides and compounds referred to collectively as protective, interactive, non-condensing compounds (PINC) could also be complexed to purified plasmid DNA to influence variables such as stability, intramuscular dispersion, or trafficking to specific organs or cell types.

Target cell sensitization can be used as a functional assay for expression and HLA  
20 class I presentation of minigene-encoded CTL epitopes. For example, the plasmid DNA is introduced into a mammalian cell line that is suitable as a target for standard CTL chromium release assays. The transfection method used will be dependent on the final formulation. Electroporation can be used for "naked" DNA, whereas cationic lipids allow direct *in vitro* transfection. A plasmid expressing green fluorescent protein (GFP) can be  
25 co-transfected to allow enrichment of transfected cells using fluorescence activated cell sorting (FACS). These cells are then chromium-51 ( $^{51}\text{Cr}$ ) labeled and used as target cells for epitope-specific CTL lines; cytotoxicity, detected by  $^{51}\text{Cr}$  release, indicates both production of, and HLA presentation of, minigene-encoded CTL epitopes. Expression of HTL epitopes may be evaluated in an analogous manner using assays to assess HTL  
30 activity.

*In vivo* immunogenicity is a second approach for functional testing of minigene DNA formulations. Transgenic mice expressing appropriate human HLA proteins are immunized with the DNA product. The dose and route of administration are formulation dependent (*e.g.*, IM for DNA in PBS, intraperitoneal (IP) for lipid-complexed DNA).



Twenty-one days after immunization, splenocytes are harvested and restimulated for 1 week in the presence of peptides encoding each epitope being tested. Thereafter, for CTL effector cells, assays are conducted for cytolysis of peptide-loaded, <sup>51</sup>Cr-labeled target cells using standard techniques. Lysis of target cells that were sensitized by HLA loaded  
5 with peptide epitopes, corresponding to minigene-encoded epitopes, demonstrates DNA vaccine function for *in vivo* induction of CTLs. Immunogenicity of HTL epitopes is evaluated in transgenic mice in an analogous manner.

Alternatively, the nucleic acids can be administered using ballistic delivery as described, for instance, in U.S. Patent No. 5,204,253. Using this technique, particles  
10 comprised solely of DNA are administered. In a further alternative embodiment, DNA can be adhered to particles, such as gold particles.

#### IV.K.2. Combinations of CTL Peptides with Helper Peptides

Vaccine compositions comprising the peptides of the present invention, or analogs  
15 thereof, which have immunostimulatory activity may be modified to provide desired attributes, such as improved serum half life, or to enhance immunogenicity.

For instance, the ability of the peptides to induce CTL activity can be enhanced by linking the peptide to a sequence which contains at least one epitope that is capable of inducing a T helper cell response. The use of T helper epitopes in conjunction with CTL  
20 epitopes to enhance immunogenicity is illustrated, for example, in co-pending U.S.S.N. 08/820360, U.S.S.N. 08/197,484, and U.S.S.N. 08/464,234.

Particularly preferred CTL epitope/HTL epitope conjugates are linked by a spacer molecule. The spacer is typically comprised of relatively small, neutral molecules, such as amino acids or amino acid mimetics, which are substantially uncharged under  
25 physiological conditions. The spacers are typically selected from, *e.g.*, Ala, Gly, or other neutral spacers of nonpolar amino acids or neutral polar amino acids. It will be understood that the optionally present spacer need not be comprised of the same residues and thus may be a hetero- or homo-oligomer. When present, the spacer will usually be at least one or two residues, more usually three to six residues. Alternatively, the CTL  
30 peptide may be linked to the T helper peptide without a spacer.

Although the CTL peptide epitope can be linked directly to the T helper peptide epitope, often CTL epitope/HTL epitope conjugates are linked by a spacer molecule. The spacer is typically comprised of relatively small, neutral molecules, such as amino acids or amino acid mimetics, which are substantially uncharged under physiological

conditions. The spacers are typically selected from, *e.g.*, Ala, Gly, or other neutral spacers of nonpolar amino acids or neutral polar amino acids. It will be understood that the optionally present spacer need not be comprised of the same residues and thus may be a hetero- or homo-oligomer. When present, the spacer will usually be at least one or two  
5 residues, more usually three to six residues. The CTL peptide epitope can be linked to the T helper peptide epitope either directly or via a spacer either at the amino or carboxy terminus of the CTL peptide. The amino terminus of either the immunogenic peptide or the T helper peptide may be acylated.

HTL peptide epitopes can also be modified to alter their biological properties. For  
10 example, peptides comprising HTL epitopes can contain D-amino acids to increase their resistance to proteases and thus extend their serum half-life. Also, the epitope peptides of the invention can be conjugated to other molecules such as lipids, proteins or sugars, or any other synthetic compounds, to increase their biological activity. Specifically, the T helper peptide can be conjugated to one or more palmitic acid chains at either the amino  
15 or carboxyl termini.

In certain embodiments, the T helper peptide is one that is recognized by T helper cells present in the majority of the population. This can be accomplished by selecting amino acid sequences that bind to many, most, or all of the HLA class II molecules. These are known as "loosely HLA-restricted" or "promiscuous" T helper sequences.  
20 Examples of amino acid sequences that are promiscuous include sequences from antigens such as tetanus toxoid at positions 830-843 (QYIKANSKFIGITE), *Plasmodium falciparum* CS protein at positions 378-398 (DIEKKIAKMEKASSVFNVNS), and Streptococcus 18kD protein at positions 116 (GAVDSILGGVATYGAA). Other examples include peptides bearing a DR 1-4-7 supermotif, or either of the DR3 motifs.

25 Alternatively, it is possible to prepare synthetic peptides capable of stimulating T helper lymphocytes, in a loosely HLA-restricted fashion, using amino acid sequences not found in nature (*see, e.g.*, PCT publication WO 95/07707). These synthetic compounds called Pan-DR-binding epitopes (*e.g.*, PADRE™, Epimmune, Inc., San Diego, CA) are designed to most preferably bind most HLA-DR (human HLA class II) molecules. For  
30 instance, a pan-DR-binding epitope peptide having the formula: aKXVWANTLKAAa, where "X" is either cyclohexylalanine, phenylalanine, or tyrosine, and a is either D-alanine or L-alanine, has been found to bind to most HLA-DR alleles, and to stimulate the response of T helper lymphocytes from most individuals, regardless of their HLA type.

An alternative of a pan-DR binding epitope comprises all "L" natural amino acids and can be provided in the form of nucleic acids that encode the epitope.

In some embodiments it may be desirable to include in the pharmaceutical compositions of the invention at least one component which primes cytotoxic T lymphocytes. Lipids have been identified as agents capable of priming CTL *in vivo* against viral antigens. For example, palmitic acid residues can be attached to the  $\epsilon$ - and  $\alpha$ -amino groups of a lysine residue and then linked, *e.g.*, via one or more linking residues such as Gly, Gly-Gly-, Ser, Ser-Ser, or the like, to an immunogenic peptide. The lipidated peptide can then be administered either directly in a micelle or particle, incorporated into a liposome, or emulsified in an adjuvant, *e.g.*, incomplete Freund's adjuvant. In a preferred embodiment, a particularly effective immunogenic comprises palmitic acid attached to  $\epsilon$ - and  $\alpha$ - amino groups of Lys, which is attached via linkage, *e.g.*, Ser-Ser, to the amino terminus of the immunogenic peptide.

As another example of lipid priming of CTL responses, *E. coli* lipoproteins, such as tripalmitoyl-S-glycerylcysteinylserine (P<sub>3</sub>CSS) can be used to prime virus specific CTL when covalently attached to an appropriate peptide. (*See, e.g., Deres, et al., Nature* 342:561, 1989). Peptides of the invention can be coupled to P<sub>3</sub>CSS, for example, and the lipopeptide administered to an individual to specifically prime a CTL response to the target antigen. Moreover, because the induction of neutralizing antibodies can also be primed with P<sub>3</sub>CSS-conjugated epitopes, two such compositions can be combined to more effectively elicit both humoral and cell-mediated responses to infection.

As noted herein, additional amino acids can be added to the termini of a peptide to provide for ease of linking peptides one to another, for coupling to a carrier support or larger peptide, for modifying the physical or chemical properties of the peptide or oligopeptide, or the like. Amino acids such as tyrosine, cysteine, lysine, glutamic or aspartic acid, or the like, can be introduced at the C- or N-terminus of the peptide or oligopeptide, particularly class I peptides. However, it is to be noted that modification at the carboxyl terminus of a CTL epitope may, in some cases, alter binding characteristics of the peptide. In addition, the peptide or oligopeptide sequences can differ from the natural sequence by being modified by terminal-NH<sub>2</sub> acylation, *e.g.*, by alkanoyl (C<sub>1</sub>-C<sub>20</sub>) or thioglycolyl acetylation, terminal-carboxyl amidation, *e.g.*, ammonia, methylamine, *etc.* In some instances these modifications may provide sites for linking to a support or other molecule.

*Vaccine Compositions Comprising Dendritic Cells Pulsed with CTL and/or HTL Peptides*

An embodiment of a vaccine composition in accordance with the invention comprises *ex vivo* administration of a cocktail of epitope-bearing peptides to PBMC, or  
5 isolated DC therefrom, from the patient's blood. A pharmaceutical to facilitate harvesting of DC can be used, such as GM-CSF/IL-4. After pulsing the DC with peptides and prior to reinfusion into patients, the DC are washed to remove unbound peptides. In this embodiment, a vaccine comprises peptide-pulsed DCs which present the pulsed peptide epitopes complexed with HLA molecules on their surfaces. The vaccine is then  
10 administered to the patient.

**IV.L. Administration of Vaccines for Therapeutic or Prophylactic Purposes**

The peptides of the present invention and pharmaceutical and vaccine compositions of the invention are useful for administration to mammals, particularly  
15 humans, to treat and/or prevent HCV infection. Vaccine compositions containing the peptides of the invention are administered to a patient infected with HCV or to an individual susceptible to, or otherwise at risk for, HCV infection to elicit an immune response against HCV antigens and thus enhance the patient's own immune response capabilities. In therapeutic applications, peptide and/or nucleic acid compositions are  
20 administered to a patient in an amount sufficient to elicit an effective CTL and/or HTL response to the virus antigen and to cure or at least partially arrest or slow symptoms and/or complications. An amount adequate to accomplish this is defined as "therapeutically effective dose." Amounts effective for this use will depend on, *e.g.*, the particular composition administered, the manner of administration, the stage and severity  
25 of the disease being treated, the weight and general state of health of the patient, and the judgment of the prescribing physician.

The vaccine compositions of the invention may also be used purely as prophylactic agents. Generally the dosage for an initial prophylactic immunization generally occurs in a unit dosage range where the lower value is about 1, 5, 50, 500, or  
30 1000 µg and the higher value is about 10,000; 20,000; 30,000; or 50,000 µg. Dosage values for a human typically range from about 500 µg to about 50,000 µg per 70 kilogram patient. This is followed by boosting dosages of between about 1.0 µg to about 50,000 µg of peptide administered at defined intervals from about four weeks to six months after the

initial administration of vaccine. The immunogenicity of the vaccine may be assessed by measuring the specific activity of CTL and HTL obtained from a sample of the patient's blood.

As noted above, peptides comprising CTL and/or HTL epitopes of the invention  
5 induce immune responses when presented by HLA molecules and contacted with a CTL  
or HTL specific for an epitope comprised by the peptide. The manner in which the  
peptide is contacted with the CTL or HTL is not critical to the invention. For instance,  
the peptide can be contacted with the CTL or HTL either *in vivo* or *in vitro*. If the  
contacting occurs *in vivo*, the peptide itself can be administered to the patient, or other  
10 vehicles, *e.g.*, DNA vectors encoding one or more peptides, viral vectors encoding the  
peptide(s), liposomes and the like, can be used, as described herein. When the peptide is  
contacted *in vitro*, the vaccinating agent can comprise a population of cells, *e.g.*, peptide-  
pulsed dendritic cells, or TAA-specific CTLs, which have been induced by pulsing  
antigen-presenting cells *in vitro* with the peptide. Such a cell population is subsequently  
15 administered to a patient in a therapeutically effective dose.

The peptides or DNA encoding them can be administered individually or as  
fusions of one or more peptide sequences.

For pharmaceutical compositions, the immunogenic peptides of the invention, or  
DNA encoding them, are generally administered to an individual already infected with  
20 HCV. The peptides or DNA encoding them can be administered individually or as  
fusions of one or more peptide sequences. Those in the incubation phase or the acute  
phase of infection can be treated with the immunogenic peptides separately or in  
conjunction with other treatments, as appropriate.

For therapeutic use, administration should generally begin at the first diagnosis of  
25 HCV infection. This is followed by boosting doses until at least symptoms are  
substantially abated and for a period thereafter. In chronic infection, loading doses  
followed by boosting doses may be required.

Treatment of an infected individual with the compositions of the invention may  
hasten resolution of the infection in acutely infected individuals. For those individuals  
30 susceptible (or predisposed) to developing chronic infection, the compositions are  
particularly useful in methods for preventing the evolution from acute to chronic  
infection. Where susceptible individuals are identified prior to or during infection, the  
composition can be targeted to them, thus minimizing the need for administration to a  
larger population.

The peptide or other compositions used for the treatment or prophylaxis of HCV infection can be used, *e.g.*, in persons who have not manifested symptoms of disease but who act as a disease vector. In this context, it is generally important to provide an amount of the peptide epitope delivered by a mode of administration sufficient to  
5 effectively stimulate a cytotoxic T cell response; compositions which stimulate helper T cell responses can also be given in accordance with this embodiment of the invention.

The dosage for an initial therapeutic immunization generally occurs in a unit dosage range where the lower value is about 1, 5, 50, 500, or 1000  $\mu\text{g}$  and the higher value is about 10,000; 20,000; 30,000; or 50,000  $\mu\text{g}$ . Dosage values for a human  
10 typically range from about 500  $\mu\text{g}$  to about 50,000  $\mu\text{g}$  per 70 kilogram patient. Boosting dosages of between about 1.0  $\mu\text{g}$  to about 50000  $\mu\text{g}$  of peptide pursuant to a boosting regimen over weeks to months may be administered depending upon the patient's response and condition as determined by measuring the specific activity of CTL and HTL obtained from the patient's blood. The peptides and compositions of the present  
15 invention may be employed in serious disease states, that is, life-threatening or potentially life threatening situations. In such cases, as a result of the minimal amounts of extraneous substances and the relative nontoxic nature of the peptides in preferred compositions of the invention, it is possible and may be felt desirable by the treating physician to administer substantial excesses of these peptide compositions relative to  
20 these stated dosage amounts.

Thus, for treatment of chronic infection, a representative dose is in the range disclosed above, namely where the lower value is about 1, 5, 50, 500, or 1000  $\mu\text{g}$  and the higher value is about 10,000; 20,000; 30,000; or 50,000  $\mu\text{g}$ , preferably from about 500  $\mu\text{g}$  to about 50,000  $\mu\text{g}$  per 70 kilogram patient. Initial doses followed by boosting doses at  
25 established intervals, *e.g.*, from four weeks to six months, may be required, possibly for a prolonged period of time to effectively immunize an individual. In the case of chronic infection, administration should continue until at least clinical symptoms or laboratory tests indicate that the viral infection has been eliminated or substantially abated and for a period thereafter. The dosages, routes of administration, and dose schedules are adjusted  
30 in accordance with methodologies known in the art.

The pharmaceutical compositions for therapeutic treatment are intended for parenteral, topical, oral, intrathecal, or local administration. Preferably, the pharmaceutical compositions are administered parentally, *e.g.*, intravenously,

subcutaneously, intradermally, or intramuscularly. Thus, the invention provides compositions for parenteral administration which comprise a solution of the immunogenic peptides dissolved or suspended in an acceptable carrier, preferably an aqueous carrier. A variety of aqueous carriers may be used, *e.g.*, water, buffered water, 0.8% saline, 0.3% glycine, hyaluronic acid and the like. These compositions may be sterilized by conventional, well known sterilization techniques, or may be sterile filtered. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile solution prior to administration. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH-adjusting and buffering agents, tonicity adjusting agents, wetting agents, preservatives, and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, *etc.*

The concentration of peptides of the invention in the pharmaceutical formulations can vary widely, *i.e.*, from less than about 0.1%, usually at or at least about 2% to as much as 20% to 50% or more by weight, and will be selected primarily by fluid volumes, viscosities, *etc.*, in accordance with the particular mode of administration selected.

A human unit dose form of the peptide composition is typically included in a pharmaceutical composition that comprises a human unit dose of an acceptable carrier, preferably an aqueous carrier, and is administered in a volume of fluid that is known by those of skill in the art to be used for administration of such compositions to humans (*see, e.g., Remington's Pharmaceutical Sciences*, 17<sup>th</sup> Edition, A. Gennaro, Editor, Mack Publishing Co., Easton, Pennsylvania, 1985).

The peptides of the invention may also be administered via liposomes, which serve to target the peptides to a particular tissue, such as lymphoid tissue, or to target selectively to infected cells, as well as to increase the half-life of the peptide composition. Liposomes include emulsions, foams, micelles, insoluble monolayers, liquid crystals, phospholipid dispersions, lamellar layers and the like. In these preparations, the peptide to be delivered is incorporated as part of a liposome, alone or in conjunction with a molecule which binds to a receptor prevalent among lymphoid cells, such as monoclonal antibodies which bind to the CD45 antigen, or with other therapeutic or immunogenic compositions. Thus, liposomes either filled or decorated with a desired peptide of the invention can be directed to the site of lymphoid cells, where the liposomes then deliver the peptide compositions. Liposomes for use in accordance with the invention are formed

from standard vesicle-forming lipids, which generally include neutral and negatively charged phospholipids and a sterol, such as cholesterol. The selection of lipids is generally guided by consideration of, *e.g.*, liposome size, acid lability and stability of the liposomes in the blood stream. A variety of methods are available for preparing  
5 liposomes, as described in, *e.g.*, Szoka, *et al.*, *Ann. Rev. Biophys. Bioeng.* 9:467 (1980), and U.S. Patent Nos. 4,235,871, 4,501,728, 4,837,028, and 5,019,369.

For targeting cells of the immune system, a ligand to be incorporated into the liposome can include, *e.g.*, antibodies or fragments thereof specific for cell surface determinants of the desired immune system cells. A liposome suspension containing a  
10 peptide may be administered intravenously, locally, topically, *etc.* in a dose which varies according to, *inter alia*, the manner of administration, the peptide being delivered, and the stage of the disease being treated.

For solid compositions, conventional nontoxic solid carriers may be used which include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium  
15 stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like. For oral administration, a pharmaceutically acceptable nontoxic composition is formed by incorporating any of the normally employed excipients, such as those carriers previously listed, and generally 10-95% of active ingredient, that is, one or more peptides of the invention, and more preferably at a concentration of 25%-75%.

For aerosol administration, the immunogenic peptides are preferably supplied in finely divided form along with a surfactant and propellant. Typical percentages of peptides are 0.01%-20% by weight, preferably 1%-10%. The surfactant must, of course, be nontoxic, and preferably soluble in the propellant. Representative of such agents are the esters or partial esters of fatty acids containing from 6 to 22 carbon atoms, such as  
25 caproic, octanoic, lauric, palmitic, stearic, linoleic, linolenic, olesteric and oleic acids with an aliphatic polyhydric alcohol or its cyclic anhydride. Mixed esters, such as mixed or natural glycerides may be employed. The surfactant may constitute 0.1%-20% by weight of the composition, preferably 0.25-5%. The balance of the composition is ordinarily propellant. A carrier can also be included, as desired, as with, *e.g.*, lecithin for intranasal  
30 delivery.

#### IV.M. Kits

The peptide and nucleic acid compositions of this invention can be provided in kit form together with instructions for vaccine administration. Typically the kit would



include desired peptide compositions in a container, preferably in unit dosage form and instructions for administration. An alternative kit would include a minigene construct with desired nucleic acids of the invention in a container, preferably in unit dosage form together with instructions for administration. Lymphokines such as IL-2 or IL-12 may  
5 also be included in the kit. Other kit components that may also be desirable include, for example, a sterile syringe, booster dosages, and other desired excipients.

The invention will be described in greater detail by way of specific examples. The following examples are offered for illustrative purposes, and are not intended to limit the invention in any manner. Those of skill in the art will readily recognize a variety of  
10 non-critical parameters that can be changed or modified to yield alternative embodiments in accordance with the invention.

## V. EXAMPLES

As in many viral diseases, there is evidence that clearance of HCV is mediated by  
15 CTL. In a study of primary HCV infection in six chimpanzees, four progressed to chronic infection (Cooper *et al.*, abstract, 19<sup>th</sup> US-Japan Hepatitis Joint Panel Meeting, January 27-29, 1998). It was found that these four animals showed either no CTL response or a very narrowly focused response during early infection. In contrast, in the remaining two animals that resolved the infection, a broad CTL response was observed  
20 against multiple HCV proteins, some of which were conserved. Weiner *et al.* (*Proc. Natl. Acad. Sci. USA* 92:2755-2759, 1995) demonstrated that viral escape, in which the epitopes presented to PATR class I molecules mutated, was linked with a progression toward chronic infection. These data show a role for the CTL in directing the course of HCV disease, and in shaping the genetic composition of HCV species in the persistently  
25 infected host.

In work in humans, Koziel and co-workers have established the presence of HCV-specific CTL in liver infiltrates from patients with chronic HCV infection (Koziel *et al.*, *J. Immunol.* 149:3339, 1992; and Koziel *et al.*, *J. Virol.* 67:7522, 1993), and have also identified a number of CTL epitopes recognized in the context of several different HLA  
30 class I molecules. Other investigators have shown that HCV-specific CTL can be detected in the peripheral blood of patients with chronic hepatitis C (Cerny *et al.*, *J. Clin. Invest.* 95:521, 1995; Cerny *et al.*, *Curr. Topics in Micro. and Immunol* 189:169, 1994; Cerny *et al.*, Abst. 2<sup>nd</sup> International Meeting on Hepatitis C and Related Viruses; La Jolla, CA, 1994; Battegay *et al.*, Abst. 2<sup>nd</sup> International Meeting on Hepatitis C and Related

Viruses; La Jolla, CA, 1994; Shirai *et al.*, *J. Virol.* 68:3334, 1994; Shirai *et al.*, *J. Immunol.* 154:2733, 1995; Battegay *et al.*, *J. Virol.* 69:2462, 1995). In addition, escape variants have been demonstrated in patients chronically infected with HCV (Chang *et al.*, *J. Clin. Invest.* 100:2376-2385, 1997; Tsai *et al.*, *Gastroenterology* 115:954-966, 1998).

5       The magnitude of the CTL responses observed in HCV-infected patients is, in general, higher than those observed in the case of chronic hepatitis B infection, suggesting that there is less impairment of specific T cell immunity than with HBV infection. The magnitude of CTL responses in HCV patients is, however, lower than those observed in HBV infected individuals who successfully cleared HBV infection.

10       These results support the understanding that HCV infected patients are capable of responding to active immunotherapy, and suggest that potentiation and increasing of T cell responses to HCV may be of use in therapy and prevention of chronic HCV infection (Prince, A. M. *FEMS Micro. Rev.* 14:273, 1994).

      Several groups have analyzed the potential role of HCV-specific CTL responses

15       in disease resistance and pathogenesis. In some studies no correlation was found between CTL viremia and CTL precursor frequency for individual HCV epitopes (Rehermann *et al.*, *J. Clin. Invest.* 98:1432-1440, 1996; Wong *et al.*, *J. Immunol.* 160:1479-1488, 1998). In other studies, however, it was shown that a clear correlation existed between levels of HCV infection and CTL responses, provided that the global response against multiple

20       CTL epitopes was considered (Rehermann *et al.*, *J. Virol.* 70:7092-7102, 1996). These data represent a strong rationale for development of vaccine constructs capable of inducing vigorous CTL responses directed against a multiplicity of conserved HCV-derived epitopes.

      Koziel and colleagues have demonstrated the presence of HCV-specific CTLs, as

25       well as T helper cell responses, in exposed but seronegative individuals (Koziel *et al.*, *J. Infect. Diseases* 176:859-866, 1997). In addition, HCV-specific CTLs have been detected in healthy, seronegative family members of chronically HCV-infected patients, indicating that a protective immunity is established in absence of a detectable infection (Bronowicki *et al.*, *J. Infect. Dis.* 176:518-522, 1997; Scognamiglio *et al.*, in preparation).

30       Experimental evidence also indicates that HTL epitopes play an important role in immune reactivity and defenses against HCV infection (Missale *et al.*, *J. Clin. Invest.* 98:706-714, 1996). Diepolder *et al.* (in *Lancet* 346:1006, 1995) have shown that a region of the NS3 gene (NS3 1007-1534) is recognized by patients who clear acute HCV infection, but is not seen by patients who develop chronic infection. Subsequent studies

showed that this particular region contain a highly cross-reactive HTL epitope (NS3 1248-1261), which binds with good affinity to 10 of 13 DR molecules tested, and is highly conserved in 30/33 different HCV isolates considered (Diepolder *et al.*, *J. Virol.* 71:6011-6019, 1997). These data suggested that directing HTL responses to this type of epitope (rather than to less cross-reactive and/or highly variable ones) will be of therapeutic and prophylactic benefit and strongly argue for inclusion of this and other epitopes with similar characteristics in HCV vaccine constructs.

The following examples illustrate identification, selection, and use of immunogenic Class I and Class II peptide epitopes for inclusion in vaccine compositions.

#### Example 1: HLA Class I and Class II Binding Assays

The following example of peptide binding to HLA molecules demonstrates quantification of binding affinities of HLA class I and class II peptides. Binding assays can be performed with peptides that are either motif-bearing or not motif-bearing.

Epstein-Barr virus (EBV)-transformed homozygous cell lines, fibroblasts, CIR, or 721.22 transfectants were used as sources of HLA class I molecules. The specific cell lines routinely used for purification of MHC class I and class II molecules are listed in Table XXIV. Cell lysates were prepared and HLA molecules purified in accordance with disclosed protocols (Sidney *et al.*, *Current Protocols in Immunology* 18.3.1 (1998); Sidney, *et al.*, *J. Immunol.* 154:247 (1995); Sette, *et al.*, *Mol. Immunol.* 31:813 (1994)). HLA molecules were purified from lysates by affinity chromatography. The lysate was passed over a column of Sepharose CL-4B beads coupled to an appropriate antibody. The antibodies used for the extraction of HLA from cell lysates are listed in Table XXV. The anti-HLA column was then washed with 10mM Tris-HCL, pH 8.0, in 1% NP-40, PBS, and PBS containing 0.4% n-octylglucoside and HLA molecules were eluted with 50mM diethylamine in 0.15M NaCl containing 0.4% n-octylglucoside, pH 11.5. A 1/25 volume of 2.0M Tris, pH 6.8, was added to the eluate to reduce the pH to ~8.0. Eluates were then be concentrated by centrifugation in Centriprep 30 concentrators (Amicon, Beverly, MA). Protein content was evaluated by a BCA protein assay (Pierce Chemical Co., Rockford, IL) and confirmed by SDS-PAGE.

A detailed description of the protocol utilized to measure the binding of peptides to Class I and Class II MHC has been published (Sette *et al.*, *Mol. Immunol.* 31:813, 1994; Sidney *et al.*, in *Current Protocols in Immunology*, Margulies, Ed., John Wiley & Sons, New York, Section 18.3, 1998). Briefly, purified MHC molecules (5 to 500nM)

were incubated with various unlabeled peptide inhibitors and 1-10nM  $^{125}\text{I}$ -radiolabeled probe peptides for 48h in PBS containing 0.05% Nonidet P-40 (NP40) (or 20% w/v digitonin for H-2 IA assays) in the presence of a protease inhibitor cocktail. All assays were at pH 7.0 with the exception of DRB1\*0301, which was performed at pH 4.5, and

5 DRB1\*1601 (DR2w2 $\beta_1$ ) and DRB4\*0101 (DRw53), which were performed at pH 5.0.

Following incubation, MHC-peptide complexes were separated from free peptide by gel filtration on 7.8 mm x 15 cm TSK200 columns (TosoHaas 16215, Montgomeryville, PA). Because the large size of the radiolabeled peptide used for the DRB1\*1501 (DR2w2 $\beta_1$ ) assay makes separation of bound from unbound peaks more

10 difficult under these conditions, all DRB1\*1501 (DR2w2 $\beta_1$ ) assays were performed using a 7.8mm x 30cm TSK2000 column eluted at 0.6 mls/min. The eluate from the TSK columns was passed through a Beckman 170 radioisotope detector, and radioactivity was plotted and integrated using a Hewlett-Packard 3396A integrator, and the fraction of peptide bound was determined.

15 Radiolabeled peptides were iodinated using the chloramine-T method. Representative radiolabeled probe peptides utilized in each assay, and its assay specific  $\text{IC}_{50}$  nM, are summarized in Tables IV and V. Typically, in preliminary experiments, each MHC preparation was titrated in the presence of fixed amounts of radiolabeled peptides to determine the concentration of HLA molecules necessary to bind 10-20% of

20 the total radioactivity. All subsequent inhibition and direct binding assays were performed using these HLA concentrations.

Since under these conditions  $[\text{label}] < [\text{HLA}]$  and  $\text{IC}_{50} \geq [\text{HLA}]$ , the measured  $\text{IC}_{50}$  values are reasonable approximations of the true  $K_D$  values. Peptide inhibitors are typically tested at concentrations ranging from 120  $\mu\text{g/ml}$  to 1.2  $\text{ng/ml}$ , and are tested in

25 two to four completely independent experiments. To allow comparison of the data obtained in different experiments, a relative binding figure is calculated for each peptide by dividing the  $\text{IC}_{50}$  of a positive control for inhibition by the  $\text{IC}_{50}$  for each tested peptide (typically unlabeled versions of the radiolabeled probe peptide). For database purposes, and inter-experiment comparisons, relative binding values are compiled. These values

30 can subsequently be converted back into  $\text{IC}_{50}$  nM values by dividing the  $\text{IC}_{50}$  nM of the positive controls for inhibition by the relative binding of the peptide of interest. This method of data compilation has proven to be the most accurate and consistent for

comparing peptides that have been tested on different days, or with different lots of purified MHC.

Because the antibody used for HLA-DR purification (LB3.1) is  $\alpha$ -chain specific,  $\beta_1$  molecules are not separated from  $\beta_3$  (and/or  $\beta_4$  and  $\beta_5$ ) molecules. The  $\beta_1$  specificity of the binding assay is obvious in the cases of DRB1\*0101 (DR1), DRB1\*0802 (DR8w2), and DRB1\*0803 (DR8w3), where no  $\beta_3$  is expressed. It has also been demonstrated for DRB1\*0301 (DR3) and DRB3\*0101 (DR52a), DRB1\*0401 (DR4w4), DRB1\*0404 (DR4w14), DRB1\*0405 (DR4w15), DRB1\*1101 (DR5), DRB1\*1201 (DR5w12), DRB1\*1302 (DR6w19) and DRB1\*0701 (DR7). The problem of  $\beta$  chain specificity for DRB1\*1501 (DR2w2 $\beta_1$ ), DRB5\*0101 (DR2w2 $\beta_2$ ), DRB1\*1601 (DR2w21 $\beta_1$ ), DRB5\*0201 (DR51Dw21), and DRB4\*0101 (DRw53) assays is circumvented by the use of fibroblasts. Development and validation of assays with regard to DR $\beta$  molecule specificity have been described previously (*see, e.g., Southwood et al., J. Immunol.* 160:3363-3373, 1998).

Binding assays as outlined above may be used to analyze supermotif and/or motif-bearing epitopes as, for example, described in Example 2.

Example 2. Identification of Conserved HLA Supermotif- and Motif-Bearing CTL Candidate Epitopes

Vaccine compositions of the invention may include multiple epitopes that comprise multiple HLA supermotifs or motifs to achieve broad population coverage. This example illustrates the identification of supermotif- and motif-bearing epitopes for the inclusion in such a vaccine composition. Calculation of population coverage was performed using the strategy described below.

*Computer searches and algorithms for identification of supermotif and/or motif-bearing epitopes*

Computer searches for epitopes bearing HLA Class I or Class II supermotifs or motifs were performed as follows. All translated HCV isolate sequences were analyzed using a text string search software program, *e.g., MotifSearch 1.4* (D. Brown, San Diego) to identify potential peptide sequences containing appropriate HLA binding motifs; alternative programs are readily produced in accordance with information in the art in view of the motif/supermotif disclosure herein. Furthermore, such calculations can be

made mentally. Identified A2-, A3-, and DR-supermotif sequences were scored using polynomial algorithms to predict their capacity to bind to specific HLA-Class I or Class II molecules. These polynomial algorithms take into account both extended and refined motifs (that is, to account for the impact of different amino acids at different positions),  
5 and are essentially based on the premise that the overall affinity (or  $\Delta G$ ) of peptide-HLA molecule interactions can be approximated as a linear polynomial function of the type:

$$"\Delta G" = a_{1i} \times a_{2i} \times a_{3i} \dots \times a_{ni}$$

where  $a_{ji}$  is a coefficient which represents the effect of the presence of a given amino acid ( $j$ ) at a given position ( $i$ ) along the sequence of a peptide of  $n$  amino acids. The crucial  
10 assumption of this method is that the effects at each position are essentially independent of each other (i.e., independent binding of individual side-chains). When residue  $j$  occurs at position  $i$  in the peptide, it is assumed to contribute a constant amount  $j_i$  to the free energy of binding of the peptide irrespective of the sequence of the rest of the peptide. This assumption is justified by studies from our laboratories that demonstrated that  
15 peptides are bound to MHC and recognized by T cells in essentially an extended conformation (data omitted herein).

The method of derivation of specific algorithm coefficients has been described in Gulukota *et al.*, *J. Mol. Biol.* 267:1258-126, 1997; (see also Sidney *et al.*, *Human Immunol.* 45:79-93, 1996; and Southwood *et al.*, *J. Immunol.* 160:3363-3373, 1998).  
20 Briefly, for all  $i$  positions, anchor and non-anchor alike, the geometric mean of the average relative binding (ARB) of all peptides carrying  $j$  is calculated relative to the remainder of the group, and used as the estimate of  $j_i$ . For Class II peptides, if multiple alignments are possible, only the highest scoring alignment is utilized, following an iterative procedure. To calculate an algorithm score of a given peptide in a test set, the  
25 ARB values corresponding to the sequence of the peptide are multiplied. If this product exceeds a chosen threshold, the peptide is predicted to bind. Appropriate thresholds are chosen as a function of the degree of stringency of prediction desired.

#### *Selection of HLA-A2 supertype cross-reactive peptides*

30 Complete polypeptide sequences from fourteen HCV isolates were aligned, then scanned, utilizing motif identification software, to identify conserved 9- and 10-mer sequences containing the HLA-A2-supermotif main anchor specificity.

A total of 231 conserved, HLA-A2 supermotif-positive sequences were identified. These peptides were then evaluated for the presence of A\*0201 preferred secondary anchor residues using A\*0201-specific polynomial algorithms. A total of 67 conserved, motif-bearing and algorithm-positive sequences were identified.

5        Fifty of these conserved, motif-containing 9- and 10-mer peptides were tested for their capacity to bind to purified HLA-A\*0201 molecules *in vitro* (HLA-A\*0201 is considered a prototype A2 supertype molecule). Sixteen peptides bound A\*0201 with IC<sub>50</sub> values ≤500 nM; 4 with high binding affinities (IC<sub>50</sub> values ≤50 nM) and 12 with intermediate binding affinities, in the 50-500 nM range (Table XXVI).

10        These 16 peptides were then tested for binding to additional A2-supertype molecules (A\*0202, A\*0203, A\*0206, and A\*6802). As shown in Table XXVI, most of these peptides were found to be A2-supertype cross-reactive binders. More specifically, 12/16 (75%) peptides bound at least three of the five A2-supertype molecules tested.

15    *Selection of HLA-A3 supermotif-bearing epitopes*

The sequences from the same fourteen known HCV isolates scanned above were also examined for the presence of conserved peptides with the HLA-A3-supermotif primary anchors. A total of 71 conserved 9- or 10-mer motif containing sequences were identified. Further analysis using the A03 and A11 algorithms (see, e.g., Gulukota et al,  
20    *J. Mol. Biol.* 267:1258-1267, 1997 and Sidney et al, *Human Immunol.* 45:79-93, 1996) identified 39 sequences that scored high in either or both algorithms. Twenty seven of the 39 peptides were synthesized and tested for binding to HLA-A\*03 and HLA-A\*11, the two most prevalent A3-supertype molecules. Fifteen peptides were identified which bound A3 and/or A11 with binding affinities of ≤500 nM (Table XXVII). These peptides  
25    were then tested for binding cross-reactivity to the other common A3-supertype alleles (A\*3101, A\*3301, and A\*6801). Seven of the 15 peptides bound at least three of the five HLA-A3-supertype molecules tested.

In the course of an independent series of experiments (Kubo *et al.*, *J. Immunol.* 152:3913-3924, 1994), one peptide, HCV NS3 1262, not identified by the selection  
30    criteria utilized above because it does not have the A3-supermotif main anchor specificity, was determined to be cross-reactive in the A3-supertype, binding A\*03, A\*11, and A\*6801. It is also shown in Table XXVII. Interestingly, this peptide

represents a single residue N-terminal truncation of peptide 1073.14, which is also shown in Table XXVII.

In summary, 8 peptides that bind 3 or more A3-supertype molecules derived from conserved regions of the HCV genome were identified.

5

*Selection of HLA-B7 supermotif bearing epitopes*

When the same fourteen HCV isolates were also analyzed for the presence of conserved 9- or 10-mer peptides with the HLA-B7-supermotif, 35 sequences were identified. The corresponding peptides were synthesized and tested for binding to HLA-

10 B\*0702, the most common B7-supertype allele (*i.e.*, the prototype B7 supertype allele). Thirteen peptides bound B\*0702 with  $IC_{50}$  of  $\leq 500$  nM (Table XXVIIIa). These 13 peptides were then tested for binding to other common B7-supertype molecules (B\*3501, B\*51, B\*5301, and B\*5401). As shown in Table XXVIIIa, only 1 peptide (Core 169) was capable of binding to three or more of the five B7-supertype alleles tested.

15 To identify additional B7-supertype epitopes, further studies were undertaken. The protein sequences from the fourteen HCV isolates utilized above were again examined to identify conserved, motif-containing 8- and 11-mers. The isolates were also examined for 9- and 10-mer sequences allowing for lower conservancy (51%-78%). Twenty-five 8-mers, sixteen 11-mers, and thirty-five 9- and 10-mers were identified, synthesized, and tested for binding to B\*0702. Thirteen peptides bound with high or intermediate affinity ( $IC_{50} \leq 500$  nM) (Table XXVIIIb). These peptides were additionally screened for binding to other B7-supertype molecules. Only one cross-reactive binder, the NS3 1378 8-mer (peptide 29.0035/1260.04), was identified (Table XXVIIIb).

20 In summary, a total of two cross-reactive B7-supertype binders were identified (Core 169 and NS3 1378).

*Selection of A1 and A24 motif-bearing epitopes*

To further increase population coverage, HLA-A1 and -A24 epitopes can also be incorporated into potential vaccine constructs.

30 In a previous analysis, two A1 and three A24 binders, 100% conserved among four strains of HCV, were identified (Wentworth *et al.*, *Int. Immunol.* 8:651-659, 1996). An analysis of the protein sequence data from the fourteen HCV strains utilized above demonstrated that these peptides were >79% conserved, and also identified an additional



eleven A1- and twenty five A24-motif-containing conserved sequences (see Table XXIXA and B). Eight of the additional eleven A1 peptides and seven of the additional twenty five A24 peptides were tested for binding to the appropriate HLA molecule (*i.e.*, A1 or A24). Overall, as shown in Table XXIX, four A1-motif peptides (A) and three  
5 A24-motif peptides (B) have been found with binding capacities of 500 nM or less for the appropriate allele-specific HLA molecule.

Analysis of the HLA-A2 and A3 supermotif-bearing epitopes identified above revealed that in 13/14 cases, peptides binding the supertype prototype HLA molecule (*i.e.* A\*0201 for the A2 supertype, and A\*0301 for the A3 supertype) with an IC<sub>50</sub> of less than  
10 100nM were cross-reactive and recognized by HCV-infected patients as described in Example 3, which follows. Based on these observations, two A1 peptides and one A24 peptide epitopes were also selected as candidates for inclusion in vaccine compositions; these peptides bind the appropriate HLA molecule with an IC<sub>50</sub> of less than 100nM.

### 15 Example 3: Confirmation of Immunogenicity

#### *Evaluation of A\*0201 immunogenicity*

It has been shown that CTL induced in A\*0201/K<sup>b</sup> transgenic mice exhibit specificity similar to CTL induced in the human system (*see, e.g.*, Vitiello *et al.*, *J. Exp. Med.* 173:1007-1015, 1991; Wentworth *et al.*, *Eur. J. Immunol.* 26:97-101, 1996).

20 Accordingly, these mice were used to evaluate the immunogenicity of the twelve conserved A2-supertype cross-reactive peptides identified in Example 2 above.

CTL induction in transgenic mice following peptide immunization has been described (Vitiello *et al.*, *J. Exp. Med.* 173:1007-1015, 1991; Alexander *et al.*; *J. Immunol.* 159:4753-4761, 1997). In these studies, mice were injected subcutaneously at  
25 the base of the tail with each peptide (50 µg/mouse) emulsified in IFA in the presence of an excess of an IA<sup>b</sup>-restricted helper peptide (140 µg/mouse) (HBV core 128-140, Sette *et al.*, *J. Immunol.* 153:5586-5592, 1994). Eleven days after injection, splenocytes were incubated in the presence of peptide-loaded syngenic LPS blasts. After six days, cultures were assayed for cytotoxic activity using peptide-pulsed targets. The data, summarized in  
30 Table XXX, indicate that 7 of the 12 peptides (58%) were capable of inducing primary CTL responses in A\*0201/K<sup>b</sup> transgenic mice. (For these studies, a peptide was considered positive if it induced CTL (L.U. 30/10<sup>6</sup> cells ≥2 in at least two transgenic animals (Wentworth *et al.*, *Eur. J. Immunol.* 26:97-101, 1996).

The conserved, cross reactive candidate CTL epitopes were also tested for recognition *in vitro* by PBMCs obtained from HCV-infected patients. Briefly, PBMC from patients infected with HCV were cultured in the presence of 10 µg/ml of synthetic peptide. After 7 and 14 days, the cultures were restimulated with peptide. The cultures were assayed for cytolytic activity on day 21 using target cells pulsed with the specific peptide in a standard four hour <sup>51</sup>Cr release assay. The data are summarized in Table XXX. As shown, all 12 peptides are CTL epitopes recognized by PBMC from HCV-infected patients. From the data in Table XXX, it is interesting to note that HLA transgenics did not fully reveal the immunogenicity of some peptides that were positive in recall responses. This apparent discrepancy may reflect differences in the route of immunization utilized (*e.g.*, natural infection versus peptide immunization), or CTL repertoire.

#### *Evaluation of A\*03/A11 immunogenicity*

The immunogenicity of six of the eight A3-supertype cross-reactive peptides identified in Example 2 above was evaluated in HLA-A11/K<sup>b</sup> transgenic mice, using the protocol described above for HLA-A2 transgenic mice (Alexander *et al.*, *J. Immunol.* 159:4753-4761, 1997). Five of these six peptides were able to induce primary CTL responses (Table XXXI).

All eight peptides were also studied by collaborators using PBMC cultures from HCV infected patients and contacts of such patients. This data is also summarized in Table XXXI. Briefly, all eight peptides were recognized by HCV infected individuals.

#### *Evaluation of B7 immunogenicity*

One of the two B7-supertype cross-reactive peptides (1145.12, Core 169) has been evaluated for immunogenicity in HCV-infected patients. Two independent collaborators have shown that this peptide is indeed immunogenic, and is recognized by T cells from HCV-infected patients (Chang *et al.*, *J. Immunol.* 162:1156-1164, 1999)

#### Example 4: Implementation of the Extended Supermotif to Improve the Binding Capacity of Native Epitopes by Creating Analogs

HLA motifs and supermotifs (comprising primary and/or secondary residues) are useful in the identification and preparation of highly cross-reactive native peptides, as demonstrated herein. Moreover, the definition of HLA motifs and supermotifs also

allows one to engineer highly cross-reactive epitopes by identifying residues within a native peptide sequence which can be analogued, or "fixed" to confer upon the peptide certain characteristics, *e.g.* greater cross-reactivity within the group of HLA molecules that comprise a supertype, and/or greater binding affinity for some or all of those HLA molecules. Examples of analog peptides that exhibit modulated binding affinity are set forth in this example.

#### *Analoging at Primary Anchor Residues*

As shown in Example 2, more than ten different HCV-derived, A2-supertype-restricted epitopes were identified. Peptide engineering strategies are implemented to further increase the cross-reactivity of the candidate epitopes identified above which bind 3/5 of the A2 supertype alleles tested. On the basis of the data disclosed, *e.g.*, in related and co-pending U.S.S.N 09/226,775, the main anchors of A2-supermotif-bearing peptides are altered, for example, to introduce a preferred L, I, V, or M at position 2, and I or V at the C-terminus.

To analyze the cross-reactivity of the analog peptides, each engineered analog is initially tested for binding to the prototype A2 supertype allele A\*0201, then, if A\*0201 binding capacity is maintained, for A2-supertype cross-reactivity.

Similarly, analogs of HLA-A3 supermotif-bearing epitopes may also be generated. For example, peptides binding to 3/5 of the A3-supertype molecules may be engineered at primary anchor residues to possess a preferred residue (V, S, M, or A) at position 2.

The analog peptides are then tested for the ability to bind A\*03 and A\*11 (prototype A3 supertype alleles). Those peptides that demonstrate  $\leq 500$  nM binding capacity are then tested for A3-supertype cross-reactivity.

Similarly to the A2- and A3- motif bearing peptides, peptides binding 3 or more B7-supertype alleles may be improved, where possible, to achieve increased cross-reactive binding. B7 supermotif-bearing peptides may, for example, be engineered to possess a preferred residue (V, I, L, or F) at the C-terminal primary anchor position, as demonstrated by Sidney *et al.* (*J. Immunol.* 157:3480-3490, 1996).

#### *Analoging at Secondary Anchor Residues*

Moreover, HLA supermotifs are of value in engineering highly cross-reactive peptides and/or peptides that bind HLA molecules with increased affinity by identifying

particular residues at secondary anchor positions that are associated with such properties. Demonstrating this, the binding capacity of a peptide representing a discreet single amino acid substitution at position one was analyzed. Peptide 1145.13 (Table XXVIIIc), which represents the substitution of L to F at position 1 of the core 169 sequence, binds all five  
5 B7-supertype molecules with a good affinity (all IC<sub>50</sub> values  $\leq$  132 nM), and in 3 instances has higher affinity over that of the parent peptide by >35-fold.

Because so few B7-supertype cross-reactive epitopes were identified, our results from previous binding evaluations were analyzed to identify conserved (8-, 9-, 10-, or 11-mer) peptides which bind, minimally, 3/5 B7 supertype molecules with weak affinity  
10 (IC<sub>50</sub> of 500nM-5 $\mu$ M). This analysis identified 9 peptides, 6 of which are analogued (including core 169 which had been previously analogued). These peptides are tested for enhanced binding affinity and B7-supertype cross-reactivity.

Engineered analogs with sufficiently improved binding capacity or cross-reactivity are tested for immunogenicity in HLA-B7-transgenic mice, following for  
15 example, IFA immunization or lipopeptide immunization.

In conclusion, these data demonstrate that by the use of even single amino acid substitutions, it is possible to increase the binding affinity and/or cross-reactivity of peptide ligands for HLA supertype molecules.

20 Example 5: Identification of conserved HCV-derived sequences with HLA-DR binding motifs

Peptide epitopes bearing an HLA class II supermotif or motif may also be identified as outlined below using methodology similar to that described in Examples 1-3.

25 *Selection of HLA-DR-supermotif-bearing epitopes*

To identify HCV-derived, HLA class II HTL epitopes, the same fourteen HCV polyprotein sequences used for the identification of HLA Class I supermotif/motif sequences were analyzed for the presence of sequences bearing an HLA-DR-motif or supermotif. Specifically, 15-mer sequences were selected comprising a DR-supermotif,  
30 further comprising a 9-mer core, and three-residue N- and C-terminal flanking regions (15 amino acids total). It was also required that the 15-mer sequence be conserved in at least 79% (11/14) of the HCV strains analyzed. These criteria identified a total of 49 non-redundant sequences, which are shown in Table XXXIIA. (In the context of Class II

epitopes, a sequence is considered operationally redundant if more than 80% of its sequence overlaps with another peptide.)

Protocols for predicting peptide binding to DR molecules have been developed (Southwood *et al.*, *J. Immunol.* 160:3363-3373, 1998). These protocols, specific for individual DR molecules, allow the scoring, and ranking, of 9-mer core regions. Each protocol not only scores peptide sequences for the presence of DR-supermotif primary anchors (i.e., at position 1 and position 6) within a 9-mer core, but additionally evaluates sequences for the presence of secondary anchors. Using allele specific selection tables (see, *e.g.*, Southwood *et al.*, *ibid.*), it has been found that these protocols efficiently select peptide sequences with a high probability of binding a particular DR molecule. Additionally, it has been found that performing these protocols in tandem, specifically those for DR1, DR4w4, and DR7, can efficiently select DR cross-reactive peptides.

To see if these protocols serve to identify additional epitopes, the same HCV polyproteins used above were re-scanned for the presence of 15-mer peptides with 9-mer core regions that were  $\geq 79\%$  (11/14 strains) conserved. This identified 152 sequences; 49 of which were identified previously, as described above. Next, the 9-mer core region of each of these peptides was scored using the DR1, DR4w4, and DR7 algorithms. Twenty-two peptides, including 12 new sequences (10 peptides were from the original set of 49) were found to have 9-mer cores with protocol-derived scores predictive of cross-reactive DR binders. The 12 additional sequences are shown in Table XXXIIB.

The conserved, HCV-derived peptides identified above were tested for their binding capacity for various common HLA-DR molecules. All peptides were initially tested for binding to the DR molecules in the primary panel: DR1, DR4w4, and DR7. Peptides binding at least 2 of these 3 DR molecules were then tested for binding to DR2w2  $\beta$ 1, DR2w2  $\beta$ 2, DR6w19, and DR9 molecules in secondary assays. Finally, peptides binding at least 2 of the 4 secondary panel DR molecules, and thus cumulatively at least 4 of 7 different DR molecules, were screened for binding to DR4w15, DR5w11, and DR8w2 molecules in tertiary assays. Peptides binding at least 7 of the 10 DR molecules comprising the primary, secondary, and tertiary screening assays were considered cross-reactive DR binders. The composition of these screening panels, and the phenotypic frequency of associated antigens, are shown in Table XXXIII.

Upon testing, it was found that 29 of the original 75 peptides (39%) bound two or more of the primary HLA molecules. Twenty-six of these cross-reactive binders were

then tested in the secondary assays, and nineteen were found to bind at least four of the seven HLA DR molecules in the primary and secondary panels. Finally, the nineteen peptides passing the secondary screening phase were tested for binding in the tertiary assays. As a result, nine peptides were identified which bound at least seven of ten  
5 common HLA-DR molecules. Table XXXIV shows these nine peptides and their binding capacity for each allele-specific HLA-DR molecule in the primary through tertiary panels. Also shown in Table XXXIV are two peptides (F134.05 and F134.08) for which a complete binding analysis was not performed. However, both of these peptides bound six of the seven HLA DR molecules tested. F134.08 nests peptide 1283.44, which bound  
10 eight of 10 allele-specific HLA molecules.

In conclusion, eleven cross-reactive DR-binding peptides, derived from six discrete (*i.e.* non-redundant) regions of the HCV genome, have been identified. Two of the six regions from which these epitopes were derived are covered by multiple, overlapping epitopes.

#### 15 *Selection of conserved DR3 motif peptides*

Because HLA-DR3 is an allele that is prevalent in Caucasian, Black, and Hispanic populations, DR3 binding capacity is an important criterion in the selection of HTL epitopes. However, data generated previously indicated that DR3 only rarely cross-reacts  
20 with other DR alleles (Sidney *et al.*, *J. Immunol.* 149:2634-2640, 1992; Geluk *et al.*, *J. Immunol.* 152:5742-5748, 1994; Southwood *et al.*, *J. Immunol.* 160:3363-3373, 1998). This is not entirely surprising in that the DR3 peptide-binding motif appears to be distinct from the specificity of most other DR alleles.

To efficiently identify peptides that bind DR3, target proteins were analyzed for  
25 conserved sequences carrying one of the two DR3 specific binding motifs reported by Geluk *et al.* (*J. Immunol.* 152:5742-5748, 1994). Fifteen sequences, including a peptide nested within a DR-supermotif sequence identified above (peptide Pape 22), were identified (Table XXXIId). Preferably, DR3 motifs will be found clustered in proximity with DR supermotif regions.

30 Fourteen of the fifteen peptides containing a DR3 motif were tested for their DR3 binding capacity. Two peptides (CH35.0106 and CH35.0107) were found to bind DR3 with an affinity of 1 $\mu$ M or less (Table XXXV), and thereby qualify as HLA class II high affinity binders.

DR3 binding epitopes identified in this manner may then be included in vaccine compositions with DR supermotif-bearing peptide epitopes.

Example 6: Immunogenicity of candidate HCV-derived HTL epitopes and known dominant HCV HTL epitope

In the course of collaborative studies with G. Pape and C. Ferrari, eight conserved, HCV-derived peptides have been identified which are recognized by HCV-infected individuals.

One of these studies (Diepolder *et al.*, *J. Virol.* 71:6011-6019, 1997), identified peptide F98.05, which spans residues 1248-1261 of the NS3 protein, as an immunodominant CD4+ T-cell epitope that was recognized by 14/23 NS3-specific CD4+ T-cell clones from 4/5 patients with acute hepatitis C infection. This epitope, shown above to be an HLA-DR cross-reactive binder (see Table XXXIV), was capable of being presented to helper CD4+ T cells by multiple HLA molecules (DR4, DR11, DR12, DR13, and DR16). Two other peptides, Pape 22 and Pape 29, were also recognized by CD4+ T cell clones, although, in a more limited context; correspondingly, neither of these peptides are DR-cross-reactive binders.

By direct peripheral blood T cell stimulation and by fine specificity analysis of HCV-specific T-cell lines and clones, studies done in collaboration with Ferrari's group identified 6 immunodominant epitopes, including one also identified in the Pape collaboration, that are derived from conserved regions of the core, NS3, and NS4 proteins. These epitopes were also found to be cross-reactive, being presented to T cells in the context of different Class II molecules. Three of the 6 epitopes, F98.04 (F134.03), F134.05 and F134.08, are cross-reactive HLA-DR binders (see Table XXXIV).

In conclusion, the immunogenicity of 8 epitopes derived from conserved regions of the HCV genome has been demonstrated. Three of these epitopes (F98.05, F134.05, and F134.08; see Table XXXIV) are broadly cross-reactive HLA-DR binding peptides.

Example 7. Calculation of phenotypic frequencies of HLA-supertypes in various ethnic backgrounds to determine breadth of population coverage

This example illustrates the assessment of the breadth of population coverage of a vaccine composition comprised of multiple epitopes comprising multiple supermotifs and/or motifs.

In order to analyze population coverage, gene frequencies of HLA alleles were determined. Gene frequencies for each HLA allele were calculated from antigen or allele frequencies utilizing the binomial distribution formulae  $gf=1-(\text{SQRT}(1-af))$  (see, e.g., Sidney *et al.*, *Human Immunol.* 45:79-93, 1996). To obtain overall phenotypic

5 frequencies, cumulative gene frequencies were calculated, and the cumulative antigen frequencies derived by the use of the inverse formula  $[af=1-(1-Cgf)^2]$ .

Where frequency data was not available at the level of DNA typing, correspondence to the serologically defined antigen frequencies was assumed. To obtain total potential supertype population coverage no linkage disequilibrium was assumed, and

10 only alleles confirmed to belong to each of the superotypes were included (minimal estimates). Estimates of total potential coverage achieved by inter-loci combinations were made by adding to the A coverage the proportion of the non-A covered population that could be expected to be covered by the B alleles considered (e.g.,  $\text{total}=A+B*(1-A)$ ). Confirmed members of the A3-like supertype are A3, A11, A31, A\*3301, and A\*6801.

15 Although the A3-like supertype may also include A34, A66, and A\*7401, these alleles were not included in overall frequency calculations. Likewise, confirmed members of the A2-like supertype family are A\*0201, A\*0202, A\*0203, A\*0204, A\*0205, A\*0206, A\*0207, A\*6802, and A\*6901. Finally, the B7-like supertype-confirmed alleles are: B7, B\*3501-03, B51, B\*5301, B\*5401, B\*5501-2, B\*5601, B\*6701, and B\*7801 (potentially

20 also B\*1401, B\*3504-06, B\*4201, and B\*5602).

Population coverage achieved by combining the A2-, A3- and B7-superotypes is approximately 86% in five major ethnic groups (see Table XXI). Coverage may be extended by including peptides bearing the A1 and A24 motifs. On average, A1 is present in 12% and A24 in 29% of the population across five different major ethnic

25 groups (Caucasian, North American Black, Chinese, Japanese, and Hispanic). Together, these alleles are represented with an average frequency of 39% in these same ethnic populations. The total coverage across the major ethnicities when A1 and A24 are combined with the coverage of the A2-, A3- and B7-supertype alleles is >95%. An analagous approach can be used to estimate population coverage achieved with

30 combinations of class II motif-bearing epitopes.

#### *Summary of candidate HLA class I and class II epitopes*

In summary, on the basis of the data presented in the above examples, 26 CTL candidate peptide epitopes derived from conserved regions of the HCV virus have been



identified (Table XXXVIa). These include twelve HLA-A2 supermotif-bearing epitopes, eight HLA-A3 supermotif-bearing epitopes, and one HLA-B7 supermotif-bearing epitope, each capable of binding to multiple A2-, A3-, or B7-supertype molecules, and immunogenic in HLA transgenic mice or antigenic for human PBL (with the exception of peptide 29.0035/1260.04). Additional epitopes not evaluated for immunogenicity are also included. They are an additional B7-supermotif-bearing epitope and two HLA-A1 and one HLA-A24 high-affinity binding peptides. A known HLA-A31 restricted epitope (VGIYLLPNR), which also binds HLA-A33, is also set out in Table XXXVIa and is useful in combination with other Class I or Class II epitopes.

With these 26 CTL epitopes (as disclosed herein and from the art), average population coverage, (*i.e.*, recognition of at least one HCV epitope), is predicted to be greater than 95% in each of five major ethnic populations. The potential redundancy of coverage afforded by 25 of these epitopes (the peptide 24.0086 was not included) was estimated using the game theory Monte Carlo simulation analysis, which is known in the art (see *e.g.*, Osborne, M.J. and Rubinstein, A. "A course in game theory" MIT Press, 1994). As shown in Figure 1, it is estimated that 90% of the individuals in a population comprised of the Caucasian, North American Black, Japanese, Chinese, and Hispanic ethnic groups would recognize 2 or more of the candidate epitopes described herein.

A list of HCV-derived HTL epitopes that would be preferred for use in the design of minigene constructs or other vaccine formulations is summarized in Table XXXVIb. As shown, 9 different peptide-binding regions have been identified which bind multiple HLA-DR molecules or bind HLA-DR3. (In the case of the NS4 1914-1935 region, the longer peptide, F134.08, recognized by patients, was chosen over the shorter peptide, 1283.44. The longer peptide essentially incorporates the shorter peptide, and also binds additional DR molecules that the shorter peptide does not bind.) Three of these peptides have been recognized as dominant epitopes in HCV infected patients.

It is estimated that each of 10 common DR molecules recognizing the DR supermotif, and DR3, are covered by a minimum of 2 epitopes. Correspondingly, the total estimated population coverage represented by this panel of epitopes is in excess of 91% in each of the 5 major ethnic populations (Table XXXVII).

Example 8: Recognition Of Generation Of Endogenous Processed Antigens After Priming

This example determines that CTL induced by native or analogued peptide epitopes identified and selected as described in Examples 1-6 recognize endogenously synthesized, *i.e.*, native antigens.

Effector cells isolated from transgenic mice that are immunized with peptide epitopes as in Example 3, for example HLA-A2 supermotif-bearing epitopes, are re-stimulated *in vitro* using peptide-coated stimulator cells. Six days later, effector cells are assayed for cytotoxicity and the cell lines that contain peptide-specific cytotoxic activity are further re-stimulated. An additional six days later, these cell lines are tested for cytotoxic activity on <sup>51</sup>Cr labeled Jurkat-A2.1/K<sup>b</sup> target cells in the absence or presence of peptide, and also tested on <sup>51</sup>Cr labeled target cells bearing the endogenously synthesized antigen, *i.e.* cells that are stably transfected with HCV expression vectors.

The result will demonstrate that CTL lines obtained from animals primed with peptide epitope recognize endogenously synthesized HCV antigen. The choice of transgenic mouse model to be used for such an analysis depends upon the epitope(s) that is being evaluated. In addition to HLA-A\*0201/K<sup>b</sup> transgenic mice, several other transgenic mouse models including mice with human A11, which may also be used to evaluate A3 epitopes, and B7 alleles have been characterized and others (*e.g.*, transgenic mice for HLA-A1 and A24) are being developed. HLA-DR1 and HLA-DR3 mouse models have also been developed, which may be used to evaluate HTL epitopes.

Example 9: Activity Of CTL-HTL Conjugated Epitopes In Transgenic Mice

This example illustrates the induction of CTLs and HTLs in transgenic mice by use of an HCV CTL/HTL peptide conjugate whereby the vaccine composition comprises peptides administered to an HCV-infected patient or an individual at risk for HCV. The peptide composition can comprise multiple CTL and/or HTL epitopes. This analysis demonstrates enhanced immunogenicity that can be achieved by inclusion of one or more HTL epitopes in a vaccine composition. Such a peptide composition can comprise a lipidated HTL epitope conjugated to a preferred CTL epitope containing, for example, at least one CTL epitope selected from Table XXVI-XXIX, or an analog of that epitope. The HTL epitope is, for example, selected from Table XXXII.

Lipopeptide preparation: Lipopeptides are prepared by coupling the appropriate fatty acid to the amino terminus of the resin bound peptide. A typical procedure is as

follows: A dichloromethane solution of a four-fold excess of a pre-formed symmetrical anhydride of the appropriate fatty acid is added to the resin and the mixture is allowed to react for two hours. The resin is washed with dichloromethane and dried. The resin is then treated with trifluoroacetic acid in the presence of appropriate scavengers [e.g. 5% (v/v) water] for 60 minutes at 20°C. After evaporation of excess trifluoroacetic acid, the crude peptide is washed with diethyl ether, dissolved in methanol and precipitated by the addition of water. The peptide is collected by filtration and dried.

Immunization procedures: Immunization of transgenic mice is performed as described (Alexander *et al.*, *J. Immunol.* 159:4753-4761, 1997). For example, A2/K<sup>b</sup> mice, which are transgenic for the human HLA A2.1 allele and are useful for the assessment of the immunogenicity of HLA-A\*0201 motif- or HLA-A2 supermotif-bearing epitopes, are primed subcutaneously (base of the tail) with 0.1 ml of peptide conjugate formulated in saline, or DMSO/saline. Seven days after priming, splenocytes obtained from these animals are restimulated with syngenic irradiated LPS-activated lymphoblasts coated with peptide.

Cell lines: Target cells for peptide-specific cytotoxicity assays are Jurkat cells transfected with the HLA-A2.1/K<sup>b</sup> chimeric gene (*e.g.*, Vitiello *et al.*, *J. Exp. Med.* 173:1007, 1991)

*In vitro* CTL activation: One week after priming, spleen cells (30x10<sup>6</sup> cells/flask) are co-cultured at 37°C with syngeneic, irradiated (3000 rads), peptide coated lymphoblasts (10x10<sup>6</sup> cells/flask) in 10 ml of culture medium/T25 flask. After six days, effector cells are harvested and assayed for cytotoxic activity.

Assay for cytotoxic activity: Target cells (1.0 to 1.5x10<sup>6</sup>) are incubated at 37°C in the presence of 200 µl of <sup>51</sup>Cr. After 60 minutes, cells are washed three times and resuspended in R10 medium. Peptide is added where required at a concentration of 1 µg/ml. For the assay, 10<sup>4</sup> <sup>51</sup>Cr-labeled target cells are added to different concentrations of effector cells (final volume of 200 µl) in U-bottom 96-well plates. After a 6 hour incubation period at 37°C, a 0.1 ml aliquot of supernatant is removed from each well and radioactivity is determined in a Micromedic automatic gamma counter. The percent specific lysis is determined by the formula: percent specific release = 100 x (experimental release - spontaneous release)/(maximum release - spontaneous release). To facilitate comparison between separate CTL assays run under the same conditions, % <sup>51</sup>Cr release data is expressed as lytic units/10<sup>6</sup> cells. One lytic unit is arbitrarily defined as the number of effector cells required to achieve 30% lysis of 10,000 target cells in a 6

hour  $^{51}\text{Cr}$  release assay. To obtain specific lytic units/ $10^6$ , the lytic units/ $10^6$  obtained in the absence of peptide is subtracted from the lytic units/ $10^6$  obtained in the presence of peptide. For example, if 30%  $^{51}\text{Cr}$  release is obtained at the effector (E): target (T) ratio of 50:1 (i.e.,  $5 \times 10^5$  effector cells for 10,000 targets) in the absence of peptide and 5:1 (i.e.,  $5 \times 10^4$  effector cells for 10,000 targets) in the presence of peptide, the specific lytic units would be:  $[(1/50,000) - (1/500,000)] \times 10^6 = 18 \text{ LU}$ .

The results are analyzed to assess the magnitude of the CTL responses of animals injected with the immunogenic CTL/HTL conjugate vaccine preparation and are compared to the magnitude of the CTL response achieved using the CTL epitope as outlined in Example 3. Analyses similar to this may be performed to evaluate the immunogenicity of peptide conjugates containing multiple CTL epitopes and/or multiple HTL epitopes. In accordance with these procedures it is found that a CTL response is induced, and concomitantly that an HTL response is induced upon administration of such compositions.

Example 10. Selection of CTL and HTL epitopes for inclusion in an HCV-specific vaccine.

This example illustrates the procedure for the selection of peptide epitopes for vaccine compositions of the invention. The peptides in the composition can be in the form of a nucleic acid sequence, either single or one or more sequences (i.e., minigene) that encodes peptide(s), or may be single and/or polyepitopic peptides.

Epitopes are selected which, upon administration, mimic immune responses that have been observed to be correlated with tumor clearance. For example, vaccine can include 3-4 epitopes that come from at least one HCV antigen region. Epitopes from one region can be used in combination with epitopes from one or more additional HCV antigen regions. Analogs of epitopes can also be selected for inclusion in the vaccine.

Epitopes are often selected that have a binding affinity of an  $\text{IC}_{50}$  of 500 nM or less for an HLA class I molecule, or for class II, an  $\text{IC}_{50}$  of 1000 nM or less.

Sufficient supermotif bearing peptides, or a sufficient array of allele-specific motif bearing peptides, are selected to give broad population coverage. For example, epitopes are selected to provide at least 80% population coverage. A Monte Carlo analysis, a statistical evaluation known in the art, can be employed to assess breadth, or redundancy, of population coverage.

When creating a polyepitopic compositions, *e.g.* a minigene, it is typically desirable to generate the smallest peptide possible that encompasses the epitopes of interest. The principles employed are similar, if not the same, as those employed when selecting a peptide comprising nested epitopes. Additionally, however, upon  
5 determination of the nucleic acid sequence to be provided as a minigene, the peptide sequence encoded thereby is analyzed to determine whether any "junctional epitopes" have been created. A junctional epitope is a potential HLA binding epitope, as predicted, *e.g.*, by motif analysis. Junctional epitopes are generally to be avoided because the recipient may bind to an HLA molecule and generate an immune response to that epitope,  
10 which is not present in a native protein sequence.

Peptide epitopes for inclusion in vaccine compositions are, for example, selected from those listed in Tables XXVI-XXIX and Table XXXII. A vaccine composition comprised of selected peptides, when administered, is safe, efficacious, and elicits an immune response similar in magnitude of an immune response that clears an acute HCV  
15 infection.

#### Example 11: Construction of Minigene Multi-Epitope DNA Plasmids

This example provides guidance for the construction of a minigene expression plasmid. Minigene plasmids may, of course, contain various configurations of CTL  
20 and/or HTL epitopes or epitope analogs as described herein. Examples of the construction and evaluation of expression plasmids are described, for example, in co-pending U.S.S.N. 09/311,784 filed 5/13/99. An example of such a plasmid for the expression of HCV epitopes is shown in Figure 2, which illustrates the orientation of HCV peptide epitopes in a minigene construct.

25 A minigene expression plasmid may include multiple CTL and HTL peptide epitopes. In the present example, HLA-A2, -A3, -B7 supermotif-bearing peptide epitopes and HLA-A1 and -A24 motif-bearing peptide epitopes are used in conjunction with DR supermotif-bearing epitopes and/or DR3 epitopes (Figure 2). Preferred epitopes are identified, for example, in Tables XXVI-XXIX and XXXII. HLA class I supermotif or  
30 motif-bearing peptide epitopes derived from multiple HCV antigens, *e.g.*, the core, NS4, NS3, NS5, NS1/E2, are selected such that multiple supermotifs/motifs are represented to ensure broad population coverage. Similarly, HLA class II epitopes are selected from multiple HCV antigens to provide broad population coverage, *i.e.* both HLA DR-1-4-7 supermotif-bearing epitopes and HLA DR-3 motif-bearing epitopes are selected for

inclusion in the minigene construct. The selected CTL and HTL epitopes are then incorporated into a minigene for expression in an expression vector.

This example illustrates the methods to be used for construction of such a minigene-bearing expression plasmid. Other expression vectors that may be used for minigene compositions are available and known to those of skill in the art.

The minigene DNA plasmid contains a consensus Kozak sequence and a consensus murine kappa Ig-light chain signal sequence followed by CTL and/or HTL epitopes selected in accordance with principles disclosed herein. The sequence encodes an open reading frame fused to the Myc and His antibody epitope tag coded for by the pcDNA 3.1 Myc-His vector.

Overlapping oligonucleotides, for example eight oligonucleotides, averaging approximately 70 nucleotides in length with 15 nucleotide overlaps, are synthesized and HPLC-purified. The oligonucleotides encode the selected peptide epitopes as well as appropriate linker nucleotides, Kozak sequence, and signal sequence. The final multiepitope minigene is assembled by extending the overlapping oligonucleotides in three sets of reactions using PCR. A Perkin/Elmer 9600 PCR machine is used and a total of 30 cycles are performed using the following conditions: 95°C for 15 sec, annealing temperature (5° below the lowest calculated T<sub>m</sub> of each primer pair) for 30 sec, and 72°C for 1 min.

For the first PCR reaction, 5 µg of each of two oligonucleotides, *i.e.*, an amplification primer pair, are annealed and extended: Oligonucleotides 1+2, 3+4, 5+6, and 7+8 are combined in 100 µl reactions containing *Pfu* polymerase buffer (1x= 10 mM KCL, 10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 20 mM Tris-chloride, pH 8.75, 2 mM MgSO<sub>4</sub>, 0.1% Triton X-100, 100 µg/ml BSA), 0.25 mM each dNTP, and 2.5 U of *Pfu* polymerase. The full-length dimer products are gel-purified, and two reactions containing the product of 1+2 and 3+4, and the product of 5+6 and 7+8 are mixed, annealed, and extended for 10 cycles. Half of the two reactions are then mixed, and 5 cycles of annealing and extension carried out before flanking primers are added to amplify the full length product for 25 additional cycles. The full-length product is gel-purified and cloned into pCR-blunt (Invitrogen) and individual clones are screened by sequencing.

Example 12. The plasmid construct and the degree to which it induces immunogenicity.

The degree to which the plasmid construct prepared using the methodology outlined in Example 11 is able to induce immunogenicity is evaluated through *in vivo*

injections into mice and subsequent *in vitro* assessment of CTL and HTL activity, which are analysed using cytotoxicity and proliferation assays, respectively, as detailed *e.g.* in U.S.S.N. 09/311,784 filed 5/13/99 and Alexander *et al.*, *Immunity* 1:751-761, 1994. For example, to assess the capacity of a pMin minigene construct that contains HLA-A2 supermotif epitopes to induce CTLs *in vivo*, HLA-A2.1/K<sup>b</sup> transgenic mice are immunized intramuscularly with 100 µg of naked cDNA. As a means of comparing the level of CTLs induced by cDNA immunization, a control group of animals is also immunized with an actual peptide composition that comprises multiple epitopes synthesized as a single polypeptide as they would be encoded by the minigene.

Splenocytes from immunized animals are stimulated twice with each of the respective compositions (peptide epitopes encoded in the minigene or the polypeptidic peptide), then assayed for peptide-specific cytotoxic activity in a <sup>51</sup>Cr release assay. The results indicate the magnitude of the CTL response directed against the A3-restricted epitope, thus indicating the *in vivo* immunogenicity of the minigene vaccine and polypeptidic vaccine. It is, therefore, found that the minigene elicits immune responses directed toward the HLA-A2 supermotif peptide epitopes as does the polypeptidic peptide vaccine. A similar analysis is also performed using other HLA-A3 and HLA-B7 transgenic mouse models to assess CTL induction by HLA-A3 and HLA-B7 motif or supermotif epitopes.

To assess the capacity of a class II epitope encoding minigene to induce HTLs *in vivo*, I-A<sup>b</sup> restricted mice, for example, are immunized intramuscularly with 100 µg of plasmid DNA. As a means of comparing the level of HTLs induced by DNA immunization, a group of control animals is also immunized with an actual peptide composition emulsified in complete Freund's adjuvant.

CD4+ T cells, *i.e.* HTLs, are purified from splenocytes of immunized animals and stimulated with each of the respective compositions (peptides encoded in the minigene). The HTL response is measured using a <sup>3</sup>H-thymidine incorporation proliferation assay, (*see, e.g.*, Alexander *et al.* *Immunity* 1:751-761, 1994). the results indicate the magnitude of the HTL response, thus demonstrating the *in vivo* immunogenicity of the minigene.

Alternatively, plasmid constructs can be evaluated *in vitro* by testing for epitope presentation by APC following transduction or transfection of the APC with an epitope-expressing nucleic acid construct. Such a study determines "antigenicity" and allows the use of human APC. The assay determines the ability of the epitope to be presented by the

APC in a context that is recognized by a T cell by quantifying the density of epitope-HLA class I complexes on the cell surface. Quantitation can be performed by directly measuring the amount of peptide eluted from the APC (*see, e.g., Sijts et al., J. Immunol.* 156:683-692, 1996; Demotz *et al., Nature* 342:682-684, 1989); or the number of peptide-HLA class I complexes can be estimated by measuring the amount of lysis or lymphokine release induced by infected or transfected target cells, and then determining the concentration of peptide necessary to obtain equivalent levels of lysis or lymphokine release (*see, e.g., Kageyama et al., J. Immunol.* 154:567-576, 1995).

10 Example 13: Peptide Composition for Prophylactic Uses

Vaccine compositions of the present invention are used to prevent HCV infection in persons who are at risk for such infection. For example, a polyepitopic peptide epitope composition (or a nucleic acid comprising the same) containing multiple CTL and HTL epitopes such as those selected in Examples 9 and/or 10, which are also selected to target greater than 80% of the population, is administered to individuals at risk for HCV infection. The composition is provided as a single lipidated polypeptide that encompasses multiple epitopes. The vaccine is administered in an aqueous carrier comprised of Freund's Incomplete Adjuvant. The dose of peptide for the initial immunization is from about 1 to about 50,000 µg, generally 100-5,000 µg, for a 70 kg patient. The initial administration of vaccine is followed by booster dosages at 4 weeks followed by evaluation of the magnitude of the immune response in the patient, by techniques that determine the presence of epitope-specific CTL populations in a PBMC sample. Additional booster doses are administered as required. The composition is found to be both safe and efficacious as a prophylaxis against HCV infection.

25 Alternatively, the polyepitopic peptide composition can be administered as a nucleic acid in accordance with methodologies known in the art and disclosed herein.

Example 14: Polyepitopic Vaccine Compositions Derived from Native HCV Sequences

A native HCV polyprotein sequence is screened, preferably using computer algorithms defined for each class I and/or class II supermotif or motif, to identify "relatively short" regions of the polyprotein that comprise multiple epitopes and is preferably less in length than an entire native antigen. This relatively short sequence that contains multiple distinct, even overlapping, epitopes is selected and used to generate a minigene construct. The construct is engineered to express the peptide, which



corresponds to the native protein sequence. The "relatively short" peptide is generally less than 250 amino acids in length, often less than 100 amino acids in length, preferably less than 75 amino acids in length, and more preferably less than 50 amino acids in length. The protein sequence of the vaccine composition is selected because it has  
5 maximal number of epitopes contained within the sequence, *i.e.*, it has a high concentration of epitopes. As noted herein, epitope motifs may be nested or overlapping (*i.e.*, frame shifted relative to one another). For example, with frame shifted overlapping epitopes, two 9-mer epitopes and one 10-mer epitope can be present in a 10 amino acid peptide. Such a vaccine composition is administered for therapeutic or prophylactic  
10 purposes.

The vaccine composition will preferably include, for example, three CTL epitopes and at least one HTL epitope from an HCV antigen. This polyepitopic native sequence is administered either as a peptide or as a nucleic acid sequence which encodes the peptide. Alternatively, an analog can be made of this native sequence, whereby one or more of the  
15 epitopes comprise substitutions that alter the cross-reactivity and/or binding affinity properties of the polyepitopic peptide.

The embodiment of this example provides for the possibility that an as yet undiscovered aspect of immune system processing will apply to the native nested sequence and thereby facilitate the production of therapeutic or prophylactic immune  
20 response-inducing vaccine compositions. Additionally such an embodiment provides for the possibility of motif-bearing epitopes for an HLA makeup that is presently unknown. Furthermore, this embodiment (absent analogs) directs the immune response to multiple peptide sequences that are actually present in native HCV antigens thus avoiding the need to evaluate any junctional epitopes. Lastly, the embodiment provides an economy of  
25 scale when producing nucleic acid vaccine compositions.

Related to this embodiment, computer programs can be derived in accordance with principles in the art, which identify in a target sequence, the greatest number of epitopes per sequence length.

#### 30 Example 15. Polyepitopic Vaccine Compositions Directed To Multiple Diseases

The HCV peptide epitopes of the present invention are used in conjunction with peptide epitopes from target antigens related to one or more other diseases, to create a vaccine composition that is useful for the prevention or treatment of HCV as well as the

one or more other disease(s). Examples of the other diseases include, but are not limited to, HIV, and HBV.

For example, a polyepitopic peptide composition comprising multiple CTL and HTL epitopes that target greater than 98% of the population may be created for  
5 administration to individuals at risk for both HCV and HIV infection. The composition can be provided as a single polypeptide that incorporates the multiple epitopes from the various disease-associated sources, or can be administered as a composition comprising one or more discrete epitopes.

10 Example 16. Use of peptides to evaluate an immune response

Peptides of the invention may be used to analyze an immune response for the presence of specific CTL or HTL populations directed to a prostate cancer-associated antigen. Such an analysis may be performed using multimeric complexes as described, e.g., by Ogg *et al.*, *Science* 279:2103-2106, 1998 and Greten *et al.*, *Proc. Natl. Acad. Sci.*  
15 *USA* 95:7568-7573, 1998. In the following example, peptides in accordance with the invention are used as a reagent for diagnostic or prognostic purposes, not as an immunogen.

In this example, highly sensitive human leukocyte antigen tetrameric complexes ("tetramers") are used for a cross-sectional analysis of, for example, HCV HLA-A\*0201-specific CTL frequencies from HLA A\*0201-positive individuals at different stages of  
20 disease or following immunization using an HCV peptide containing an A\*0201 motif. Tetrameric complexes are synthesized as described (Musey *et al.*, *N. Engl. J. Med.* 337:1267, 1997). Briefly, purified HLA heavy chain (A\*0201 in this example) and  $\beta$ 2-microglobulin are synthesized by means of a prokaryotic expression system. The heavy  
25 chain is modified by deletion of the transmembrane-cytosolic tail and COOH-terminal addition of a sequence containing a BirA enzymatic biotinylation site. The heavy chain,  $\beta$ 2-microglobulin, and peptide are refolded by dilution. The 45-kD refolded product is isolated by fast protein liquid chromatography and then biotinylated by BirA in the  
30 presence of biotin (Sigma, St. Louis, Missouri), adenosine 5'triphosphate and magnesium. Streptavidin-phycoerythrin conjugate is added in a 1:4 molar ratio, and the tetrameric product is concentrated to 1 mg/ml. The resulting product is referred to as tetramer-phycoerythrin.

For the analysis of patient blood samples, approximately one million PBMCs are centrifuged at 300g for 5 minutes and resuspended in 50  $\mu$ l of cold phosphate-buffered saline. Tri-color analysis is performed with the tetramer-phycoerythrin, along with anti-CD8-Tricolor, and anti-CD38. The PBMCs are incubated with tetramer and antibodies on ice for 30 to 60 min and then washed twice before formaldehyde fixation. Gates are applied to contain >99.98% of control samples. Controls for the tetramers include both A\*0201-negative individuals and A\*0201-positive uninfected donors. The percentage of cells stained with the tetramer is then determined by flow cytometry. The results indicate the number of cells in the PBMC sample that contain epitope-restricted CTLs, thereby readily indicating the extent of immune response to the HCV epitope, and thus the stage of HCV infection or exposure to a vaccine that elicits a protective or therapeutic response.

#### Example 17: Use of Peptide Epitopes to Evaluate Recall Responses

The peptide epitopes of the invention are used as reagents to evaluate T cell responses, such as acute or recall responses, in patients. Such an analysis may be performed on patients who have recovered from infection, who are chronically infected with HCV, or who have been vaccinated with an HCV vaccine.

For example, the class I restricted CTL response of persons who have been vaccinated may be analyzed. The vaccine may be any HCV vaccine. PBMC are collected from vaccinated individuals and HLA typed. Appropriate peptide epitopes of the invention that are preferably highly conserved and, optimally, bear supermotifs to provide cross-reactivity with multiple HLA supertype family members, are then used for analysis of samples derived from individuals who bear that HLA type.

PBMC from vaccinated individuals are separated on Ficoll-Histopaque density gradients (Sigma Chemical Co., St. Louis, MO), washed three times in HBSS (GIBCO Laboratories), resuspended in RPMI-1640 (GIBCO Laboratories) supplemented with L-glutamine (2mM), penicillin (50U/ml), streptomycin (50  $\mu$ g/ml), and Hepes (10mM) containing 10% heat-inactivated human AB serum (complete RPMI) and plated using microculture formats. A synthetic peptide comprising an epitope of the invention is added at 10  $\mu$ g/ml to each well and HBV core 128-140 epitope is added at 1  $\mu$ g/ml to each well as a source of T cell help during the first week of stimulation.

In the microculture format,  $4 \times 10^5$  PBMC are stimulated with peptide in 8 replicate cultures in 96-well round bottom plate in 100  $\mu$ l/well of complete RPMI. On

days 3 and 10, 100 ml of complete RPMI and 20 U/ml final concentration of rIL-2 are added to each well. On day 7 the cultures are transferred into a 96-well flat-bottom plate and restimulated with peptide, rIL-2 and  $10^5$  irradiated (3,000 rad) autologous feeder cells. The cultures are tested for cytotoxic activity on day 14. A positive CTL response requires two or more of the eight replicate cultures to display greater than 10% specific  $^{51}\text{Cr}$  release, based on comparison with uninfected control subjects as previously described (Rehermann, *et al.*, *Nature Med.* 2:1104,1108, 1996; Rehermann *et al.*, *J. Clin. Invest.* 97:1655-1665, 1996; and Rehermann *et al.* *J. Clin. Invest.* 98:1432-1440, 1996).

Target cell lines are autologous and allogeneic EBV-transformed B-LCL that are either purchased from the American Society for Histocompatibility and Immunogenetics (ASHI, Boston, MA) or established from the pool of patients as described (Guilhot, *et al.* *J. Virol.* 66:2670-2678, 1992).

Cytotoxicity assays are performed in the following manner. Target cells consist of either allogeneic HLA-matched or autologous EBV-transformed B lymphoblastoid cell line that are incubated overnight with the synthetic peptide epitope of the invention at 10  $\mu\text{M}$ , and labeled with 100  $\mu\text{Ci}$  of  $^{51}\text{Cr}$  (Amersham Corp., Arlington Heights, IL) for 1 hour after which they are washed four times with HBSS.

Cytolytic activity is determined in a standard 4-h, split well  $^{51}\text{Cr}$  release assay using U-bottomed 96 well plates containing 3,000 targets/well. Stimulated PBMC are tested at effector/target (E/T) ratios of 20-50:1 on day 14. Percent cytotoxicity is determined from the formula:  $100 \times [(\text{experimental release} - \text{spontaneous release}) / (\text{maximum release} - \text{spontaneous release})]$ . Maximum release is determined by lysis of targets by detergent (2% Triton X-100; Sigma Chemical Co., St. Louis, MO). Spontaneous release is <25% of maximum release for all experiments.

The results of such an analysis indicate the extent to which HLA-restricted CTL populations have been stimulated by previous exposure to HCV or an HCV vaccine.

The class II restricted HTL responses may also be analyzed. Purified PBMC are cultured in a 96-well flat bottom plate at a density of  $1.5 \times 10^5$  cells/well and are stimulated with 10  $\mu\text{g}/\text{ml}$  synthetic peptide, whole antigen, or PHA. Cells are routinely plated in replicates of 4-6 wells for each condition. After seven days of culture, the medium is removed and replaced with fresh medium containing 10U/ml IL-2. Two days later, 1  $\mu\text{Ci}$   $^3\text{H}$ -thymidine is added to each well and incubation is continued for an additional 18 hours. Cellular DNA is then harvested on glass fiber mats and analyzed for  $^3\text{H}$ -thymidine

incorporation. Antigen-specific T cell proliferation is calculated as the ratio of  $^3\text{H}$ -thymidine incorporation in the presence of antigen divided by the  $^3\text{H}$ -thymidine incorporation in the absence of antigen.

5 Example 18: Induction Of Specific CTL Response In Humans

A human clinical trial for an immunogenic composition comprising CTL and HTL epitopes of the invention is set up as an IND Phase I, dose escalation study and carried out as a randomized, double-blind, placebo-controlled trial. Such a trial is designed, for example, as follows:

10 A total of about 27 subjects are enrolled and divided into 3 groups:

Group I: 3 subjects are injected with placebo and 6 subjects are injected with 5  $\mu\text{g}$  of peptide composition;

Group II: 3 subjects are injected with placebo and 6 subjects are injected with 50  $\mu\text{g}$  peptide composition;

15 Group III: 3 subjects are injected with placebo and 6 subjects are injected with 500  $\mu\text{g}$  of peptide composition.

After 4 weeks following the first injection, all subjects receive a booster inoculation at the same dosage.

20 The endpoints measured in this study relate to the safety and tolerability of the peptide composition as well as its immunogenicity. Cellular immune responses to the peptide composition are an index of the intrinsic activity of this the peptide composition, and can therefore be viewed as a measure of biological efficacy. The following summarize the clinical and laboratory data that relate to safety and efficacy endpoints.

25 Safety: The incidence of adverse events is monitored in the placebo and drug treatment group and assessed in terms of degree and reversibility.

Evaluation of Vaccine Efficacy: For evaluation of vaccine efficacy, subjects are bled before and after injection. Peripheral blood mononuclear cells are isolated from fresh heparinized blood by Ficoll-Hypaque density gradient centrifugation, aliquoted in freezing media and stored frozen. Samples are assayed for CTL and HTL activity.

30 The vaccine is found to be both safe and efficacious.

Example 19: Phase II Trials In Patients Infected With HCV

Phase II trials are performed to study the effect of administering the CTL-HTL peptide compositions to patients having chronic HCV infection. The main objectives of

the trials are to determine an effective dose and regimen for inducing CTLs in chronically infected HCV patients, to establish the safety of inducing a CTL and HTL response in these patients, and to see to what extent activation of CTLs improves the clinical picture of chronically infected CTL patients, as manifested by a transient flare in alanine aminotransferase (ALT), normalization of ALT, and reduction in HCV DNA. Such a study is designed, for example, as follows:

The studies are performed in multiple centers. The trial design is an open-label, uncontrolled, dose escalation protocol wherein the peptide composition is administered as a single dose followed six weeks later by a single booster shot of the same dose. The dosages are 50, 500 and 5,000 micrograms per injection. Drug-associated adverse effects (severity and reversibility) are recorded.

There are three patient groupings. The first group is injected with 50 micrograms of the peptide composition and the second and third groups with 500 and 5,000 micrograms of peptide composition, respectively. The patients within each group range in age from 21-65, include both males and females, and represent diverse ethnic backgrounds. All of them are infected with HCV for over five years and are HIV, HBV and delta hepatitis virus (HDV) negative, but have positive levels of HCV antigen.

The magnitude and incidence of ALT flares and the levels of HCV DNA in the blood are monitored to assess the effects of administering the peptide compositions. The levels of HCV DNA in the blood are an indirect indication of the progress of treatment. The vaccine composition is found to be both safe and efficacious in the treatment of chronic HCV infection.

#### Example 20. Induction of CTL Responses Using a Prime Boost Protocol

A prime boost protocol can also be used for the administration of the vaccine to humans. Such a vaccine regimen may include an initial administration of, for example, naked DNA followed by a boost using recombinant virus encoding the vaccine, or recombinant protein/polypeptide or a peptide mixture administered in an adjuvant.

For example, the initial immunization may be performed using an expression vector, such as that constructed in Example 11, in the form of naked nucleic acid administered IM (or SC or ID) in the amounts of 0.5-5 mg at multiple sites. The nucleic acid (0.1 to 1000  $\mu$ g) can also be administered using a gene gun. Following an incubation period of 3-4 weeks, a booster dose is administered. The booster can, e.g., be recombinant fowlpox virus administered at a dose of  $5 \cdot 10^7$  to  $5 \cdot 10^9$  pfu. An alternative

recombinant virus, such as an MVA, canarypox, adenovirus, or adeno-associated virus, can also be used for the booster, or the polyepitopic protein or a mixture of the peptides can be administered. For evaluation of vaccine efficacy, patient blood samples will be obtained before immunization as well as at intervals following administration of the initial vaccine and booster doses of the vaccine. Peripheral blood mononuclear cells are isolated from fresh heparinized blood by Ficoll-Hypaque density gradient centrifugation, aliquoted in freezing media and stored frozen. Samples are assayed for CTL and HTL activity.

Analysis of the results will indicate that a magnitude of response sufficient to achieve protective immunity or to treat HCV infection is generated.

#### Example 21. Administration of Vaccine Compositions Using Dendritic Cells

Vaccines comprising peptide epitopes of the invention may be administered using dendritic cells. In this example, the peptide-pulsed dendritic cells can be administered to a patient to stimulate a CTL response *in vivo*. In this method dendritic cells are isolated, expanded, and pulsed with a vaccine comprising peptide CTL and HTL epitopes of the invention. The dendritic cells are infused back into the patient to elicit CTL and HTL responses *in vivo*. The induced CTL and HTL then destroy (CTL) or facilitate destruction (HTL) of the specific target HCV-infected cells that bear the proteins from which the epitopes in the vaccine are derived.

Alternatively, *Ex vivo* CTL or HTL responses to a particular tumor-associated antigen can be induced by incubating in tissue culture the patient's, or genetically compatible, CTL or HTL precursor cells together with a source of antigen-presenting cells, such as dendritic cells, and the appropriate immunogenic peptides. After an appropriate incubation time (typically about 7-28 days), in which the precursor cells are activated and expanded into effector cells, the cells are infused back into the patient, where they will destroy (CTL) or facilitate destruction (HTL) of their specific target cells, *i.e.*, tumor cells.

#### Example 22: Alternative Method of Identifying Motif-Bearing Peptides

Another way of identifying motif-bearing peptides is to elute them from cells bearing defined MHC molecules. For example, EBV transformed B cell lines used for tissue typing, have been extensively characterized to determine which HLA molecules they express. In certain cases these cells express only a single type of HLA molecule.

These cells can then be infected with a pathogenic organism, *e.g.*, HCV, or transfected with nucleic acids that express the antigen of interest. Thereafter, peptides produced by endogenous antigen processing of peptides produced consequent to infection (or as a result of transfection) will bind be displayed on the cell surface. The peptides are then  
5 eluted from the HLA molecules by exposure to mild acid conditions and their amino acid sequence determined, *e.g.*, by mass spectral analysis (*e.g.*, Kubo *et al.*, *J. Immunol.* 152:3913, 1994). Because, as disclosed herein, the majority of peptides that bind a particular HLA molecule are motif-bearing, this is an alternative modality for obtaining the motif-bearing peptides correlated with the particular HLA molecule expressed on the  
10 cell.

Alternatively, cell lines that do not express any endogenous HLA molecules can be transfected with an expression construct encoding a single HLA allele. These cells may then be used as described, *i.e.*, they may be infected with a pathogenic organism or transfected with nucleic acid encoding an antigen of interest to isolate peptides  
15 corresponding to the pathogen or antigen of interest that have been presented on the cell surface. Peptides obtained from such an analysis will bear motif(s) that correspond to binding to the single HLA allele that is expressed in the cell.

As appreciated by one in the art, one can perform a similar analysis on a cell bearing more than one HLA allele and subsequently determine peptides specific for each  
20 HLA allele expressed. Moreover, one of skill would also recognize that means other than infection or transfection, such as loading with a protein antigen, can be used to provide a source of antigen to the cell.

The above examples are provided to illustrate the invention but not to limit its scope. For example, the human terminology for the Major Histocompatibility Complex, namely HLA, is used throughout this document. It is to be appreciated that these  
25 principles can be extended to other species as well. Thus, other variants of the invention will be readily apparent to one of ordinary skill in the art and are encompassed by the appended claims. All publications, patents, and patent application cited herein are hereby incorporated by reference for all purposes.



TABLE I

SUPERMOTIFS	POSITION	POSITION	POSITION
	2 (Primary Anchor)	3 (Primary Anchor)	C Terminus (Primary Anchor)
A1	<b>T, I, L, V, M, S</b>		<b>F, W, Y</b>
A2	<b>L, I, V, M, A, T, Q</b>		<b>I, V, M, A, T, L</b>
A3	<b>V, S, M, A, T, L, I</b>		<b>R, K</b>
A24	<b>Y, F, W, I, V, L, M, T</b>		<b>F, I, Y, W, L, M</b>
B7	<b>P</b>		<b>V, I, L, F, M, W, Y, A</b>
B27	<b>R, H, K</b>		<b>F, Y, L, W, M, I, V, A</b>
B44	<b>E, D</b>		<b>F, W, L, I, M, V, A</b>
B58	<b>A, T, S</b>		<b>F, W, Y, L, I, V, M, A</b>
B62	<b>Q, L, I, V, M, P</b>		<b>F, W, Y, M, I, V, L, A</b>
MOTIFS			
A1	<b>T, S, M</b>		<b>Y</b>
A1		<b>D, E, A, S</b>	<b>Y</b>
A2.1	<b>L, M, V, Q, I, A, T</b>		<b>V, L, I, M, A, T</b>
A3	<b>L, M, V, I, S, A, T, F, C, G, D</b>		<b>K, Y, R, H, F, A</b>
A11	<b>V, T, M, L, I, S, A, G, N, C, D, F</b>		<b>K, R, Y, H</b>
A24	<b>Y, F, W, M</b>		<b>F, L, I, W</b>
A*3101	<b>M, V, T, A, L, I, S</b>		<b>R, K</b>
A*3301	<b>M, V, A, L, F, I, S, T</b>		<b>R, K</b>
A*6801	<b>A, V, T, M, S, L, I</b>		<b>R, K</b>
B*0702	<b>P</b>		<b>L, M, F, W, Y, A, I, V</b>
B*3501	<b>P</b>		<b>L, M, F, W, Y, I, V, A</b>
B51	<b>P</b>		<b>L, I, V, F, W, Y, A, M</b>
B*5301	<b>P</b>		<b>I, M, F, W, Y, A, L, V</b>
B*5401	<b>P</b>		<b>A, T, I, V, L, M, F, W, Y</b>

Bolded residues are preferred, italicized residues are less preferred: A peptide is considered motif-bearing if it has primary anchors at each primary anchor position for a motif or supermotif as specified in the above table.

TABLE II

	POSITION							
	1	2	3	4	5	6	7	8 C-terminus
<b>SUPERMOTIFS</b>								
A1		1° Anchor T,I,L,V,M,S						1° Anchor F,W,Y
A2		1° Anchor L,I,V,M,A, T,Q						1° Anchor L,I,V,M,A,T
A3	preferred	1° Anchor V,S,M,A,T, L,I	Y,F,W (4/5)		Y,F,W (3/5)		Y,F,W (4/5)	1° Anchor R,K
	deleterious	D,E (3/5); P (5/5)	D,E (4/5)					
A24		1° Anchor Y,F,W,I,V, L,M,T						1° Anchor F,I,Y,W,L,M
B7	preferred	F,W,Y (5/5) L,I,V,M (3/5)	1° Anchor P	F,W,Y (4/5)			F,W,Y (3/5)	1° Anchor V,I,L,F,M,W,Y,A
	deleterious	D,E (3/5); P (5/5); G (4/5); A (3/5); Q,N (3/5)			D,E (3/5)	G (4/5)	Q,N (4/5)	D,E (4/5)
B27		1° Anchor R,H,K						1° Anchor F,Y,L,W,M,V,A
B44		1° Anchor E,D						1° Anchor F,W,Y,L,I,M,V,A
B58		1° Anchor A,T,S						1° Anchor F,W,Y,L,I,V,M,A
B62		1° Anchor Q,L,I,V,M, P						1° Anchor F,W,Y,M,I,V,L,A

## POSITION

	1	2	3	4	5	6	7	8	C-terminus
A1 preferred 9-mer	G,F,Y,W	I°Anchor S,T,M	D,E,A	Y,F,W		P	D,E,Q,N	Y,F,W	I°Anchor Y
	deleterious	D,E	R,H,K,L,I,V M,P	A	G	A			

## MOTIFS

A1 preferred 9-mer	G,R,H,K	A,S,T,C,L,I V,M,	I°Anchor D,E,A,S	G,S,T,C		A,S,T,C	L,I,V,M	D,E	I°Anchor Y
	deleterious	A	R,H,K,D,E, P,Y,F,W	D,E	P,Q,N	R,H,K	P,G	G,P	

POSITION									
	1	2	3	4	5	6	7	8	9 or C-terminus
A1 preferred	Y,F,W	<u>1°Anchor</u> S,T,M	D,E,A,Q,N	A	Y,F,W,Q,N		P,A,S,T,C	G,D,E	P
deleterious	G,P		R,H,K,G,L,I V,M	D,E	R,H,K	Q,N,A	R,H,K,Y,F, W	R,H,K	A
A1 preferred	Y,F,W	S,T,C,L,I,V M	<u>1°Anchor</u> D,E,A,S	A	Y,F,W		P,G	G	Y,F,W
deleterious	R,H,K	R,H,K,D,E, P,Y,F,W			P	G		P,R,H,K	Q,N
A2.1 preferred	Y,F,W	<u>1°Anchor</u> L,M,I,V,Q, A,T	Y,F,W	S,T,C	Y,F,W		A	P	<u>1°Anchor</u> V,L,I,M,A,T
deleterious	D,E,P		D,E,R,K,H			R,K,H	D,E,R,K,H		
A2.1 preferred	A,Y,F,W	<u>1°Anchor</u> L,M,I,V,Q, A,T	L,V,I,M	G		G		F,Y,W, L,V,I,M	<u>1°Anchor</u> V,L,I,M,A,T
deleterious	D,E,P		D,E	R,K,H,A	P		R,K,H	D,E,R, K,H	R,K,H

POSITION									
	1	2	3	4	5	6	7	8	9 or C-terminus
A3	preferred R,H,K	1°Anchor L,M,V,I,S, A,T,F,C,G D	Y,F,W	P,R,H,K,Y, F,W	A	Y,F,W		P	C-terminus 1°Anchor K,Y,R,H,F,A
	deleterious D,E,P		D,E						
A11	preferred A	1°Anchor V,T,L,M,I, S,A,G,N,C, D,F	Y,F,W	Y,F,W	A	Y,F,W	Y,F,W	P	1°Anchor K,,R,Y,H
	deleterious D,E,P						A	G	
A24 9-mer	preferred Y,F,W,R,H,K	1°Anchor Y,F,W,M		S,T,C			Y,F,W	Y,F,W	1°Anchor F,L,I,W
	deleterious D,E,G		D,E	G	Q,N,P	D,E,R,H,K	G	A,Q,N	
A24 10-mer	preferred	1°Anchor Y,F,W,M		P	Y,F,W,P		P		1°Anchor F,L,I,W
	deleterious		G,D,E	Q,N	R,H,K	D,E	A	Q,N	D,E,A
A3101	preferred R,H,K	1°Anchor M,V,T,A,L, I,S	Y,F,W	P		Y,F,W	Y,F,W	A,P	1°Anchor R,K
	deleterious D,E,P		D,E		A,D,E	D,E	D,E	D,E	

## POSITION

	1	2	3	4	5	6	7	8	9 or C-terminus	C-terminus 1°Anchor R,K
A3301 preferred		1°Anchor M,V,A,L,F, I,S,T	Y,F,W				A,Y,F,W			
deleterious	G,P		D,E							
A6801 preferred	Y,F,W,S,T,C	1°Anchor A,V,T,M,S, L,I			Y,F,W,L,I, V,M		Y,F,W	P		1°Anchor R,K
deleterious	G,P		D,E,G		R,H,K			A		
B0702 preferred	R,H,K,F,W,Y	1°Anchor P	R,H,K		R,H,K	R,H,K	R,H,K	P,A		1°Anchor L,M,F,W,Y,A, I,V
deleterious	D,E,Q,N,P		D,E,P	D,E	D,E	G,D,E	Q,N	D,E		
B3501 preferred	F,W,Y,L,I,V,M	1°Anchor P	F,W,Y				F,W,Y			1°Anchor L,M,F,W,Y,I, V,A
deleterious	A,G,P				G	G				

POSITION									
	1	2	3	4	5	6	7	8	9 or C-terminus
B51 preferred	L,I,V,M,F,W,Y	<u>1°Anchor</u> P	F,W,Y	S,T,C	F,W,Y		G	F,W,Y	C-terminus <u>1°Anchor</u> L,I,V,F,W, Y,A,M
deleterious	A,G,P,D,E,R,H,K, S,T,C				D,E	G	D,E,Q,N	G,D,E	
B5301 preferred	L,I,V,M,F,W,Y	<u>1°Anchor</u> P	F,W,Y	S,T,C	F,W,Y		L,I,V,M,F, W,Y	F,W,Y	<u>1°Anchor</u> I,M,F,W,Y, A,L,V
deleterious	A,G,P,Q,N					G	R,H,K,Q,N	D,E	
B5401 preferred	F,W,Y	<u>1°Anchor</u> P	F,W,Y,I,L,V M		L,I,V,M		A,L,I,V,M	F,W,Y,A,P	<u>1°Anchor</u> A,T,I,V,L, M,F,W,Y
deleterious	G,P,Q,N,D,E		G,D,E,S,T,C		R,H,K,D,E	D,E	Q,N,D,G,E	D,E	

Italicized residues indicate less preferred or "tolerated" residues.  
The information in Table II is specific for 9-mers unless otherwise specified.

Table III

POSITION									
MOTIFS	<u>1° anchor 1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>1° anchor 6</u>	<u>7</u>	<u>8</u>	<u>9</u>
DR4 preferred	F, M, Y, L, I, V, W	M	T		I	V, S, T, C, P, A, L, I, M	M, H, R,		M, H
deleterious				W,					W, D, E
DR1 preferred	M, F, L, I, V, W, Y			P, A, M, Q		V, M, A, T, S, P, L, I, C	M,		A, V, M
deleterious		C	C, H	F, D	C, W, D		G, D, E, D		
DR7 preferred	M, F, L, I, V, W, Y	M	W	A		I, V, M, S, A, C, T, P, L	M		I, V
deleterious		C,		G,			G, R, D N		G
DR Supermotif	M, F, L, I, V, W, Y					V, M, S, T, A, C, P, L, I			
DR3 MOTIFS	<u>1° anchor 1</u>	<u>2</u>	<u>3</u>	<u>1° anchor 4</u>	<u>5</u>	<u>1° anchor 6</u>			
motif a preferred	L, I, V, M, F, Y			D					
motif b preferred	L, I, V, M, F, A, Y			D, N, Q, E, S, T		K, R, H			

Italicized residues indicate less preferred or "tolerated" residues.



**Table IV: HLA Class I Standard Peptide Binding Affinity.**

ALLELE	STANDARD PEPTIDE	SEQUENCE (SEQ ID NO:)	STANDARD BINDING AFFINITY (nM)
A*0101	944.02	YLEPAIAKY	25
A*0201	941.01	FLPSDYFPSV	5.0
A*0202	941.01	FLPSDYFPSV	4.3
A*0203	941.01	FLPSDYFPSV	10
A*0205	941.01	FLPSDYFPSV	4.3
A*0206	941.01	FLPSDYFPSV	3.7
A*0207	941.01	FLPSDYFPSV	23
A*6802	1072.34	YVIKVSARV	8.0
A*0301	941.12	KVFPYALINK	11
A*1101	940.06	AVDLYHFLK	6.0
A*3101	941.12	KVFPYALINK	18
A*3301	1083.02	STLPETYVRR	29
A*6801	941.12	KVFPYALINK	8.0
A*2402	979.02	AYIDNYNKF	12
B*0702	1075.23	APRTLVL	5.5
B*3501	1021.05	FPFKYAAAF	7.2
B51	1021.05	FPFKYAAAF	5.5
B*5301	1021.05	FPFKYAAAF	9.3
B*5401	1021.05	FPFKYAAAF	10

**Table V. HLA Class II Standard Peptide Binding Affinity.**

Allele	Nomenclature	Standard Peptide	Sequence (SEQ ID NO:)	Binding Affinity (nM)
DRB1*0101	DR1	515.01	PKYVKQNTLKLAT	5.0
DRB1*0301	DR3	829.02	YKTIAFDEEARR	300
DRB1*0401	DR4w4	515.01	PKYVKQNTLKLAT	45
DRB1*0404	DR4w14	717.01	YARFQSQTTLKQKT	50
DRB1*0405	DR4w15	717.01	YARFQSQTTLKQKT	38
DRB1*0701	DR7	553.01	QYIKANSKFIGITE	25
DRB1*0802	DR8w2	553.01	QYIKANSKFIGITE	49
DRB1*0803	DR8w3	553.01	QYIKANSKFIGITE	1600
DRB1*0901	DR9	553.01	QYIKANSKFIGITE	75
DRB1*1101	DR5w11	553.01	QYIKANSKFIGITE	20
DRB1*1201	DR5w12	1200.05	EALIHQLKINPYVLS	298
DRB1*1302	DR6w19	650.22	QYIKANAKFIGITE	3.5
DRB1*1501	DR2w2 $\beta$ 1	507.02	GRTQDENPVVHFFKNIV TPRTPPP	9.1
DRB3*0101	DR52a	511	NGQIGNDPNRDIL	470
DRB4*0101	DRw53	717.01	YARFQSQTTLKQKT	58
DRB5*0101	DR2w2 $\beta$ 2	553.01	QYIKANSKFIGITE	20

Table VI

HLA-supertype	Allele-specific HLA-supertype members	
	Verified <sup>a</sup>	Predicted <sup>b</sup>
A1	A*0101, A*2501, A*2601, A*2602, A*3201	A*0102, A*2604, A*3601, A*4301, A*8001
A2	A*0201, A*0202, A*0203, A*0204, A*0205, A*0206, A*0207, A*0209, A*0214, A*6802, A*6901	A*0208, A*0210, A*0211, A*0212, A*0213
A3	A*0301, A*1101, A*3101, A*3301, A*6801	A*0302, A*1102, A*2603, A*3302, A*3303, A*3401, A*3402, A*6601, A*6602, A*7401
A24	A*2301, A*2402, A*3001	A*2403, A*2404, A*3002, A*3003
B7	B*0702, B*0703, B*0704, B*0705, B*1508, B*3501, B*3502, B*3503, B*3503, B*3504, B*3505, B*3506, B*3507, B*3508, B*5101, B*5102, B*5103, B*5104, B*5105, B*5301, B*5401, B*5501, B*5502, B*5601, B*5602, B*6701, B*7801	B*1511, B*4201, B*5901
B27	B*1401, B*1402, B*1509, B*2702, B*2703, B*2704, B*2705, B*2706, B*3801, B*3901, B*3902, B*7301	B*2701, B*2707, B*2708, B*3802, B*3903, B*3904, B*3905, B*4801, B*4802, B*1510, B*1518, B*1503
B44	B*1801, B*1802, B*3701, B*4402, B*4403, B*4404, B*4001, B*4002, B*4006	B*4101, B*4501, B*4701, B*4901, B*5001
B58	B*5701, B*5702, B*5801, B*5802, B*1516, B*1517	
B62	B*1501, B*1502, B*1513, B*5201	B*1301, B*1302, B*1504, B*1505, B*1506, B*1507, B*1515, B*1520, B*1521, B*1512, B*1514, B*1510

- a. Verified alleles include alleles whose specificity has been determined by pool sequencing analysis, peptide binding assays, or by analysis of the sequences of CTL epitopes.
- b. Predicted alleles are alleles whose specificity is predicted on the basis of B and F pocket structure to overlap with the supertype specificity.

Table VII

## HCV A01 Super Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0101
ATGNLPGCSF	165	10	13	93	
ATLGFAY	1265	8	14	100	
AVQWMNRLIAF	1917	11	14	100	
CTCGSSQLY	1128	9	11	79	0.3700
CTRGVAKVDF	1190	11	11	79	
CTWMNSTGF	555	9	11	79	
CVTOTVDF	1462	8	12	88	
DLEWISTW	1857	9	12	86	
EITMRSPVF	1207	9	12	86	
FSYDTRCF	2870	8	11	79	
FTAMITRY	2792	8	14	100	
FTGLTHDAHF	1567	11	13	93	
GLPVOCHLEF	1552	11	12	86	
GLSAFSLHSY	2821	10	11	79	0.0029
GLTHIDAHF	1569	9	13	93	
GSSYGFQY	2841	8	11	79	
GTFPINAY	2063	8	11	79	
GVAGALVAF	1863	9	12	86	
GVAKAVDF	1193	8	11	79	
GVLAALAAV	1670	9	12	86	
GVRVCEKMAV	2619	11	14	100	
GVRVLEQWNY	154	11	12	86	
HLHQIMOVQY	696	11	11	79	
HMMNFRSGQY	1769	11	13	93	
HMGEGEAVQW	1910	11	11	79	
IMAKNEVF	2591	8	12	86	
ITYSTYQKF	1296	8	12	86	
NDVQVLY	701	8	12	86	
KSTKVPAAV	1241	9	12	86	0.0130
KVIDTLTQGF	121	10	12	86	
LIEANLW	2235	8	12	86	
LINTNGSW	414	8	11	79	
LLAPITAY	1030	8	14	100	
LLFNILGGW	1812	9	12	86	
LSPRGSRPSW	97	11	11	79	0.8100
LSAFSLHSY	2922	9	11	79	
LSPRGSRPSW	98	10	11	79	
LTCGFADLMGY	126	11	12	86	
LTTHDAHF	1570	8	13	93	
LVDLIAGY	1853	8	11	79	
MILMTHFF	2878	8	12	86	
NIVDQVLY	700	9	12	86	0.0980
NLPGCSFIF	168	10	13	93	
NTCVTQTVDF	1480	10	12	86	
NTNRRPOQWKF	14	11	11	79	

## HCV AD1 Super Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0101
NQDQVGW	1108	9	11	79	
PITYSTYQKF	1295	10	11	79	
PNQFSYOTRCF	2667	11	11	79	
PSVAATLGF	1281	9	14	100	
PTLHGPTPLLY	1621	11	11	79	
PVOOCHLEF	1554	9	12	86	
PVOOCHLEW	1554	10	12	86	
QTVQFSLDPTF	1465	11	12	86	
RJHGLSAF	2918	8	12	86	
RLAPITAY	1029	9	12	86	
RMAWDMMNNW	317	10	12	86	
RMILMTIF	2875	8	12	86	
RMILMTIFF	2875	9	12	86	
RVCEKVALY	2621	8	14	100	
RMLDXVNY	156	9	12	86	
STKVPAAV	1242	8	12	86	
SVAATLGF	1262	8	14	100	
SVAATLGFQAY	1262	11	14	100	
TIMAKNEVF	2590	9	11	79	0.0300
TLHGPTPLLY	1622	10	11	79	
TLLNLGGW	1811	10	12	86	
TTIMAKNEVF	2509	10	11	79	
TTMRSPVF	1208	8	12	86	
TVDPSLDPTF	1466	10	12	86	
VIDTLTQGF	122	9	12	86	
VLAALAY	1871	8	12	86	
VLEDGVNY	167	8	12	86	
VLVDILAGY	1852	9	11	79	
VN3SSYGF	2639	8	11	79	
VN3SSYGFQY	2639	10	11	79	
WMNRLIAF	1820	8	14	100	
YSPGQRYEF	2648	9	11	79	
YTNVDQDLVGW	1106	11	11	79	
YVGDLCGSVF	276	10	12	86	

79

Table VIII  
HCV A02 Super Motif with Binding Information

Conservancy	Freq.	Position	Sequence	A*0201	A*0202	A*0203	A*0206	A*0802
93	13	1904	ANLRHHV					
86	12	1673	AALAAYCL					
79	11	1250	AAQGYKVL					
79	11	1250	AAQGYKVLV					
79	11	1250	AAQGYKVLVL					
79	11	147	AARALAHGV					
79	11	147	AARALAHGVRV					
100	14	1264	AATLGFGA					
93	13	1264	AATLGFGAYM					
86	12	1187	AVCTRGV					
79	11	1187	AVCTRGVA					
79	11	1187	AVCTRGVAKA					
93	13	1890	AILSPGAL	0.0014				
86	12	1880	AILSPGALV	0.0035				
85	12	1880	AILSPGALVV					
100	14	150	ALAHGVRV	0.0037				
100	14	150	ALAHGVRVL					
86	12	1737	ALGLDTA					
86	12	689	ALSTGLHL					
79	11	1896	ALVGVWCA					
79	11	1896	ALVGVWCAV					
78	11	1898	ALVGVWCAAI					
86	12	1602	AOAPPSWDCM					
79	11	1251	AQGYKVL	0.0160	0.0008	0.2200	0.0002	0.0039
79	11	1251	AQGYKVLV	0.0010				
86	12	77	AQGYKVLVL					
86	12	77	AQGYKVLVL					
93	13	1285	ATLFGAYM					
79	11	1354	ATPGSVT					
79	11	1596	ATVCARAGA					
100	14	1419	AVAYYRGL	0.0002				
100	14	1419	AVAYYRGLDV					
79	11	1188	AVCTRGVA					
79	11	1188	AVCTRGVAKA					
79	11	1188	AVCTRGVAKAV					
100	14	1917	AVCHHNTL					
100	14	1917	AVQWNNRLI	0.0001				
100	14	1917	AVQWNNRLIA					
93	13	1903	CAAILRHHV					
79	11	1530	CAWYELTPA					
86	12	2941	CLFKLGVPPL	0.0002				
86	12	739	CLWNNILLI					
79	11	1653	CMSADLEV					

## HCY A02 Super Motif with Binding Information

Conservancy	Freq.	Position	Sequence	A*0201	A*0202	A*0203	A*0206	A*6802
79	11	1653	CMSADLEW	0.0067				
79	11	1653	CMSADLEVVT					
79	11	1128	CTCGSSDL					
79	11	1128	CTCGSSDLYL					
78	11	1126	CTCGSSDLYLV					
79	11	1190	CTRGVAKA					
79	11	1190	CTRGVAKAV					
79	11	555	CTWMNSTGFT	0.0006				
86	12	1462	CVTQTVDFSL					
79	11	1527	DAGCAWYEL					
100	14	1574	DAHFLSOT					
86	12	1855	DILAGYGA	0.0002				
79	11	1855	DILAGYGAGV					
79	11	1855	DILAGYGAGVA					
86	12	279	DLCGSVFL	0.0007				
79	11	279	DLCGSVFLV					
86	12	1657	DLEVSTST	0.0002				
86	12	1657	DLEVSTSTW					
86	12	1657	DLEVSTSTWL					
93	13	2617	-DLGVRVGEKMA-					
93	13	2617	DLGVRVCEKMA					
79	11	132	DLMGYIPL	0.0530	0.0009	0.0480	0.0077	3.3000
79	11	132	DLMGYIPLV					
79	11	132	DLMGYIPLVGA	0.0008				
79	11	2412	DLSDGSWST					
79	11	2412	DLSDGSWSTV					
79	11	1883	DLNLLPA					
79	11	1883	DLNLLPAI					
79	11	1883	DLNLLPAIL					
79	11	2772	DLWICESA					
86	12	1134	DLYLVTRHA					
86	12	1134	DLYLVTRHADV					
86	12	321	DMMNWSPT	0.0001				
86	12	1339	DOAETAGA	0.0001				
86	12	1339	DOAETAGATL					
86	12	1339	DOAETAGARLV					
86	12	994	DTAACGDI					
86	12	994	DTAACGDII					
86	12	124	DTLTCGFA					
86	12	124	DTLTCGFADL					
86	12	124	DTLTCGFADLM					
93	13	2673	DTNCFDST	0.0001				

## HCV AD2 Super Motif with Binding Information

Conservancy	Freq.	Position	Sequence	A*0201	A*0202	A*0203	A*0206	A*8B02
93	13	2673	DTRCFDSTV					
93	13	2673	DTRCFDSTV					
86	12	21	DVRFPGGQI	0.0001				
86	12	21	DVRFPGGQI					
79	11	750	EALENLV					
100	14	2794	EAMTRYSA					
86	12	2237	EANLLWROEM					
93	13	1377	EIPFYGKA					
93	13	1377	EIPFYGKA	0.0001				
100	14	2814	ELITSCSSNV	0.0002				
79	11	666	ELSPILLST					
79	11	666	ELSPILLST					
86	12	2245	EMGNITRV	0.0003				
86	12	1731	EOROKAL					
85	12	1731	EOROKALGL					
86	12	1731	EOROKALGLL					
86	12	1342	ETAGARLV					
86	12	1342	ETAGARLV					
86	12	1342	ETAGARLVWL					
86	12	1342	ETAGARLVLA					
86	12	1207	ETIMRSPV					
86	12	1207	ETIMRSPVFT					
88	12	1659	EVTSTW	0.0001				
86	12	1659	EVTSTWVL	0.0004				
86	12	1659	EVTSTWVL					
93	13	130	FADLMGYI					
79	11	130	FADLMGYIPL					
79	11	130	FADLMGYIPL					
100	14	1927	FASRGNV					
86	12	1927	FASRGNVSP					
100	14	1773	FISGQIYL	0.1000				
100	14	1773	FISGQIYLA					
100	14	1773	FISGQIYLA					
79	11	1304	FLADGGCSGA					
86	12	177	FLALLSCL	0.0048				
86	12	177	FLALLSCL					
86	12	177	FLALLSCL					
93	13	728	FLLLADARV					
86	12	1228	FOVAHLHA					
86	12	1228	FOVAHLHAPT					
78	11	2645	FOYSPGORV					
100	14	2782	FTEAMTRYSA	0.2800	0.0480	0.0670	0.0150	0.3600
93	13	1587	FTGLTHIDA					



## HCV A92 Super Motif with Binding Information

Conservancy	Freq.	Position	Sequence	A*0201	A*0202	A*0203	A*0206	A*8802
93	13	512	FTPSPVV					
93	13	512	FTPSPVWGT					
93	13	512	FTPSPVWGT					
79	11	684	FTLPALST					
79	11	684	FTLPALSTGL					
79	11	146	GAARALAHGV					
86	12	992	GAOTAAACGDI					
86	12	992	GAOTAAACGDI					
86	12	1861	GAGVAGAL					
88	12	1861	GAGVAGALV					
86	12	1861	GAGVAGALVA					
86	12	350	GAHWGVLA					
79	11	1895	GALVGVW					
79	11	1895	GALVGVWCA					
79	11	1895	GALVGVWCAA					
86	12	1345	GARLVLA					
79	11	1345	GARLVLAT					
78	11	1345	GARLVLATA					
79	11	1345	GARLVVLATAT					
100	14	1916	GAVQWMTL					
100	14	1916	GAVQWMTL					
100	14	1916	GAVQWMTL					
100	14	1333	GIGTVLDOA					
100	14	1333	GIGTVLDOA					
100	14	1333	GIGTVLDOAET					
100	14	1776	GIGYLAGL					
100	14	1776	GIGYLAGIST					
100	14	1776	GIGYLAGLSTL					
79	11	1425	GLDVS/PT					
93	13	1552	GLPVOCDL					
79	11	968	GLRDLAVA					
79	11	968	GLRDLAVAV					
100	14	1782	GLSTLPNPA					
79	11	1782	GLSTLPNPAI					
93	13	1589	GLTHDAHFL					
93	13	26	GGRCCVTL					
93	13	26	GONGGVIL					
79	11	2063	GTFPINAYT					
79	11	2063	GTFPINAYT					
100	14	1335	GTVLDNAET					
100	14	1335	GTVLDNAETA					
86	12	1863	GVAGALVA					
79	11	1081	GVQWTVYHGA					

0.0001

0.0001

0.0034

0.0007

## ILCV A02 Super Motif with Binding Information

Conservancy	Freq.	Position	Sequence	A*0201	A*0202	A*0203	A*0208	A*6802
86	12	1670	GVLAALAA					
86	12	1670	GVLAALAYCL					
79	11	161	GVNYATGNL	0.0001				
86	12	45	GVRATRTK					
100	14	2619	GVRCEKM					
100	14	2619	GVRCEKMA					
100	14	2619	GVRCEKMA	0.0002				
83	13	154	GVRCEKMA	0.0001				
79	11	1900	GVRLEDGV					
100	14	1234	GVVCAAIL					
100	14	1234	HAPTSGKST					
100	14	1572	HDAHFLSQT					
86	12	696	HLIQNVIV	0.0100	0.0014	0.5400	0.0027	0.0037
79	11	1719	HLPIEQQM					
93	13	1769	HMMNFISG					
79	11	698	IQNVIVVYL	0.3300	0.0004	0.1300	0.0280	0.0053
79	11	222	HTPGCVPCV					
86	12	2855	HTPVNSWL					
86	12	2855	HTPVNSWLGN					
79	11	1810	HVGPGECA					
79	11	1910	HVSPTHV					
86	12	1933	HVSPTHV					
100	14	1925	IAPSTGNHV					
79	11	1858	ILAGYGAGV	0.0430	0.0300	2.0000	0.0048	0.0450
79	11	1858	ILAGYGAGV	0.0002				
86	12	1816	ILGGWVA					
86	12	1816	ILGGWVAQL	0.0430	0.0024	0.0190	0.0005	0.0038
86	12	1816	ILGGWVAQL					
86	12	1331	ILGIGTVL					
86	12	1331	ILGIGTVLDOA					
93	13	1891	ILSPGALV					
93	13	1891	ILSPGALV					
93	13	1891	ILSPGALV					
93	13	1891	ILSPGALVGV					
79	11	2591	IMAKNEVFCV	0.0210	0.0004	0.3700	0.0036	0.0130
100	14	1777	IQYLAGLST	0.0088				
100	14	1777	IQYLAGLSTL					
86	12	2250	ITRVESENKV					
86	12	2250	ITRVESENKV					
100	14	2816	ITSCSSNV					
100	14	2816	ITSCSSNVSV					
100	14	2816	ITSCSSNVSV					
86	12	969	ITWGADTA					
86	12	969	ITWGADTAA					

HCY A02 Super Motif with Binding Information

Conservancy	Freq.	Position	Sequence	A*0201	A*0202	A*0203	A*0206	A*6802
79	11	1296	ITYSTYCKFL					
79	11	1296	ITYSTYCKFLA					
79	11	2613	NFPDLGV					
79	11	2613	NFPDLGVRV	0.0016				
93	13	30	NGGVYLL					
86	12	1738	KALGILLOT					
86	12	1738	KALGILLOTA					
86	12	2625	KMALYDVV					
86	12	1734	KOKALGIL					
86	12	1734	KOKALGILLOT					
86	12	1734	KOKALGILLOTA					
86	12	1734	KVIDTLTCGFA					
86	12	121	KVLVNPSV	0.0048				
100	14	1255	KVLVNPSVA					
100	14	1255	KVLVNPSVAA					
100	14	1255	KVPAAYAA	0.0011				
79	11	1244	LAALAYCL					
86	12	1872	LADGCSGGA					
79	11	1305	LAOFKOKA					
86	12	1729	LAOFKOKAL					
86	12	1729	LAGYGAGV					
79	11	1857	LAGYGAGVA					
79	11	1857	LAGYGAGVAGA					
79	11	1857	LAHGRVL					
100	14	151	LALLSCLT					
86	12	179	LAVAVEPV					
79	11	972	LWAFASRGNHV					
100	14	1924	LITSCSSNV	0.0004				
100	14	2815	LITSCSSNVSV					
100	14	2815	LWFFDLGV	0.0002				
79	11	2612	LWFFDLGVRV					
79	11	2612	LLALLSCL					
86	12	178	LLALLSCLT					
86	12	178	LLALLSCLT	0.0230	0.0150	0.0220	0.0011	0.0130
100	14	726	LLFLLADA					
93	13	726	LLFLLADARV					
86	12	1812	LLFLLGGW	1.2000	0.0380	3.1000	0.1900	1.2000
86	12	1812	LLFLLGGWA					
93	13	729	LLADARV	0.0061				
93	13	1887	LLPAILSPGA					
93	13	1887	LLPAILSPGAL	0.0025				
83	13	36	LUPRRGPRL					
83	13	36	LUPRRGPRLGV					

HCV A02 Super Motif with Binding Information

Conservancy	Freq.	Position	Sequence	A*0201	A*0202	A*0203	A*0206	A*8802
86	12	2240	LLWREMGNN					
93	13	1629	LLYRLGAV					
79	11	133	LMGYRLV					
79	11	133	LMGYPLVGA					
86	12	2761	LOOCTMLV					
86	12	126	LTCGFADL					
86	12	126	LTCGFADLM					
100	14	2180	LTDP SHIT					
100	14	2180	LTDP SHITA					
88	12	1052	LTGRDKNV					
93	13	1570	LTIDAHFL					
93	13	2176	LTSMLTDP SHI					
79	11	2738	LTTSCGNT					
79	11	2738	LTTSCGNLT					
79	11	2738	LTTSCGNLT					
88	12	1591	LVAYQATV	0.0002				
88	12	1591	LVAYQATVGA	-0.0001				
79	11	1853	LVDLAGYGA					
86	12	1867	LVGGVLA	0.0003				
86	12	1867	LVGGVLAAL					
86	12	1867	LVGGVLAALA					
100	14	1257	LVLPNSVA					
100	14	1257	LVLPNSVAA					
100	14	1257	LVLPNSVAAT					
100	14	1257	LVLPNSVAATL					
79	11	1884	LVNLLPAI	0.0002				
79	11	1884	LVNLLPAIL					
86	12	1137	LVTRHADV	0.0001				
79	11	1137	LVTRHADVI					
79	11	1137	LVTRHADVIPV					
79	11	1897	LVGVVCA					
79	11	1897	LVGVVCAA					
79	11	1897	LVGVVCAAI					
79	11	1897	LVGVVCAAIL	0.0011				
79	11	2773	LVVCEA					
86	12	1348	LVVLATAT					
86	12	2582	LVVNEVFCV	0.0022				
100	14	2179	MLTDP SHI					
100	14	2179	MLTDP SHIT	0.0002				
100	14	2179	MLTDP SHITA					
93	13	322	MMNNWSPT					

## HCY A02 Super Motif with Binding Information

Conservancy	Freq.	Position	Sequence	A*0201	A*0202	A*0203	A*0206	A*6802
93	13	1418	NAVAYYRGL					
93	13	1418	NAVAYYRGLOV					
86	12	2068	NAYTTGPCT					
86	12	1815	NILGGWA					
86	12	1815	NILGGWAA					
86	12	1815	NILGGWAAQL					
93	13	1282	NIRTVRT	0.0001				
79	11	1282	NIRTVRTI					
79	11	1282	NIRTVRTIT					
79	11	1282	NIRTVRTITT					
86	12	2249	NITRVESENKV					
86	12	700	NVDVOYL					
86	12	118	NLGKVIDT	0.0006				
86	12	118	NLGKVIDTL					
86	12	118	NLGKVIDTLT					
93	13	1888	NLLPAILSPGA					
86	12	2239	NLIWFOEM					
93	13	168	NLPGCCFSI	0.0041				
93	13	168	NLPGCCFSIL					
86	12	1480	NTCVTOTV					
93	13	416	NTNGSWHI					
86	12	14	NTNRTPODV					
93	13	1889	PAILSPGA					
93	13	1889	PAILSPGAL					
86	12	1889	PAILSPGALV					
86	12	1889	PAILSPGALVV					
86	12	588	PALSTGLI					
86	12	688	PALSTGLJHL					
79	11	2609	PARLVFPDL					
79	11	2066	PINAYTTGPCT					
79	11	1295	PITYSTYGRFL					
93	13	2403	PLEGEPGDFL					
79	11	143	PLGGAARA	0.0001				
79	11	143	PLGGAARAL					
79	11	143	PLGGAATALA					
93	13	1628	PLYRLGA	0.0001				
93	13	1628	PLYRLGAV					
79	11	2667	PMGFSYDT					
79	11	2807	POPEYOLEL					
79	11	2807	POPEYOLEU					
79	11	2807	POPEYOLEUT					
93	13	7	POKTKRNT					

## HCV A02 Super Motif with Binding Information

Conservancy	Freq.	Position	Sequence	A*0201	A*0202	A*0203	A*0206	A*6802
86	12	109	PTDPRRSRL					
79	11	1473	PTFTIET					
79	11	1473	PTFTIETT					
100	14	1238	PTSGKST					
93	13	1236	PTSGKSTKV					
86	12	1936	PTHVVPESDA					
86	12	1936	PTHVVPESDMA					
79	11	1621	PTLHGPTPL					
79	11	1621	PTLHGPTPL					
79	11	2870	PTLWARM					
79	11	2870	PTLWARMIL					
79	11	2870	PTLWARMILM					
79	11	2870	PTLWARMILMT					
100	14	1628	PTLLYRL					
93	13	1826	PTLLYRLGA					
93	13	1826	PTLLYRLGAV					
100	14	2857	PVNSWLGNI	0.0001				
100	14	2857	PVNSWLGNI	0.0001				
86	12	2857	PVNSWLGNIIM					
79	11	2018	PVHGCP					
93	13	508	PVYCFPSFV	0.0004				
93	13	508	PVYCFPSFW					
86	12	1340	QAETAGARL					
86	12	1340	QAETAGARLV					
88	12	1340	QAETAGARLV					
86	12	1603	QAPPSWDOM					
93	13	1595	QATVCANIA					
79	11	1595	QATVCARAQA					
93	13	29	QINGGVYL	0.0015				
93	13	29	QINGGVYL					
86	12	338	QLLRPOA					
86	12	2184	QLPCEPEPV					
79	11	2210	QLSAPSLKA	0.0002				
79	11	2210	QLSAPSLKAT					
86	12	1455	QVDFSLDT					
86	12	1229	QVAHLHPT					
86	12	1186	RAAVCTRGV					
79	11	1186	RAAVCTRGVA					
100	14	149	RALAHGVRV	0.0001				
100	14	149	RALAHGVRVL					
86	12	2733	RASGYLT					
79	11	43	RLGVRATRK					

## HCV A02 Super Motif with Binding Information

Conservancy	Freq.	Position	Sequence	A*0201	A*0202	A*0203	A*0206	A*5802
78	11	2918	RLHGLSAFSL					
79	11	2811	RLNFPDL	0.0280	0.0055	0.0180	0.0002	0.0032
79	11	2611	RLNFPDLGV	0.0890	0.0110	1.0000	0.0100	0.0050
79	11	1618	RLKPTLHGFT					
86	12	1029	RLAPITA					
86	12	1347	RLVLATA					
86	12	1347	RLVVLATAT					
100	14	619	RLWHYPCT					
86	12	317	RLAWDMMIM					
93	13	635	RLMYGSGVEHL					
86	12	2243	RLQEMGGNI					
86	12	2243	RLQEMGGNIT					
86	12	2243	RLQEMGGNITRV					
79	11	1284	RLTGVRITIT					
79	11	1284	RLTGVRITIT					
100	14	2621	RLVCEKXIAL					
86	12	2621	RLVCEKXIALYDV					
86	12	2252	RLVSEKIV	0.0001				
86	12	2252	RLVSEKIV					
79	11	2100	RLVGDRIY					
86	12	156	RLVLEDGVNYA					
86	12	156	RLVLEDGVNYAT					
86	12	2833	RLVYLLTROPT					
79	11	1655	RLADLEWT					
79	11	1655	RLADLEWTST					
79	11	2212	RLSAPSLKAT					
79	11	2212	RLSAPSLKATCT					
93	13	2207	RLSAPSLKAT					
100	14	175	RLSIFLALL					
86	12	175	RLSIFLALLSGL					
100	14	1470	RLSDPTFTI					
86	12	1470	RLSDPTFTIET					
79	11	1470	RLSDPTFTIETT					
79	11	2926	RLSHSYSPGEI	0.0008				
86	12	1051	RLSHSYSPGEI	0.0002				
86	12	1051	RLSHSYSPGEI	0.0053				
100	14	2178	RLSMLTDPDSHI					
100	14	2178	RLSMLTDPDSHIIT					
100	14	2178	RLSMLTDPDSHITA					
86	12	2163	RLSOLPCEPEOV					
93	13	2209	RLSOLPCEPEOV					
79	11	2209	RLSOLPCEPEOV					
78	11	2209	RLSOLPCEPEOV					

## UCV A02 Super Motif with Binding Information

Conservancy	Freq.	Position	Sequence	A*0201	A*0202	A*0203	A*0206	A*6802
93	13	56	SOPRGRROPI					
86	12	1242	STKVPAAYA					
79	11	1242	STKVPAAYAA					
100	14	1784	STLPGNPA					
79	11	1784	STLPGNPAI					
79	11	2	STNPKPORIKT	0.0007				
86	12	1663	STWLVGGV					
86	12	1663	STWLVGGVL					
86	12	1663	STWLVGGVLA					
88	12	1299	STYCKFLA					
100	14	1262	SVAATLGFCA					
86	12	1455	SVIDCNTCV					
86	12	1455	SVIDNTCVT	0.0088				
88	12	995	TAACGDII					
86	12	1343	TAGARLVV					
86	12	1343	TAGARLVWL					
88	12	1343	TAGARLVVLA					
79	11	1343	TAGARLVVLAT					
79	11	2852	TARHPVNSWL					
79	11	2590	TIMAKNEV					
93	13	1266	TLGFGAYM					
88	12	1266	TLGFGAYMSKA					
79	11	1822	TLHGPTPL					
79	11	1822	TLHGPTPLC	0.0070				
88	12	1811	TLHPNLGGW					
79	11	686	TLPALSTGL					
79	11	688	TLPALSTGLI					
79	11	1785	TLPGNPAI	0.0003				
86	12	125	TLTGGFADL	0.0004				
86	12	125	TLTGGFADLM					
79	11	2871	TLWARMIL	0.0003				
79	11	2871	TLWARMILM					
79	11	2871	TLWARMILMT					
88	12	1209	TMRSVPFT					
86	12	1484	TQTYDLSL					
86	12	1484	TQTYDLSLPT					
79	11	2589	TTIMAKNEV					
79	11	685	TTLPALST					
79	11	685	TTLPALSTGL					
79	11	685	TTLPALSTGLI					
86	12	1208	TTMRSVPFT					
79	11	2739	TTSCGNIL					



## HCY A02 Super Motif with Binding Information

Conservancy	Freq.	Position	Sequence	A*0201	A*0202	A*0203	A*0208	A*6802
79	11	2739	TTSCGNLT					
79	11	1597	TVCARAOA					
86	12	1466	TVDFSLDPT					
86	12	1466	TVDFSLDPTFT					
100	14	1336	TVLDOAET					
100	14	1336	TVLDOAETA					
86	12	1336	TVLDOAETAGA					
100	14	1263	VAATLGFGA					
93	13	1263	VAATLGFAYM					
88	12	1230	VAHLHAPT					
86	12	1440	VATDALMT					
86	12	1592	VAYQATVCA					
79	11	1592	VAYQATVCARA	0.0005				
100	14	1420	VAYYRGLDV	0.0001				
100	14	1420	VAYYRGLDVSV					
86	12	1456	VIDNTCV					
86	12	1456	VIDNTCVT					
88	12	1456	VIDNTCVTOT					
86	12	122	VIDTLTGFA					
86	12	1671	VLAALAYCL					0.0550
93	13	1521	VLCECYDA					
79	11	1521	VLCECYDAGCA					
100	14	1337	VLDIAETA					
86	12	1337	VLDIAETAGA					
86	12	157	VLEDGWNYA					
86	12	157	VLEDGWNYAT					
100	14	1258	VLNPSVAA					
100	14	1258	VLNPSVAAT					
100	14	1258	VLNPSVAATL	0.0015				
79	11	2737	VLTTSCGNT	0.0002				
79	11	2737	VLTTSCGNLT					
79	11	2737	VLTTSCGNLT					
79	11	1852	VLVDILAGYA					
86	12	1666	VLVGGVLA					
86	12	1666	VLVGGVLAAL					
86	12	1666	VLVGGVLAAL					
86	12	1666	VLVGGVLAAL					
100	14	1256	VLVLPVSV					
100	14	1256	VLVLPVSA					
100	14	1256	VLVLPVSA					
100	14	1256	VLVLPVSAAT	0.0270	0.0130	0.3100	0.0120	0.0130
79	11	2600	VOPEKGGKPA	0.0084				
				0.0009				

## HCY A02 Super Motif with Binding Information

Conservancy	Freq.	Position	Sequence	A*0201	A*0202	A*0203	A*0208	A*6802
100	14	1918	VQWNNRIU					
100	14	1918	VQWNNRIUA					
100	14	1918	VQWNNRIUFA					
86	12	1463	VTOIVDFSL					
79	11	1138	VTRHAQVI					
79	11	1138	VTRHADVPV					
85	12	1661	VTSTWVLV					
86	12	1661	VTSTWLVGGV					
79	11	1439	VVATDALM					
79	11	1438	VVATDALMT					
79	11	1901	VVCAAILRHV					
79	11	1898	VVGWCAAI					
79	11	1898	VVGWCAAIL					
86	12	1660	VVTSTWVL					
86	12	1660	VVTSTWVLV	0.0003				
86	12	1766	WAKHWNFR	0.0001				
86	12	76	WQPGYRPL					
86	12	2573	WARMILMT					
79	11	2287	WARPYNPPL					
100	14	1920	WMNRUFA	0.0410	0.0330	3.0000	0.0023	0.1000
79	11	557	WMNSTGFT					
86	12	1665	WVLGGVL					
86	12	1665	WVLGGVLA	0.0005				
86	12	1665	WVLGGVLA	0.0015				
86	12	1665	WVLGGVLAAL					
79	11	1249	YAAQGYKV					
79	11	1249	YAAQGYKVL					
79	11	1249	YAAQGYKVLV					
79	11	1249	YAAQGYKVLV	0.0050				
79	11	136	YIPLVGAPL					
100	14	1779	YLAGLSTL					
86	12	1165	YKGGSGGPL	0.0802				
86	12	1165	YKGGSGGPL					
93	13	35	YLFTHCTIL	0.0400	0.0007	0.0220	0.0089	0.0039
79	11	2836	YLRDPTT					
86	12	1590	YLVAQAT					
86	12	1590	YLVAQATV	0.2500	0.1100	0.6300	0.0450	1.2000
86	12	1590	YLVAQATVCA					
86	12	1138	YLVTRHADV	0.0110	0.0021	2.8000	0.0520	0.0130
79	11	1136	YLVTRHADVI					
93	13	1594	YQATVCARA					

HCY A02 Super Motif with Binding Information

Conservancy	Freq.	Position	Sequence	A*0201	A*0202	A*0203	A*0205	A*6802
79	11	1594	YQATVCARACA					
79	11	1106	YTRWOODL					
79	11	1106	YTRVDDXLV					
86	12	276	YVGDLCGSV	0.0018				
86	12	278	YVGDLCGSVFL					
93	13	637	YVGGVHFL	0.0008				
86	12	1939	YVPESDAA					
86	12	1939	YVPESDAAA					
86	12	1939	YVPESDAARV					
			555					

Table IX

HCY A03 Super Motif (With Binding Information)

Conservancy	Freq.	Position	Sequence	A*0301	A*1101	A*3101	A*3301	A*8901
86	12	847	AACAWTRGER					
79	11	147	AARALAHGVR					
79	11	1187	AAVCTRGVAK					
79	11	2208	ASQASPSLK					
86	12	1265	ATLGFAYMSK					
79	11	49	ATRIKTSER					
79	11	1168	AVCTRGVAK	0.0260	0.0250	0.0011	0.0004	0.0001
86	12	2941	CLRLGVPRUR					
79	11	565	CTWIMSTGFTK					
79	11	2598	CVPEKGGK	0.7600	0.7500			
79	11	2599	CVPEKGGK	0.0008	0.0005			
79	11	2599	CVPEKGGK	0.0011	0.0008			
100	14	1574	DAHFLSQTK	0.0003	0.0005			
93	13	2617	DGVRVCEK	0.0003	0.0002	0.0006	0.0440	0.0002
79	11	1143	DVIVPRR					
86	12	2245	EMGNITN					
86	12	2598	EYFCVCEK					
100	14	728	FLLDATN					
79	11	146	GAARALAHGVR	0.0008	0.0270	0.0003	0.0005	0.4500
100	14	1918	GAACWNR					
79	11	3037	GILLPNR					
79	11	1004	GLPSARR					
86	12	1131	GSSDLYVTR					
86	12	1883	GVAGALVAFK					
79	11	3035	GVGYLLPNR	0.3900	1.4000	0.0055	0.0011	0.0880
79	11	45	GVRATKTSER	0.0014	0.0140	0.1500	0.0130	0.0007
79	11	1900	GVCAAILR					
79	11	1900	GVCAAILR					
93	13	33	GVYLLPRR					
93	13	33	GVYLLPRGR					
79	11	1141	HADVIVR					
79	11	1141	HADVIVR					
79	11	1141	HADVIVR					
79	11	1141	HADVIVR					
100	14	1234	HAPTGSCK					
93	13	1234	HAPTGSCK					
100	14	1572	HIDAHFLSQTK					
86	12	1232	HUWPTGSCK	0.5900	0.0024	0.0005	0.0006	0.0028
100	14	1395	HLFCHSK					
100	14	1395	HLFCHSK	0.0250	0.0006	0.0003	0.0004	0.0010
100	14	1395	HLFCHSK	0.0260	0.0002	0.0009	0.0006	0.0001
79	11	2928	HSYSGEINR					
79	11	222	HTPGCVPR	0.0004	0.0012	0.0007	0.0006	0.0092
86	12	2250	ITRVESENK	0.0150	0.0079			
86	12	1298	ITYSTYCK					
79	11	2813	MFPLGVR					
93	13	30	NGGVYLLPR	0.0036	0.0044			
93	13	30	NGGVYLLPR	0.0008	0.0058			
86	12	2944	KLGVPRR					
86	12	10	KTKRNTNR					
86	12	10	KTKRNTNR					
93	13	51	KTSEROPR	0.0110	0.0100	0.2700	0.0160	0.0550
86	12	51	KTSEROPR	0.1600	0.0640			
86	12	1729	LAECFCK					

## ICV ADJ Sauer Motif (With Binding Information)

Conservancy	Freq.	Position	Sequence	A*0301	A*1101	A*3101	A*3301	A*8801
86	12	2235	UEANLLWR					
100	14	1388	LIFCHSKK					
100	14	1388	LIFCHSKK					
79	11	2612	LVFPDLGVR	0.5400	0.1800	0.0071	0.0089	0.0008
100	14	726	LLRLADAR	0.0003	0.0001		0.0012	0.0240
93	13	36	LLPRGPR					
86	12	87	LLSPGSR					
79	11	1591	LVAYQATVCAR					
79	11	1	MSINPKPOR					
79	11	1	MSINPKPOR					
86	12	2249	NITRVESENK	0.0010	0.0062			
79	11	14	NINRPOOVK	0.0010	0.0007			
78	11	1295	PITYSTYCK					
79	11	2667	PMGFSYOTR					
93	13	514	PSPVWGTIDR					
79	11	1607	PSYDQMKK					
86	12	109	PTDPRFTR					
93	13	1238	PTSGSKTK					
86	13	516	PVWGTIDR					
88	12	1340	QAEIAGAR					
93	13	28	QNGGYLLPR					
86	12	289	QUTFSR					
79	11	289	QUTFSR					
79	11	2210	QLSAPSLK	0.0008	0.0005	0.0008	0.0008	0.0002
79	11	1168	RAAVCTRGVAK	0.0008	0.0005			
100	14	149	RAIAGVR	0.7500	0.0330	0.0290	0.0077	3.1000
79	11	47	RATKTSER					
79	11	43	RLGVRATR					
78	11	43	RLGVRATR					
100	14	1923	RLNFASR	0.9400	0.0280	0.0420	0.0004	0.0001
79	11	2611	RLVFPDLGVR					
100	14	635	RMVGGVDR	0.7200	0.0200	0.1900	0.0030	0.0045
93	13	55	RSQRTGR					
79	11	2207	SASQLSAPSLK					
86	12	1132	SSDLVLTR	0.0003	0.0044			
79	11	2	STNPKPOR					
79	11	2	STNPKPOR					
79	11	2	STNPKPORKTK					
86	12	1266	TLGFGAYMSK	0.0810	0.0610	0.0005	0.0013	0.0009
79	11	1622	TLHGFTLLYR					
93	13	52	TSESRPH					
86	12	52	TSESRPH					
86	12	52	TSESRPH					
86	12	1050	TSITGRDK					
86	12	1864	VAGALVFK					
79	11	1592	VAYQATVCAR	0.2400	0.8900	0.0048	0.0025	0.0310
86	12	1337	VLDQAEIAGAR	0.0005	0.0038	0.0580	0.0720	0.0280
79	11	1138	VITHADVPVR					
79	11	1901	VYCAAILR					
79	11	1901	VYCAAILR					
79	11	1898	VGVVCAAILR					
93	13	517	VWGTIDR					

## IICV A01 Super Motif (With Blinding Information)

Conservancy	Freq.	Position	Sequence	A*0301	A*1101	A*3101	A*3301	A*8801
86	12	93	WAGWLLSPR					
86	12	86	WLLSPRGR	0.0008	0.0005			
100	14	1920	WMNRLAFASR					
79	11	557	WMNSTGFTK	0.0530	0.0810	0.0014	0.0420	0.0058
93	13	35	YLLPRGRPR	0.0054	0.0005			
79	11	2930	YSPGENR					
100	14	637	YGGVGR					
86	12	1939	YVPESDAAF	0.0003	0.0001			
			112					

Table X  
HCV A24 Super Motif With Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*2401
AILSPGAL	1890	8	13	93	
ALAHGVRVL	150	9	14	100	
ALSTGLHL	689	9	12	86	
ALVVGVCAAI	1896	11	79	79	
ATGNLPGCSF	165	10	13	93	
ATLFGAY	1265	6	12	100	
ATLFGAYM	1265	9	13	93	
AVAYRGL	1419	8	14	100	
AVQWNNRL	1917	8	14	100	
AVQWNNRLI	1917	9	14	100	
AVQWNNRLIAF	1917	11	14	100	
AWQMMMNW	319	8	12	86	
AYAAQGYKVL	1248	10	11	79	0.0009
AYYRGLDVSII	1421	11	14	100	
CLRKLGVPPL	2941	10	12	86	
CLWIMLLI	739	8	12	86	
CTGSSDL	1128	8	11	79	
CTGSSDLY	1128	9	11	79	
CTGSSDLYL	1128	10	11	79	
CTRGVAKAVDF	1190	11	11	79	
CTWNNSTGF	555	9	11	79	
CYTQTVDF	1462	8	12	86	
CYTQTVDFSL	1462	10	12	86	
CYDAGCANY	1525	8	11	79	
CYDAGCAWY	1525	9	11	79	
CYDAGCAWYEL	1525	11	11	79	
DFSLDPTF	1468	8	14	100	
DFSLDPTFTI	1468	10	14	100	
DLCGSVRL	279	8	12	86	
DLEWYTSW	1657	9	12	86	
DLEWYTSWVL	1657	11	12	86	
DLGVAVCEQM	2617	10	13	93	
DLMGYIPL	132	8	11	79	
DLVNLPAI	1883	9	11	79	
DLVNLPAI	1883	10	11	79	
DTAACGDI	994	8	12	86	
DTAACGDI	994	9	12	86	
DTLTCGFADL	124	10	12	86	
DTLTCGFADLM	124	11	12	86	
DWRFPGGGI	21	10	12	86	
DYPYRLWHY	615	9	14	100	
EIPFYGKAI	1377	9	13	93	
ETAGARLVVL	1342	10	12	86	
ETMRSPIF	1207	9	12	86	
EVTSTWVL	1659	9	12	86	

## HCY A24 Super Motif With Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*2401
FSGQYL	1773	8	14	100	
FSGIYLAGL	1773	11	14	100	
FLALLSL	177	9	12	86	
FTEAMTRY	2792	8	14	100	
FTGLTHDAHF	1567	11	13	93	
FTLPALSTGL	884	11	11	79	
FWAGHMMNF	1765	9	12	86	6.9000
FWAGHMMNF	1765	10	12	86	
GFAULMGY	129	8	13	93	
GFAULMGYI	129	9	13	93	
GFAULMGYIPL	129	11	11	79	
GFSYDITCF	2669	8	11	79	
GIQYLAGL	1776	8	14	100	
GIQYLAGLSTL	1776	11	14	100	
GLPVQDHL	1552	0	13	93	
GLPVQDHLF	1552	11	12	86	
GLSAFSLHSY	2921	10	11	79	0.0001
GLSTLPGNPI	1782	11	11	79	
GLTHDAHF	1568	9	13	93	
GLTHDAHFL	1568	10	13	93	
GTFPINAY	2063	8	11	79	
GVAGALVAF	1863	8	12	86	
GVAKAYDF	1193	8	11	79	
GVLAALAAY	1670	9	12	86	
GVLAALAAYCL	1670	11	12	86	
GVNYATGNL	161	9	11	79	
GVRVCEGM	2619	8	14	100	
GVRVCEKWL	2619	10	14	100	
GVRVCEKMLY	2619	11	14	100	
GVRLEDQNY	154	11	12	86	
GWCAAL	1800	8	11	79	
GWILLAPI	1027	8	11	79	
GWILLAPITAY	1027	11	11	79	
GYGAGVAGAL	1859	10	12	86	0.0003
GYPLVCAPL	135	10	10	79	0.0057
GYRFRASGVL	2728	11	12	86	
HLEHNMVQY	696	11	11	79	
HLPIECGM	1719	9	11	79	
HMMNFSGI	1769	9	13	93	
HMMNFSGIY	1788	11	13	93	
HMPNLSWL	2855	8	12	86	
HMPNLSWLN	2855	11	12	86	
HMPNLSWLN	1910	11	11	79	
HMPNLSWLN	176	10	12	86	
ILGGWAAQL	1816	10	12	86	0.0026



124

HCY A24 Super Motif With Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*2401
ILGIGTIL	1331	8	12	86	
IMANNEVF	2591	8	12	86	
ITYSTYKGF	1296	9	12	86	
ITYSTYKGL	1296	10	11	79	
NDVQVLY	701	8	12	86	
NGGVYLL	30	8	13	93	
KFFGGGCI	23	8	13	93	
KVIDTLTCGF	121	10	12	86	
LFNLGGW	1813	8	12	86	
LEANLLW	2235	8	12	86	
LININGSW	414	8	11	79	
LLALLSCL	170	8	12	86	
LLAPITAY	1030	8	14	00	
LLNLGGW	1812	9	12	86	
LLPALSPGAL	1897	11	13	93	
LLPRGGPRL	36	9	13	93	
LLSPRGSRPSW	97	11	11	79	
LLWQDEMGNI	2240	11	12	06	
LTCGFADL	126	8	12	86	
LTCGFADLM	126	9	12	86	
LTCGFADLMGY	126	11	12	86	
LTHIDAHF	1670	8	13	93	
LTHIDANFL	1570	8	13	93	
LTSMLTDPSSH	2176	11	13	83	
LTISCGNTL	2738	9	11	79	
LVDLAGY	1853	8	11	79	
LVGGVLAAL	1887	9	12	86	
LVNPSVAATL	1257	11	14	100	
LVNLPAL	1884	8	11	79	
LVNLLPAIL	1884	9	11	79	
LVTRHADVI	1137	9	11	79	
LVGVVCAAI	1897	10	11	79	
LVGVVCAAIL	1897	11	11	79	
LWARMILM	2672	8	12	86	
LWARMILMTHF	2672	11	12	86	
LWROEMGGN	2241	10	12	86	
LYLYTRHADVI	1135	11	11	79	
MILMTHFF	2876	8	12	86	
MLTOPSHI	2179	8	14	100	
MWNFSGI	1770	8	14	100	
MWNFSGIQY	1770	10	14	100	
MWNFSGIQYL	1770	11	14	100	
MYGVGVHFL	636	10	13	93	0.0270
NFSGIQY	1772	8	14	100	
NFSGIQYL	1772	9	14	100	0.0170

HCV A24 Super Motif With Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*2401
NILGHWAAQL	1815	11	12	86	
NRTGVRTI	1282	9	11	79	
NMDVQYL	700	8	12	86	
NNDVQYLY	700	9	12	86	0.0001
NLGRYDITL	118	9	12	86	
NLWROEM	2239	8	12	86	
NLPGCSFSI	168	9	13	86	
NLPGCSFSF	188	10	13	93	
NLPGCSFSL	188	11	13	93	
NITVITQVDF	1460	10	12	86	
NINGSWHI	416	8	13	93	
NTNRFPDVKF	14	11	11	79	
NMDLVGW	1108	9	11	79	
NWFGCTMM	551	8	12	86	
PITYSTYKGF	1295	10	11	79	
PITYSTYKGL	1295	11	11	79	
PLEGGQXDL	2403	11	13	93	
PLGGAARAL	143	9	11	79	
PKGFSYDTRCF	2667	11	11	79	
PIDRRRSRL	109	11	12	86	
PTLHGPTPL	1621	9	11	79	
PTLHGPTPL	1821	10	11	79	
PTLHGPTPL	1621	11	11	79	
PTLWARM	2870	8	11	79	
PTLWARMIL	2870	9	11	79	
PTLWARMILM	2870	10	11	79	
PTLLYRL	1626	8	14	100	
PVODDIEF	1554	9	12	86	
PVODDLEPW	1554	10	12	86	
PVNSWLGNI	2857	9	14	100	
PVNSWLGNI	2857	10	14	100	
PVNSWLGNIIM	2857	11	12	86	
PVHGGPL	2318	8	11	79	
OFKXALGL	1732	9	12	86	
OFKXALGLL	1732	10	12	86	
QVGGVYL	29	8	13	93	
QVGGVYL	29	9	13	93	
QVDFSLDPTF	1465	11	12	86	
QWNRUAF	1919	9	14	100	
QYLAGLSTL	1778	9	14	100	0.0480
QYSPQRVEF	2847	10	11	79	0.0180
QYSPQRVEFL	2647	11	11	79	
RLHGLSAF	2918	8	12	86	
RLHGLSAFSL	2918	10	11	79	0.0001
RLVFPDL	2611	8	11	79	

## HCV A24 Super Motif With Binding Information

Sequence	Position	Pepide No.	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*2401
RLAPITAY	1029		9	12		
RMAWDMM	317		8	12	86	
RMAWDMMNW	317		10	12	86	
RMILMTIF	2875		8	12	86	
RMILMTIFF	2875		9	12	86	
RMVGGVEHL	635		11	13	86	
RYCEKML	2621		8	14	93	
RYCEKMLY	2621		9	14	100	
RYLEDGVNY	156		9	12	100	
SFSIFLLAL	173		9	12	86	
SFSIFLLALL	173		10	14	100	
SIFLLALL	175		8	14	100	0.0041
SIFLLALLSCL	175		11	14	100	
SLDPTFTI	1470		8	12	86	
SLHSYSPGEI	2928		10	11	100	
SMLTDPISH	2178		9	14	79	
STKVPAAY	1242		8	12	100	
STLPGNPAI	1784		9	11	79	
STWLVGGVL	1683		10	12	86	
SVAATLGF	1262		8	14	100	
SVAATLGFAY	1262		11	14	100	
SWDDMMKQL	1608		9	11	79	
SWLGNIM	2860		8	12	86	
SYLKGSSGGPL	1164		11	12	86	
TIMAKNEVF	2590		9	11	79	
TLGFGAYM	1266		8	13	93	
TLHGPTPL	1622		8	11	79	
TLHGPTILL	1622		9	11	79	
TUIGPTILLY	1822		10	11	79	
TLFNILGGW	1811		10	12	60	0.0001
TLPALSTGL	586		9	11	79	
TLPALSTGLI	586		10	11	79	
TLPGNPAI	1786		8	11	79	
TLTCSFADL	125		9	12	86	
TLTCSFADUM	125		10	12	86	
TLWARMIL	2871		8	11	86	
TLWARMILM	2871		9	11	79	
TLIMAKNEVF	2589		10	11	79	
TLIPALSTGL	685		10	11	79	
TLIPALSTGLI	685		11	11	79	
TTMPSPVF	1208		8	12	86	
TTSCGNIL	2739		8	11	79	
TVDFSLDPTF	1466		10	12	86	
TWAKNSTGF	556		8	11	79	
TWLVGGVL	1664		9	12	86	

## HCY A24 Super Motif With Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*2401
TYSTYKIF	1297	8	13	93	
TYSTYGRRL	1297	9	12	86	0.0230
VFTGLTHI	1566	8	13	93	
VIDILTCGF	122	9	12	86	
VLAALAAY	1671	8	12	86	
VLAALAAYCL	1671	10	12	86	0.0070
VLEDGVNY	167	8	12	86	
VLNPSVAATL	1258	10	14	100	
VLITSCGNIL	2737	10	11	79	
VLVDILAGY	1852	9	11	79	
VLVGGVLAAL	1868	10	12	86	
VMGSSYGF	2639	8	11	79	
VMGSSYGFQY	2639	10	11	79	
VTQVDFSL	1463	9	12	86	
VTRHADVI	1138	8	11	79	
VWATDALM	1439	8	11	79	
VWGWCAAI	1898	9	11	79	
VWGWCAAIL	1898	10	11	79	
WTSTWVL	1660	8	12	86	
WLLPRGPRLL	34	11	13	93	0.0016
WNNRLAF	1920	8	14	100	
WLVGGVL	1665	8	12	86	
WLVGGVLAAL	1665	11	12	86	
YIPLVGAPL	136	9	11	79	
YLAGLSTL	1779	8	14	100	
YKSSGGPL	1165	10	12	86	
YKSSGGPRL	1165	11	12	86	
YLPTIRGPTL	35	10	13	93	0.0001
YLVTRHADVI	1136	10	11	79	
YTNDDCL	1106	8	11	79	
YTNDDCLVGW	1106	11	11	79	
YWGDCSSVF	276	10	12	86	
YWGDCGSVFL	276	11	12	86	
YWGVEHFL	637	9	13	93	
YVRGLDVSI	1422	10	14	100	
260		3			

Table XI

HCV B07 Super Motif (with Binding Information)

Conservancy	Freq.	Position	Sequence	B*0702	B*3501	B*5101	B*5301	B*5401
86	12	1804	APPSWDCM	0.0028	0.0002	0.0002	0.0001	0.0002
79	11	1804	APPSWDCM	0.0001	0.0001	0.0002	0.0006	0.0003
93	13	1235	APTSCKSTKV	0.0001	0.0001	0.0012	-0.0002	0.0023
79	11	2869	APTLWARM	0.4300	0.0002	0.0012	0.0001	0.0002
79	11	2869	APTLWARM	0.0160	0.0001	0.0010	0.0001	0.0003
79	11	2869	APTLWARMIL	0.8000	0.0001	-0.0003	-0.0002	0.0033
79	11	2869	APTLWARMILM	0.0130	0.0001	0.0002	0.0005	0.0002
79	11	2410	DPQLSDGSW	0.0001	0.0002	0.0002	0.0001	0.0002
86	12	111	DPFRSRML	0.0170	0.0002	0.0001	0.0001	0.0002
79	11	2815	FPOLGVV	0.0001				
100	14	24	FRGGGV	0.0001				
100	14	24	FRGGGVGV	0.0001				
86	12	1912	FRGGGVGV	0.0001	0.0002	0.0002	0.0001	0.0002
86	12	1912	FRGGGVGV	0.0001	0.0001	0.0002	0.0001	0.0003
93	13	41	GPGLGVRA	0.0001	0.0002	0.0002	0.0001	0.0002
100	14	1825	GPPLLYRL	0.0024	0.0002	0.0002	0.0001	0.0002
93	13	1625	GPPLLYRLGA	0.0005				
93	13	507	GPVYCFPSFV	0.0001	0.0001	0.1200	-0.0002	0.2000
82	13	1378	IFPYGKAI	0.0120	0.0032	0.0700	0.0003	0.0035
79	11	137	IPLVGAPL	0.4400	0.0002	0.0017	-0.0002	0.0009
86	12	2608	KPARLVF	0.0001	0.0001	0.0002	0.0001	0.0003
79	11	2608	KPARLVF	0.0003	0.0001	0.0002	0.0002	0.0003
79	11	1820	KPARLVF	0.0003	0.0001	0.0002	0.0001	0.0003
79	11	1820	KPARLVF	1.4150	0.0001	0.0002	0.0001	0.0003
93	13	1088	KPARLVF	0.0021	0.0001	0.0001	0.0002	0.0002
93	13	1088	KPARLVF	0.0001	0.0001	0.0001	0.0002	0.0002
93	13	1888	KPARLVF	0.0053	0.0001	0.0036	0.0001	0.0001
86	12	1888	KPARLVF	0.0003	0.0001	0.0001	0.0001	0.0001
100	14	807	KPARLVF	0.0020	0.0002	0.0002	0.0002	0.0005
86	12	887	KPARLVF	0.0350	0.0002	0.0002	0.0002	0.0005
86	12	887	KPARLVF	0.0011	0.0002	0.0002	0.0002	0.0005
86	12	2165	KPARLVF	0.0001	0.0002	0.0001	0.0001	0.0002
93	13	169	KPARLVF	0.0110	0.0360	0.0059	0.0150	0.0016
93	13	169	KPARLVF	0.1950	0.0796	0.0550	0.0013	0.0015
93	13	169	KPARLVF	0.0022	0.0009	0.0100	0.0140	0.0012
93	13	37	KPARLVF	0.0007	0.0001	0.0180	-0.0002	0.0020
93	13	37	KPARLVF	0.5000	0.0001	0.0009	0.0001	0.0025
93	13	37	KPARLVF	0.0005	0.0001	0.0009	0.0001	0.0025
93	13	1553	KPARLVF	0.0001	0.0046	0.0002	0.0110	0.0003
86	12	1553	KPARLVF	0.0001	0.0001	0.0002	0.0110	0.0003
86	12	1553	KPARLVF	0.0001	0.0001	0.0002	0.0110	0.0003
86	12	1720	KPARLVF	0.0130	0.0001	0.0040	-0.0002	0.0013
100	14	1260	KPARLVF	0.0011	0.0001	0.0002	0.0001	0.0003
100	14	1260	KPARLVF	0.0001	0.0001	0.0002	0.0001	0.0003
86	12	1605	KPARLVF	0.0001	0.0002	0.0001	0.0001	0.0002
79	11	1605	KPARLVF	0.0001	0.0001	0.0002	0.0001	0.0002
79	11	1605	KPARLVF	0.0001	0.0001	0.0002	0.0001	0.0002
79	11	1605	KPARLVF	0.0001	0.0001	0.0002	0.0001	0.0002
79	11	2317	KPARLVF	0.0140	0.0001	0.0001	0.0001	0.0002
79	11	2601	KPARLVF	0.0011	0.0001	0.0001	0.0001	0.0002
79	11	2809	KPARLVF	0.0002	0.0001	0.0001	0.0001	0.0002
79	11	2808	KPARLVF	0.0001	0.0002	0.0002	0.0001	0.0002
86	12	78	KPARLVF	0.0006	0.0002	0.0002	0.0001	0.0002

## UCV B07 Super Motif (with Blinding Information)

Conservancy	Freq.	Position	Sequence	B*0702	B*3501	B*5101	B*5301	B*5401
86	12	78	QPGVPMPLY	0.0001	0.0011	0.0002	0.0001	0.0002
93	13	57	QPGRRQR	0.2300	0.0002	0.0001	0.0001	0.0002
79	11	2298	HPQYNPPL	0.0050				
93	13	1893	SPGALVGV	0.0001	0.0002	0.0002	0.1200	0.0002
79	11	1893	SPGALVGVW	0.0130	0.0001	0.0016	0.0001	0.0003
79	11	2931	SPGEINRV	0.0007				
79	11	2931	SPGEINRVA	0.0003	0.0001	0.0001	0.0002	0.0037
79	11	2649	SPQORVEF	0.0027				
79	11	2649	SPQORVERL	0.1200	0.0002	0.0002	0.0001	0.0002
79	11	99	SPRGSRPSW	0.3600	0.0002	0.0005	0.0001	0.0002
86	12	1935	SPTHVVPESDA	0.0001				
86	12	1975	TPCSGSWL	0.0028				
79	11	1126	TPCTGSSDL	0.0005	0.0001	0.0002	0.0001	0.0003
79	11	1126	TPCTGSSDLY	0.0001				
86	12	223	TPGVPCV	0.0001				
93	13	1550	TPGLVQDHL	0.0001				
93	13	1627	TPLLYRLGA	0.0063	0.0001	0.0001	0.0002	0.2300
93	13	1627	TPLLYRLGAV	0.0120	0.0001	0.0008	0.0001	0.0110
86	12	2056	TPVNSWLGNH	0.0001	0.0001	0.0053	0.0008	0.0003
86	12	2856	TPVNSWLGNH	0.0001				
86	12	1940	VPESDAAA	0.0022				
86	12	1940	VPESDAARV	0.0001	0.0001	0.0010	0.0001	0.0003
86	12	789	WPLLLLL	0.0021				
100	14	616	YPVRLWHY	0.0001				

76

HCV B27 Super Motif

Table XII

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
AKHWINFI	1767	8	12	86
AKNEVFCV	2593	8	12	86
ARALAHGV	148	8	14	100
DRSELSPL	663	8	11	79
EKGGKQPA	2603	8	11	79
EKMALYDV	2624	8	12	86
FKCKALGL	1733	8	12	86
GHRMAWDM	315	8	13	93
GKSTKVPA	1240	8	12	86
GRKPARLU	2806	8	11	79
HRMAWDMAM	316	8	13	93
KGGKHLI	1390	8	11	79
ITGVRTI	1283	8	11	79
KKCDELA	1403	8	14	100
KKCDELA	1402	8	14	100
LHGFTPLL	1623	8	11	79
LHONVDV	697	8	12	86
LRLAVAV	989	8	11	79
NHVSPTHY	1832	8	12	86
PRGRPRPI	58	8	13	93
PRGSRPSW	100	8	11	79
PRRSRL	112	8	12	86
RHADVPV	1140	8	11	79
RHTPVNSW	2854	8	12	86
RKLGVPPL	2943	8	12	86
RKPARLIV	2607	8	11	79
RICRASGV	2730	8	13	93
RKGFTLGV	38	8	13	93
RRFQDKF	17	8	12	86
SKKCDL	1401	8	14	100
SRNLGKVI	118	8	12	86
THDAHL	1571	8	13	93
TKLKLPI	2985	8	12	86
TKVPAAYA	1243	8	12	86
TRCFDSTV	2674	8	14	100
TRGVAKAV	1191	8	11	79
VFVCEKMA	2620	8	14	100
VWLEDGV	155	8	13	93
YRGDVS	1423	8	14	100
ARHTPVNSW	2853	9	11	79
ARLVFPDL	2610	9	11	79
ARLVULATA	1348	9	11	79
ARMLMTHF	2874	9	12	86
APPDYNPPL	2298	9	11	79
DRSELSPL	663	9	11	79

HCV B27 Super Motif

Sequence	Position	Peptide No.	No. of Amino Acids	Sequence Frequency	Conservancy (%)
EKMALYDW	2624		9	12	86
FKKALGLL	1733		9	12	86
GHMAWDMM	315		9	13	93
GKSTKYPAA	1240		9	12	86
GRKPARLV	2608		9	11	79
HRMAWDMM	316		9	12	86
IKGGRHLF	1390		9	11	79
KKKDELA	1402		9	14	100
LHGLSAFL	2919		9	11	79
LHGPTLLY	1623		9	11	79
LHSYSPGEI	2927		9	11	79
LKSSGGFL	1166		9	12	86
LRLGVPL	2942		9	12	86
NHVSPTHY	1932		9	12	86
NRRQDWKF	16		9	11	79
PRGRPLGV	38		9	13	93
RHTPVNSWL	2854		9	12	86
RHNGFGQA	1909		9	11	79
RKPARLVF	2607		9	11	79
RRCRASGL	2730		9	12	86
RPSRLGV	114		9	12	86
SKKKDELA	1401		9	14	100
THYVPESDA	1937		9	12	86
TKYPAAYAA	1243		9	11	79
TRHADVPV	1139		9	11	79
TRVESENK	2251		9	12	86
WTFPGGCI	22		9	13	93
VRVCEKML	2620		9	14	100
WRLAPITA	1028		9	11	79
WRQVGGN	2242		9	11	79
YRGLDVM	1423		9	12	86
YRCRASGV	2729		9	14	100
YRALAHGVRV	148		9	13	93
ARAQAPPSW	1600		10	14	100
ARHTPVNSWL	2853		10	11	79
ARMILMTHFF	2874		10	11	79
CHSKKKDEL	1399		10	12	86
DHDSLSPL	661		10	14	100
DRSELSPL	663		10	11	79
EXGGRKPARL	2603		10	11	79
FRAAVCTRGV	1185		10	12	86
GHMAWDMM	315		10	12	86
GKSTKYPAA	1240		10	12	86
GRKPARLV	2606		10	11	79
KHWNFRSGI	1768		10	13	93



HCV B27 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy [%]
KKCDELAALK	1403	10	12	86
LHONVMOY	697	10	11	79
LKSSGGRL	1168	10	12	86
OKALGILQTA	1735	10	12	86
RRVGPGEAV	1909	10	11	79
RRGPRLGVRA	39	10	13	93
RRHNGPGEA	1908	10	11	79
RRHSRLGKV	113	10	12	86
RRSRNLGKI	114	10	12	86
SKFGYGAQV	2552	10	12	86
SKKKCDELA	1401	10	14	100
THYVPESDAA	1937	10	12	86
TRGVAKAVDF	1191	10	11	79
TRVESENKV	2251	10	12	86
VKFGGGGV	22	10	13	93
VRYCEKNALY	2620	10	14	100
VRLVDGVY	155	10	12	86
WRLAPITAY	1028	10	11	79
YKVLVNPVS	1254	10	14	100
YRRCRASGV	2729	10	13	86
YGVVRLDGV	152	11	12	86
YKHMVNFSGI	1767	11	14	100
ARALAHGVRL	148	11	11	79
ARLVFPDLGV	2610	11	14	100
CHSKKKCDELA	1399	11	11	79
DHRSLSRL	651	11	11	79
EKGGRKPARJ	2603	11	11	79
FRAAVCTRGVA	1185	11	12	86
GKSTKVPAAVA	1240	11	12	86
GKVIDLTGCF	120	11	12	86
HRMANVDMANNW	316	11	12	86
KKCDELAALK	1402	11	12	86
KRNTNRPPQDV	12	11	12	86
LHGPTPLLYRL	1623	11	11	79
LHONVMOYL	697	11	11	79
LKPTLHGPTPL	1619	11	11	79
LRFHNGPGEA	1907	11	13	93
PRGPRLGVRA	38	11	12	86
PRRSRLGKV	112	11	12	86
RRHNGPGEAV	1908	11	11	79
RRSRNLGKI	113	11	12	86
SFGNHSPTHY	1829	11	12	86
SRNLGKVIDTL	116	11	12	86
THYVPESDAA	1937	11	12	86
VRLVDGVNYA	155	11	12	86

HCY B27 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
YKVLVLPNSVA 136	1254	11	14	100

HCV B58 Super Motif Table XIII

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
ANLRRHV	1804	8	13	93
AALAAVCL	1673	8	12	86
AAQGYKVL	1250	8	11	79
AATLGFGA	1264	8	14	100
AAVCTRGV	1187	8	12	88
ASLMAFTA	1793	9	11	79
ASSASQSL	2204	8	14	100
ATLGFAGY	1265	8	14	100
CSFSIRLL	172	8	14	100
CSGGAYDI	1310	8	12	86
CSSINWSVA	2819	8	14	100
CTCGSSDL	1128	8	11	79
CTRGVAKA	1190	8	11	79
DTAACGDI	994	8	12	86
DTLCGFA	124	8	12	86
EAALENLV	750	8	11	79
EAMTRYSA	2794	8	14	100
ESDAARV	1942	8	12	86
ETAGARLV	1342	8	12	88
ETMRSVP	1207	8	12	88
FADUMGYI	130	8	13	93
FASRGNIV	1927	8	14	100
FSIRLLAL	174	8	14	100
FSYDTRCF	2570	8	11	79
FTEAMTRY	2792	8	14	100
FTSPVW	512	8	13	93
GAGVAGAL	1861	8	12	86
GAHWGLA	350	8	12	86
GALWGW	1885	8	11	79
GARLVLA	1345	8	12	88
GSGKSTRV	1238	8	13	93
GSSDLVIV	1131	8	12	86
GSSGGFLL	1188	8	12	86
GSSYGFQY	2841	8	11	79
GTFPINAY	2083	8	11	79
HSYSPGEI	2928	8	11	79
HTPVNSWL	2855	8	12	86
ISGIQYLA	1774	8	14	100
ITSCSNW	2816	8	14	100
ITWGAQTA	989	8	12	86
KSTRKVPAA	1241	8	12	86
LAGYGAGV	1857	8	11	79
LAHGVRL	151	8	14	100
LAVAEPV	972	8	11	79
LSAPSLKA	2211	8	11	79

## HCY B58 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
LSPGALW	1892	8	13	93
LSTGLHL	690	8	12	86
LTCGFADL	126	8	12	86
LTHIDAHF	1570	8	13	93
MSADLEW	1654	8	11	79
NSWLGNI	2859	9	14	100
NTCVTQTV	1450	8	12	88
NTGSGWH	416	8	13	93
PAILSPGA	1889	8	13	93
PALSTGLI	668	8	12	86
PTLWARM	2870	8	11	79
PTLLYRL	1626	8	14	100
QATVCARA	1595	8	13	93
RAPPRWFM	3019	8	14	100
RSELPIL	664	8	11	79
RSRLQKV	115	9	12	86
SAFSLHSY	2923	8	11	70
SSASOLSA	2206	8	14	100
STKVPAAV	1242	8	12	86
STLPGNPA	1784	8	14	100
STLPOAVM	2633	8	12	86
STYGFCLA	1299	8	12	86
TACGSDII	895	8	12	86
TAGARLW	1343	8	12	86
TTMRSPVF	1208	8	12	86
TTSCGNTL	2739	8	11	79
VAGALVAF	1864	8	12	86
VTRHADVI	1138	8	11	79
VISTWYLV	1681	8	12	86
WAKHMMNF	1766	8	12	86
WAKVLVIM	368	8	14	100
WAGGYVW	78	8	12	86
YAAQGYKV	1249	8	11	79
YSIEFLD	2905	8	11	79
YSTYKRL	1298	8	12	88
YTWDOOL	1106	8	11	79
AAKLDDCTM	2758	9	16	114
AAQGYKLV	1250	9	11	79
AARALAHGV	147	9	11	79
AATLGFAGV	1264	9	14	100
AAVCTRGA	1187	9	11	79
ASQLSAPSL	2208	9	13	93
ATLGFAYM	1265	9	26	186
ATVCARAQA	1586	9	11	79
CAAILRRHV	1903	9	13	93

HCV B58 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
CAWYELTPA	1630	9	11	79
CSFSIELLA	172	9	14	100
CSGGAYDII	1310	9	12	86
CTCGSSDLY	1128	8	11	79
CTRGVAKAV	1190	8	11	79
CTWKNSTGF	555	2	11	79
DAGCAWYEL	1527	9	11	79
DTAACGDII	994	8	12	86
DTRCFDSIV	2673	8	13	93
ETAGARLW	1342	9	12	86
ETTMFSPVF	1207	9	12	86
FSRLALL	174	9	14	100
FSLDPTFTI	1469	9	14	100
FTGLTHIDA	1567	9	13	93
GAGVAGALV	1861	8	12	86
GALVAFKIM	1866	9	12	86
GALVAFKVM	1868	9	14	100
GAVOWWVRL	1916	9	14	100
HSKKKCOEL	1400	8	14	100
HIPGVPCV	222	9	11	79
ITWGAITAA	989	8	12	86
ITYSTYKGF	1296	9	12	86
KALGLLOTA	1736	9	12	86
KSTKVPAAV	1241	9	12	86
LAALAAAYCL	1672	9	12	86
LAQQRKOKA	1729	9	12	86
LAGLAYYSM	356	9	14	100
LAGYGAGVA	1857	9	11	79
LSAFSLHSY	2922	9	11	79
LSTLPGNPA	1783	9	14	100
LTCGFADLM	126	9	24	171
LTDPSHITA	2180	9	14	100
LTFGRKNQV	1052	9	12	86
LTHIDAHFL	1570	9	13	93
LITSCGNIL	2738	9	11	79
MAQNEVFCV	2592	9	12	86
MAVDMMMNW	318	8	12	86
NAVAYYRGL	1418	9	13	93
NSLRUFTNM	2481	9	14	100
NSWLGNIIM	2659	9	24	171
NINRPQDQV	14	9	12	86
PALSPGAL	1889	9	13	93
PSVAATLGF	1261	9	14	100
PTLHGPTPL	1621	9	11	79
PTLWARMIL	2870	9	11	79

HCV B58 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy [%]
QAETAGARL	1340	9	12	86
RAAVCTRGV	1186	9	12	86
RAAHGVGV	148	8	14	100
RAQAPPSW	1601	9	11	79
RAYAMDREH	811	9	16	114
RSELSPLL	664	2	11	79
RSNLTGKI	115	9	12	86
SSSASQLSA	2205	9	14	100
STKVPAAYA	1242	9	12	86
STLPGNPAI	1784	9	11	79
STWLVGGV	1663	9	12	88
TAGARLVVL	1343	9	12	86
TSCSSNVSV	2817	9	14	100
TTIMAKNEV	2589	9	11	79
VAATLGFGA	1263	9	14	100
VAGGHYVOM	933	9	14	100
VAYOATVCA	1592	9	14	86
VAYYRGLOV	1420	9	12	100
VSTLPOAVM	2832	9	12	86
VQTQVDFSL	1463	9	12	86
WAKHMMNFI	1768	9	12	86
YAAGYKVL	1249	8	11	79
YAPTLWARM	2868	9	14	100
YSPGEINRV	2930	9	11	79
YSPGRVEF	2848	9	11	78
YSTYKFLA	1298	9	12	86
YTNVDDOLV	1106	9	11	78
YAQGYKVL	1250	10	11	79
AATLGFGAYM	1264	10	28	186
ASLRVTEAM	2787	10	12	86
ASSSASQLSA	2204	10	14	100
ATGNLPCCSF	165	10	13	93
CSFSIRLAL	172	10	14	100
CTCGSSQLYL	1128	10	11	79
DARVCACLWM	733	10	18	129
DSVIDQNTCV	1454	10	12	86
DTLTCGFADL	124	10	12	86
ENALLWRQBA	2237	10	24	171
ETAGARLVVL	1342	10	12	86
FACLMGYIFL	130	10	11	79
FTEAMTRYSA	2792	10	14	100
GAARLAHGV	146	10	11	79
GADTAACGDI	992	10	12	86
GAGVAGALVA	1861	10	12	86
GALVGVVCA	1895	10	11	79

## HCV B58 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
GARLVLTATA	1346	10	11	79
GAQVNNRLI	1916	10	14	100
GSGKSTKPA	1238	10	12	86
GTVLOAETA	1335	10	14	100
HSKKKDELA	1400	10	14	100
IFASRGNHV	1925	10	14	100
ISGIYLAGL	1774	10	14	100
IIRVESNKV	2250	10	12	86
ITSCSNVSV	2818	10	14	100
ITYSTYKFL	1296	10	11	79
KSTKVPAAVA	1241	10	12	86
LADGGCSGA	1305	10	11	79
LAEOFKOKAL	1729	10	12	88
LALPPRAYAM	806	10	12	86
LSPGALWGV	1892	10	13	93
LSPRGSRPSW	88	10	11	79
LSRAPRWFM	3017	10	14	100
LSTLPGMPAI	1783	10	11	70
LTHPTIKYM	1842	10	16	114
NTCVTQVDF	1460	10	12	86
PAISPGALV	1888	10	12	86
PALSTGLIHL	888	10	12	86
PARLVFPDL	2609	10	11	79
PSWDMMKQL	1607	10	11	79
PTSGKSTRV	1236	10	13	93
PTHYVPESDA	1936	10	12	86
PTLHGPTPL	1621	10	11	79
PTLWARMILM	2870	10	22	157
PTLLYHLGA	1628	10	13	93
QAEAGARLV	1340	10	12	86
QAPPPSWDXM	1603	10	24	171
QATYCARAQ	1595	10	11	79
RAAKLDCTM	2757	10	16	114
RAAVCTRGVA	1186	10	11	79
RAAHGVRVL	148	10	14	100
SASQLSAPSL	2207	10	13	93
STKVPAAVA	1242	10	11	79
STWLVGGVL	1663	10	12	86
TAGARLVLA	1343	10	12	86
TARHTPNVSW	2852	10	11	79
TSCSNVSWA	2817	10	14	100
TSMITDPFHI	2177	10	13	93
TSTWLVGGV	1662	10	12	86
TTIMAKNEVF	2589	10	11	79
TTLPALSTGL	685	10	11	79

## ICV B58 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
VAATLGFGAY	1263	10	14	100
VTPGERPSGM	1507	10	16	114
VTRHADVPV	1138	10	11	79
WAOPGYWPL	76	10	12	86
WARMILMTHF	2873	10	12	86
WARPDYNPFL	2297	10	11	79
YAAQGYKVLV	1249	10	11	79
YSPGENRVA	2930	10	11	79
YSPCORVERL	2648	10	11	79
AARALNHGVRV	147	11	11	79
AASLRVFTTEAM	2788	11	12	88
AAVCTRGVAKA	1187	11	11	79
ASHLPYIEOGM	1717	11	14	100
ASCLSAFSLKA	2208	11	11	79
CARAQAAPPFSW	1599	11	11	79
CSFSIFLLALL	172	11	14	100
CTGSSSLV	1128	11	11	79
CTRGVAVAVDF	1190	11	11	79
DARVCACLWMM	733	11	16	114
DTLTCGFADLM	124	11	24	171
ETAGARLVVLA	1342	11	12	86
FADLMGYRLV	130	11	11	79
FSLSHSPGEI	2925	11	11	79
FTGLTHIDAHF	1567	11	13	93
FTLTPALSTGL	884	11	11	79
GADTAACGDII	992	11	12	86
GAGVAGALVAF	1861	11	12	86
GALVGVWCNA	1895	11	11	79
GAVQWNRILJA	1918	11	14	100
GSGKSTKVPAA	1238	11	12	86
HSKKKCDLAA	1400	11	14	100
HSYSPGENRV	2928	11	11	79
HTPVNSWLGNI	2855	11	12	86
ITRVESENKV	2250	11	12	86
ITSCSSNSVA	2816	11	14	100
ITYSTYKFLA	1296	11	11	79
KSTKVPAAAYAA	1241	11	11	79
LADGGGSGGAY	1305	11	11	79
LAGYGAGVACA	1857	11	11	79
LSNELLRRHWM	2479	11	14	100
LSPGALVGVW	1892	11	11	79
LTCGFADLMGY	126	11	12	86
LTSMLTDPISH	2178	11	13	93
NAVAYYRGLDV	1418	11	13	93
NTNRRPOONKE	14	11	11	79



ICV B58 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
PAILSPGALVV	1889	11	12	86
PSVAATLGFGA	1261	11	14	100
PTDPRRSRNL	109	11	12	86
PTHYVPESDAA	1936	11	12	86
PTLHGPTFLY	1621	11	11	79
PTPLLYRLGAV	1628	11	13	93
QAETAGARLVV	1340	11	12	86
QAPPSWDQMW	1603	11	11	79
QIVDFSLDPTF	1465	11	12	86
RSCPRGRORF	55	11	13	93
SADLEVVTSTW	1655	11	11	79
SSASQLSAPSL	2208	11	13	93
SSDLVYTRHA	1132	11	12	86
STWVLVGGVLA	1663	11	12	86
TARHIPVNSWL	2852	11	11	79
TSIGRDKNQV	1050	11	12	86
TSTWVLVGGV	1662	11	11	79
TLPLALSTGL	885	11	26	186
VAATLGFGAYM	1283	11	14	100
VAGALVAFKVM	1864	11	12	86
VAVEPWFSOM	974	11	11	79
VAYOATVCARA	1592	11	14	100
VAYRGLDVSV	1420	11	12	86
VTSTWVLVGGV	1661	11	12	86
WAQFGYPWPLY	78	11	12	86
WARMILMTHFF	2873	11	11	79
YAAQGYKVLVL	1249	11	12	86
YATQNLPGCSF	164	11	11	79
YTNVDDLVGV	1106	11	11	79

289

HCV B62 Super Motif Table XIV

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
AILSPGAL	1890	8	13	93
ALAHQVRV	150	8	14	100
ALGILQTA	1737	8	12	88
APTLWARM	2869	8	11	79
AOAPPPSW	1602	8	12	88
AGGYKVLV	1251	8	11	79
AVAYYRGL	1419	8	14	100
AVCTRGVA	1188	8	11	79
AVQWIMRL	1917	8	14	100
CLWIMILLI	739	8	12	86
CMSADLEV	1853	8	11	79
OODHLEPW	1556	8	12	86
CVTOTVDF	1462	8	12	88
DILAGYGA	1855	8	12	86
DLOGSVRL	279	8	12	86
DLNGYIPL	132	8	11	79
DLVNLIPA	1883	8	11	79
DOAETAGA	1339	8	12	86
EIPFYGKA	1377	8	13	93
EOPKOKAL	1731	8	12	86
EWTSTTW	1659	8	12	88
FISGIQYL	1773	8	14	100
FFDLGVRV	2615	8	11	79
FTGGGV	24	8	14	100
FOVAHLHA	1228	8	12	86
GIQYLGL	1778	8	14	100
GLRDLAVA	968	8	11	79
GPTLGVRV	41	8	13	93
GQNGGV	28	8	14	100
GVAGALVA	1863	8	12	86
GVAKAVDF	1193	8	11	79
GVLAALAA	1670	8	12	86
GVRVCEKM	2619	8	14	100
GVNCAIL	1800	8	11	79
HVGFGEKA	1910	8	11	79
HVSPTHV	1933	8	12	86
ILGGWVAA	1816	8	12	86
ILGIGTYL	1331	8	12	86
ILSPGALV	1891	8	13	93
IMAKNEVF	2581	8	12	86
IPFYGKAI	1378	8	13	93
IPLVGAPL	137	8	11	79
NOVOYLY	701	8	12	86
NPFDLGV	2613	8	11	79
IVGGVILL	30	8	13	93

## HCY B62 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
KMALYDVV	2625	8	12	86
KPARLIVF	2608	8	12	86
KQKAGLL	1734	8	12	86
KVPAAYAA	1244	8	11	79
LEANLLW	2235	8	12	86
LINTNGSW	414	8	11	79
LLALLSCL	178	8	12	86
LLAPITAY	1030	8	14	100
LLADARV	729	8	13	93
LLYRLGV	1629	8	13	93
LMGYPLV	133	8	11	79
LPALSTGL	687	8	14	100
LPGCSFSI	169	8	13	93
LPRRGFPL	37	8	13	93
LPVCOOHL	1553	8	13	93
LPYIEOGM	1720	8	12	86
LOCTMLV	2761	8	12	86
LVAYQATV	1591	8	12	86
LVDIILAGY	1853	8	11	79
LVGGVLA	1667	8	12	86
LVNPSVA	1257	8	14	100
LVNLLPAI	1884	8	11	79
LVTRHADV	1137	8	12	86
LVGVVCA	1897	8	11	79
LWICESA	2773	8	11	79
MLMTHFF	2878	8	12	86
MLTDPHSI	2179	8	14	100
NILGGWA	1815	8	12	86
NIVDVQTL	700	8	12	86
NLWFOEM	2239	8	12	86
NPSVAATL	1260	8	14	100
PLGGAARA	143	8	11	79
PLYRLGA	1628	8	13	93
PPSWDOOM	1805	8	12	86
PPSWDOMW	1606	8	11	79
PVHIGQPL	2318	8	11	79
QVGGVYL	29	8	13	93
QLLRPOA	336	8	12	86
QREYQEL	2808	8	11	79
QRYVWPL	78	8	12	86
RLHGLSAF	2918	8	12	86
RLIVFPDL	2611	8	11	79
RLAPITA	1029	8	12	86
RLVLATA	1347	8	12	86
RMAVDMMM	317	8	12	86

## HCV B62 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
RMILMTHF	2875	8	12	86
RPOYNPPL	2299	8	11	79
ROBMGGNI	2243	8	12	86
RVCERKIAL	2621	8	14	100
RVESENKV	2252	8	12	86
RVGGRHYV	2100	5	11	79
SIFLLALL	175	8	14	100
SLOPTFTI	1470	8	14	100
SPGENRV	2931	8	11	79
SPGRVEF	2649	8	11	79
SQLSAPSL	2209	8	13	93
SVAATLGF	1262	8	14	100
TIMAKNEV	2590	8	11	79
TLGFGAYM	1266	8	13	93
TLHGPTPL	1622	8	11	79
TLPGNPAL	1785	8	11	79
TLWARMIL	2871	8	11	79
TPCGSMIL	1875	8	12	86
TPGQVPCV	223	8	12	86
TQTVDFSL	1464	8	12	86
TVCARADA	1597	8	11	79
VIDQNTCV	1456	8	12	86
VLAALAAY	1671	8	12	86
VLCQCYDA	1521	8	13	93
VLDQAETA	1337	8	14	100
VLEGGVNY	157	8	12	86
VLNPSVAA	1258	8	14	100
VLGGMLA	1666	8	12	86
VLVNPVS	1258	8	14	100
VMGSSVGF	2639	8	11	79
VPESDAAA	1940	8	12	86
VQMMNRLL	1918	8	14	100
VWATDALM	1439	8	11	79
VVGWCAAA	1898	8	11	79
VVTSTWVL	1660	8	12	86
WMNRUJAF	1920	8	14	100
WPLLUILL	799	8	12	86
WLVGGVL	1665	8	12	86
YLAGLSTL	1779	8	14	100
YPYRLWHY	618	8	14	100
YPESDAAA	1939	8	12	86
AILSPGALV	1890	9	12	86
ALAHGVRVL	150	8	14	100
ALSTGLIHL	689	9	12	86
ALVVGWCA	1898	9	11	79

## HCY B62 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
APPSWDOM	1804	9	12	86
APTLWARM	2869	9	11	79
AOGYKVL	1251	9	11	79
AOGYWPL	77	9	12	86
AVOWMRL	1917	9	14	100
CMSADLEW	1653	9	11	79
DLCGSRLV	279	9	11	79
DLEVTSTW	1657	9	12	88
DLNGYPLV	132	9	11	79
DLNLLPAI	1883	9	11	79
DLWICESA	2772	9	11	78
DLYLVTRHA	1134	9	12	86
DPQLSDGSW	2410	9	11	79
DPRRSRL	111	9	12	88
EIPFYGKAI	1377	9	13	93
EMGNITRV	2245	9	12	86
EVTSTWVL	1658	9	12	86
FISGIOYLA	1773	8	14	100
FLLALSCL	177	8	12	86
FLLADARV	728	9	13	93
FQYSGQRY	2646	9	11	79
GIGYLDQA	1333	9	14	100
GLPVODHL	1552	9	13	93
GLRLAVAV	968	9	11	79
GLTHIDAHF	1569	9	13	93
GPGEAVOW	1912	8	12	86
GPTLLYRL	1625	9	14	100
GQVGGVYL	28	9	13	93
GVAGALVAF	1863	8	12	86
GVLAALAY	1670	8	12	86
GVNYATGNL	161	9	11	79
GVNCEKMA	2619	9	14	100
GVVLEDGV	154	8	13	93
HLHONVDV	696	9	12	86
HLPIEODM	1718	9	11	79
HMWNFISQI	1769	9	13	93
HONVNOY	688	9	11	79
HVGPGEAV	1910	9	11	79
ILAGYGAGV	1856	9	11	79
ILSPGALW	1891	9	13	93
KVLYLNPSV	1255	9	14	100
LITSCSSNV	2815	9	14	100
LWFPDLGV	2812	8	11	79
LLFLLADA	726	8	14	100
LLFNILGGW	1812	8	12	86

HCV B62 Super Motif (No binding data)

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
LLPRRGRL	36	9	13	93
LPAILSPGA	1888	8	13	93
LPALSTGLI	587	9	12	86
LPCEPEPV	2165	9	12	86
LPGCSFSIF	169	9	13	93
LVGGVLAAL	1667	9	12	86
LVLPNSVAA	1257	9	14	100
LVNLLPAIL	1884	9	11	79
LVTRHADVI	1137	9	11	79
LVGVVCAA	1897	9	11	79
NILGGWAA	1815	9	12	86
NIRTVRIT	1282	9	11	79
NIVDVOYLY	700	9	12	86
NLGKVIDTL	118	9	12	86
NLPCCSFSI	168	9	13	93
NVDDLGVW	1108	9	11	79
PLGGAARAL	143	8	11	79
PILYRLGAV	1628	9	13	93
PPPSWDOAMV	1605	9	11	79
PPVWHGQPL	2317	9	11	79
PPFVYDLI	2807	9	11	78
PVODHLEF	1534	9	12	86
PVNSWLGNI	2857	8	14	100
QVGGVYLL	29	9	13	93
QLSAPSLKA	2210	9	11	79
QPEYDLI	2808	9	11	79
QPGYFWPLY	78	9	12	86
QPRGRQPI	57	9	13	93
RLAPITAY	1029	9	12	86
RLMLMTHFF	2875	9	12	86
RVCEKMAFY	2621	9	14	100
RVESENKVV	2252	9	12	86
RMLDGWNY	156	9	12	86
SMLTDPSHI	2178	9	14	100
SPGALWGV	1893	9	13	93
SPGENRVA	2931	9	11	79
SPQORVERL	2649	9	11	79
SPRGSRPSW	99	9	11	79
SVDDNTGV	1455	9	12	86
TIMAKNEVF	2590	9	11	79
TLHGPTPLL	1622	9	11	79
TLPALSTGL	686	9	11	79
TLTCGFAOL	125	9	12	86
TLWARMILM	2871	9	11	79
TPLLYRLGA	1627	9	13	93

## HCV B62 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
TVLDOAETA	1336	9	14	100
VIOITCGF	122	9	12	86
VLEDGNYA	157	9	12	86
VLVDILAGY	1852	9	11	79
VLVGGVLA	1666	9	12	86
VLVLPNSA	1256	9	14	100
VOMNRLJA	1918	9	14	100
WGWCAAI	1898	9	11	79
WTSTWLV	1660	9	12	86
WMNRLAFA	1920	9	14	100
WMLVGGVLA	1665	8	12	86
YIPVGAFL	136	9	11	79
YLVAYCATV	1590	9	12	86
YLVTRHADV	1136	9	12	86
YOATVCARA	1594	8	13	93
YWGDLGSV	276	9	12	86
YWGVBRL	637	9	13	93
YVPESDAAA	1939	8	12	86
YLSPGALV	1880	10	12	86
ALWVGWCA	1896	10	11	79
APPSWDQNW	1604	10	11	79
APTLWARMIL	2869	10	11	79
AGPGYPWPLY	77	10	12	86
AVAYYRGLDV	1419	10	14	100
AVCTRGVAKA	1188	10	11	79
AVQWMNRLJA	1817	10	14	100
CLRKLGPPL	2941	10	12	86
CVTQTVD/SL	1462	10	12	86
DLAGYGAGV	1855	10	11	79
DLEVTSTW	1657	10	12	86
DLGVRVCEM	2617	10	13	93
DLSDGSWSTV	2412	10	11	79
DLVNLPAI	1883	10	11	79
DOAETAGARL	1339	10	12	86
DKWFGGQI	21	10	12	86
ELTSCSNV	2814	10	14	100
EDPKKALGL	1731	10	12	86
EWVSTWMLV	1659	10	12	86
GLSAFLHSY	2921	10	11	79
GLSTLPGNPA	1782	10	14	100
GLTHIDHFL	1569	10	13	93
GPGEQAVQWMA	1912	10	12	86
GVVGGVYLL	28	10	13	93
GVQWTVYHGA	1081	10	11	79
GVRVCEKMAI	2619	10	14	100

## HCY R62 Super Motif

Sequence	Position	Peptide No.	No. of Amino Acids	Sequence Frequency	Conservancy (%)
HONVDVYL	898		10	11	79
ILAGYGAGVA	1856		10	11	79
ILGGWVAQL	1818		10	12	86
IMAKNEVFCV	2591		10	11	79
KYLAGLSTL	1777		10	14	100
NFDLGVVRV	2813		10	11	79
KPTLHGTPPL	1620		10	11	79
KVIDTLGGF	121		10	12	86
KVLVLPNSVA	1255		10	14	100
LLFNLGWV	1812		10	12	86
LLFALSPGÁ	1887		10	13	93
LMGYPLVGA	133		10	11	79
LPAILSPGAL	1888		10	13	93
LPGCSFSIL	169		10	13	93
LPRGRRLGV	37		10	13	93
LPVCOHLEF	1553		10	12	86
LVAYQATVCA	1591		10	12	86
LVOLAGYGA	1853		10	11	79
LVGGVJAALA	1667		10	12	86
LVVGWVCAAI	1897		10	11	79
MLYDPSHITA	2179		10	14	100
NLPGCSFSIF	168		10	13	93
NPSVAATLGF	1260		10	14	100
PITYSTYKGF	1285		10	11	79
PLGGAARALA	143		10	11	79
POPEYDLBU	2807		10	11	79
PVCOHLEW	1554		10	12	86
PVNSWLGNII	2857		10	14	100
PVYCFTSPV	508		10	13	93
QJPCPEPQV	2164		10	12	86
QPEKGGKPA	2601		10	11	79
RLHLSAFSL	2918		10	11	79
RLVFPLGV	2611		10	11	79
RMWDMMNW	317		10	12	86
RLLEDGVYA	156		10	12	86
SLHSYSPGEI	2926		10	11	79
SLTGRDKQV	1051		10	12	86
SPGALVGVV	1893		10	11	79
SOLSAPSLKA	2208		10	11	79
SOPRRKQFI	56		10	13	93
SVAAATLGFGA	1262		10	14	100
TLHGPTPLLY	1622		10	11	79
TLFNILGGW	1811		10	12	86
TLPALSTGLI	686		10	11	79
TLTGGFADLM	125		10	12	86



## HCY B62 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
TPCTGSSDL	1126	10	11	79
TPLLYRLGAV	1627	10	13	93
TPNSNHLGNI	2856	10	12	86
TVDFSLDPTF	1486	10	12	86
VIDTLTCGFA	122	10	12	86
VLAALAAAYCL	1871	10	12	86
VLDQAEATAGA	1337	10	12	86
VLNPSVAATL	1258	10	14	100
VLITSCGNIL	2737	10	11	79
VLVGGVLAAL	1566	10	12	86
VLVLNPSVAA	1256	10	14	100
VMSSSYGFOY	2838	10	11	79
VPSDAAARV	1840	10	12	86
VQWMNRLJAF	1818	10	14	100
VGVVCAAIL	1898	10	11	79
WVLVGGVLAAL	1565	10	12	86
YKGGSSGGFL	35	10	13	93
YLLPHRGRL	1136	10	11	79
YLVTRHADVI	276	10	12	86
YGDLDSVF	1896	11	11	79
ALVGVVCAAI	1235	11	13	93
APTGGKSTKV	2889	11	11	79
APTLWARMILM	1602	11	12	86
QAAPPSPWDOM	1188	11	11	79
AVCTRGVAKAV	1917	11	14	100
AVQWMNRLJAF	1917	11	14	100
DILAGYGAGVA	1855	11	11	79
DLEVTISTWL	1657	11	12	86
DLGVRVCEUMA	2617	11	13	83
DLMGVYRLVGA	132	11	11	78
DLVLTTRHADV	1134	11	12	86
DQAEIAGARLV	1339	11	12	86
DWFRGGGQV	21	11	12	86
EQRKQKALGL	1731	11	12	86
FISGIDYLAGL	1773	11	14	100
FLADGGCSGGA	1304	11	11	79
FRGGQNGGV	24	11	14	100
FOYSRCQRIEF	2646	11	11	79
GVYLAGLSTL	1778	11	14	100
GLPQDQHLF	1552	11	12	86
GLSTLPGNPAI	1782	11	11	79
GPTLLYRLGA	1625	11	13	93
GPVYCFPSPV	507	11	13	93
GVLAALAAAYCL	1670	11	12	86
GVRVCEUMALY	2618	11	14	100

## HCV H62 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
GVRLEDGVNY	154	11	12	86
HLQNVIVQY	696	11	11	79
HMNVFSGCY	1769	11	13	93
HMNVIVQYLY	698	11	11	79
HYGPEGACW	1910	11	11	79
ILGGWVAQLA	1515	11	12	86
ILGISTVLDQA	1331	11	12	86
ILSPGALWGV	1891	11	13	93
KPARLIVFDL	2608	11	11	79
KPTLHGPTPL	1620	11	11	79
KOKALGLLOTA	1734	11	12	86
KVIDILTCGFA	121	11	12	86
KVLYLNP-SVAA	1255	11	14	100
LIAFASRGNHV	1824	11	14	100
LITSCSSNVSV	2815	11	14	100
LIVFDLGVHV	2612	11	11	79
LLFLLLADARV	726	11	13	93
LLNLGGWVA	1812	11	12	86
LLPAILSPGAL	1887	11	13	93
LLPRGPRGV	36	11	13	93
LLSPRGSPSW	87	11	11	79
LLWROBMGGV	2240	11	12	86
LPAILSPGALV	1888	11	12	86
LPALSTGLHL	687	11	12	86
LPGCSFSIFLL	169	11	13	93
LPVQDHLFPW	1553	11	12	86
LVGGVLAALAA	1667	11	12	86
LVLPNSVAATL	1257	11	14	100
LVTREHADVIV	1137	11	11	79
LVGVVVCAIL	1897	11	11	79
NILGGWVAQCL	1815	11	12	86
NITRVESENKV	2249	11	12	86
NLLPAILSPGA	1886	11	13	93
NLPGCSFSIFL	188	11	13	93
PITYSTYGKFL	1295	11	11	79
PLEGEGDPL	2403	11	11	79
PMGFSYDTRCF	2667..	11	11	79
PPSHDQMWKCL	1806	11	11	79
PVNSWLGNIIM	2857	11	12	86
PVYCFTPSPW	508	11	13	93
RMVGGVHFL	635	11	13	93
ROEMGGMTRV	2243	11	12	86
RVCXKALYDV	2821	11	12	86
SIFLLALLSQL	175	11	12	86
SMLTDPSHITA	2178	11	14	100

HCV B62 Super Motif

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)
SPTHVYPESDA	1935	11	12	86
SQLPCEPEOV	2163	11	12	86
SVAAITLFGAY	1262	11	14	100
TLGFGAYMSKA	1266	11	12	86
TLLENILGGWV	1811	11	12	86
TPCTCGSSDLY	1126	11	11	79
TPGLPVCOCHL	1550	11	13	93
TPVNSWLGNI	2856	11	12	88
TVLDCAETAGA	1336	11	12	86
VLCCEYDAGCA	1521	11	11	79
VLVDILAGYGA	1852	11	11	79
VLVGGVLAALA	1866	11	12	86
VOPEKGGKPKA	2600	11	11	79
VOWNNRLIAPA	1918	11	14	100
VWCAAILRRHV	1901	11	11	79
WVLVGGVLAAL	1655	11	12	88
YKGSSEGPLL	1165	11	12	86
YLVAYQATVCA	1590	11	12	86
YQATVCARAGA	1594	11	11	79
YWBQSGSVFL	276	11	12	88
YWPESDAAARV	1939	11	12	86

428

Table XV  
IICV Δ01 Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0101
ASFGSPY	166	8	20	100	
ONSVLSRKY	737	10	18	90	0.0001
FAAPFTOCGY	631	10	19	95	0.0680
GFAAPFTOCGY	630	11	19	95	
GREVLEY	140	8	15	75	
GYSNFMGY	579	9	17	85	
HTLWKAGILY	149	10	20	100	
KOAFIFSPTY	653	10	19	95	0.1100
LLDTASALY	30	9	17	85	0.0001
LSLDVSAIFY	415	10	19	95	12.0000
LTFGREVLEY	137	11	15	75	0.0150
MMWYWGPSLY	360	10	17	85	
MSTTLEAY	103	9	15	75	0.0810
NSVLSRKY	738	9	10	90	0.8500
PLOKGIKPY	124	9	20	100	0.0005
PLOKGIKPY	124	10	20	100	
PTTGRTSLY	797	9	17	85	0.1700
SASFGSPY	165	9	20	100	0.2100
SLOVSAIFY	416	9	19	85	5.2000
STTDLEAY	104	8	15	75	
TTGRTSLY	798	8	17	85	
WLSLDVSAIFY	414	11	19	95	
WMWYWGPS	359	11	17	85	
YPALMPLY	640	8	19	95	0.3200
YSLNFMGY	580	8	17	85	

25

Table XVI  
HCV Δ03 Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0301
AACWTRGER	647	10	12	86	0.0003
AARALAHGVR	147	10	11	79	
AATLGFGA	1264	8	14	100	
AATLGFGAY	1264	9	14	100	
AAVCTRGVA	1187	9	11	79	
AAVCTRGVAK	1187	10	11	79	
AAVCTRGVAKA	1187	11	11	79	
ACNWTRGER	648	9	12	86	
ADGGCSGGA	1306	9	11	79	
ADGGCSGGAY	1306	10	11	79	
ADVIPVRR	1142	8	12	86	
ADVIPVRRR	1142	9	11	79	
AFASRGNH	1926	8	14	100	
AGALVAFK	1865	8	12	86	
AGARLVVLA	1344	9	12	86	
AGARLVVLATA	1344	11	11	79	
AGLSTLPGNPA	1781	11	14	100	
AGVAGALVA	1862	9	12	86	
AGVAGALVAF	1862	10	12	86	
AGVAGALVAFK	1862	11	12	86	
AGWLLSPR	94	8	12	86	
AGWLLSPRGR	94	11	12	86	
AGYGAGVA	1858	8	12	86	
AGYGAGVAGA	1858	10	12	86	
ALGLLOTA	1737	8	12	86	
ALSTGLIH	689	8	12	86	
ALSTGLIHH	689	10	12	86	
ALVGVWCA	1896	8	11	79	
ALVGVWCAA	1896	10	11	79	
ASLMAFTA	1793	8	11	79	
ASQLSAPSLK	2208	10	11	79	
ASQLSAPSLKA	2208	11	11	79	
ASRGNHVSPH	1928	11	12	86	
ASSASQLSA	2204	10	14	100	
ATGNLPQCSF	165	10	13	93	
ATLGFGAY	1265	8	14	100	
ATLGFGAYMSK	1265	11	12	86	
ATRKTSER	48	8	11	79	
ATVCARAGA	1596	9	11	79	
AVCTRGVA	1188	8	11	79	
AVCTRGVAK	1188	9	11	79	
AVCTRGVAKA	1188	10	11	79	
AVQWMNRLIA	1917	10	14	100	0.0260
AVQWMNRLIAF	1917	11	14	100	
CAALIRRH	1903	8	13	93	

## HCV A03 Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0301
CANYELTPA	1530	9	11	79	
CGFADLMGY	120	9	13	93	
CGNLTICV	2742	8	11	79	
CGSSDLYLVR	1130	11	11	79	
CGYRRCRA	2727	8	14	100	
CLRLGVPLRL	2941	11	12	86	
CSFSIFLA	172	9	14	100	
CSSNVSA	2819	8	14	100	
CSSNVSAH	2819	9	12	86	
CTGSSDLY	1128	9	11	79	0.0001
CTRGVAKA	1190	8	11	79	
CTRGVAKAVDF	1190	11	11	79	
CTWMNSTGF	555	9	11	79	
CTWMNSTGFTK	555	11	11	79	
CVQPEKGG	2599	9	11	79	0.7600
CVQPEKGGK	2599	10	11	79	0.0008
CVTQTVDF	1462	8	11	79	0.0011
DAHFLSOTK	1574	9	12	86	
DLLVVCESA	2771	10	14	100	0.0003
DFSLDPTF	1460	8	11	79	
DGCGCGGA	1307	8	14	100	
DGCGCGGAY	1307	9	11	79	
DIICDECH	1310	9	11	79	
DILAGYGA	1855	8	12	86	
DILAGYGAGVA	1055	11	12	86	
DLGVRVCEK	2617	9	11	79	
DLGVRVCEKMA	2617	11	13	93	0.0003
DLMGVPLVGA	132	11	13	93	
DLVNLPLA	1803	11	11	79	
DLVVCESA	2772	8	11	79	
DLVLTTRH	1134	9	11	79	
DLVLTTRHA	1134	8	12	86	
DLTLCGFA	124	8	12	86	0.0003
DVIPVRRR	1143	8	12	86	
EAMTRYSA	2794	8	11	79	
ECYDAGCA	1524	8	14	100	
ECYDAGCAW	1524	10	11	79	
EDLVNLLPA	1882	9	11	79	
EGAVQWNR	1915	9	14	100	0.0004
EIPFYGKA	1377	8	13	93	
EMGGINTR	2245	8	12	86	
ETAGARLVLA	1342	11	12	86	
ETMRSPVF	1207	9	12	86	
EVFOVQPEK	2596	8	12	86	0.0008
FOVQPEKGGI	2588	10	11	79	

## HCV A03 Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0301
FCVPEKGGHK	2590	11	11	79	
FGAYMSKA	1269	8	12	86	
FGAYMSKAH	1268	9	12	86	
FGCTWNSITGF	553	11	11	79	
FGYGAKQVR	2554	9	12	86	
FISGIOYLA	1773	9	14	100	0.0008
FLADGCGSGA	1304	11	11	79	
FLLLADAR	728	8	14	100	
FSYDTRCF	2670	8	11	79	
FTEAMTRY	2792	8	14	100	
FTEAMTRYSA	2792	10	14	100	
FTGLTHIDA	1567	9	13	93	
FTGLTHIDAH	1567	10	13	93	
FTGLTHIDAHF	1567	11	13	93	
GAARALAH	146	8	11	79	
GAARALAHGVR	146	11	11	79	
GAGVAGALVA	1861	10	12	86	
GAGVAGALVAF	1861	11	12	86	
GAHWGVLA	350	8	12	86	
GALWGVVCA	1895	10	11	79	
GALWGVWCAA	1095	11	11	79	
GARLVLA	1345	8	12	86	
GARLVWLATA	1345	10	11	79	
GAQWNNR	1916	8	14	100	
GAQWNNRLA	1916	11	14	100	
GAYMSKAH	1270	8	12	86	
GCAWVELTPA	1529	10	11	79	
GCSFIFLLA	171	10	14	100	
GCTWNSITGF	554	10	11	79	
GDDLWICESA	2770	11	11	79	
GOLOGSVF	278	8	12	86	
GFADLMGY	129	8	13	93	
GFGAYMSK	1268	8	12	86	
GFGAYMSKA	1268	9	12	86	
GFGAYMSKAH	1268	10	12	86	
GFOYSPGOR	2645	9	11	79	
GFSYDTRCF	2669	9	11	79	
GGAARALA	145	8	11	79	
GGAARALAH	145	9	11	79	
GGCSGGAY	1308	8	11	79	
GGGCVGGW	26	10	14	100	
GGHYQMA	935	8	14	100	
GGQVGGW	27	9	14	100	
GGRIHFOH	1392	9	14	100	
GGRIHFOHSH	1392	11	14	100	0.0003

# YICV A03 Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0301
GGKPKARLVF	2505	11	11	79	
GGVLAALA	1669	8	12	86	
GGVLAALAA	1669	9	12	86	
GGVLAALAA	1669	10	12	86	
GGVLLPR	32	8	13	93	
GGVLLPRR	32	9	13	93	0.0003
GGVAAQLA	1818	9	12	86	
GIGTLDQA	1333	9	14	100	
GVLVPR	3037	8	11	79	
GLPVCOCH	1552	8	13	93	
GLPVCOCHLEF	1552	11	12	86	
GLPVSARR	1004	8	11	79	
GLRLAVA	988	8	11	79	
GLSAFSLH	2921	8	11	79	
GLSAFSLHSY	2921	10	11	79	0.0100
GLSTLPNPA	1782	10	14	100	
GLTHDAH	1569	8	13	93	
GLTHDAHF	1569	9	13	93	
GSGKSTKPPA	1238	10	12	86	
GSGKSTKPPAA	1238	11	12	86	
GSSDLVLTIR	1131	10	12	86	
GSSDLVLTIRH	1131	11	12	86	
GSSYGFOY	2641	8	11	79	
GTFFINAY	2063	8	11	79	
GTVDOAETA	1335	10	14	100	
GVAGALVA	1863	8	12	86	
GVAGALVAF	1863	9	12	86	
GVAGALVAFK	1863	10	12	86	0.3900
GVAKAVDF	1193	8	11	79	
GVQWTVYH	1081	8	11	79	
GVQWTVYHGA	1081	10	11	79	
GVGYLLPNR	3035	10	11	79	0.0014
GVLAALAA	1670	8	12	86	
GVLAALAA	1670	9	12	86	0.0046
GVRATRTSER	45	11	11	79	
GVRVCEKMA	2619	9	14	100	
GVRVCEKMA	2619	11	14	100	
GVRLEDGWY	154	11	12	86	
GVWCAAILR	1900	9	11	79	
GVWCAAILRR	1900	10	11	79	
GVWCAAILRRH	1900	11	11	79	
GVYLLPRR	33	8	13	93	
GVYLLPRRGP	33	11	13	93	
HAQVIVR	1141	8	11	79	
HAQVIVRR	1141	9	11	79	



## HCV A03 Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0301
HAQVIVRRR	1141	10	11	79	
HAPTGSCK	1234	8	14	100	
HAPTGSCKSTK	1234	11	13	93	
HGLSAFSLH	2920	9	11	79	
HGLSAFSLHSY	2920	11	11	79	
HGTPILLY	1624	8	11	79	
HGTPILLYR	1624	9	11	79	
HIDAI-FLSOTK	1572	11	14	100	
HLHAPTGSCK	1232	10	12	86	0.5900
HLHONIVDQY	696	11	11	79	
HLFCHSK	1395	8	14	100	
HLFCHSKK	1395	9	14	100	0.0250
HLFCHSKKK	1395	10	14	100	0.0260
HMWNFSGIQY	1769	11	13	93	
HSKKKCDLA	1400	10	14	100	
HSKKKCDLAA	1400	11	14	100	
HSVSGEINR	2928	10	11	79	
HTPGVPCVR	222	10	11	79	0.0004
HWGREGA	1910	8	11	79	0.0003
IAFASRGNH	1925	9	14	100	
IDAHLSTK	1573	10	14	100	
IDTLTCGF	123	8	12	86	
IDTLTCGFA	123	9	12	86	
IFCHSKK	1397	8	14	100	
IGTVLOOA	1334	8	14	100	
IGTVLDOAETA	1334	11	14	100	
IICDECH	1317	8	12	86	
ILAGYGAGVA	1656	10	11	79	
ILGGWVAA	1816	8	12	86	
ILGGWVAAQLA	1816	11	12	86	
ILGIGTVLOOA	1331	11	12	86	
IMAKNEVF	2591	8	12	86	
ISGIQYLA	1774	8	14	100	0.0150
ITRVESENK	2250	9	12	86	
ITSCSSNVSA	2816	11	14	100	
ITWGADTA	989	8	12	86	
ITWGADTAA	989	9	12	86	
ITYSTYCK	1296	9	12	86	
ITYSTYCKF	1296	9	12	86	
ITYSTYCKELA	1296	11	11	79	
NDVQVLY	701	8	12	86	
NFPDLGVR	2613	9	11	79	0.0036
IVGGVLLPR	30	10	13	93	0.0008
IVGGVLLPRR	30	11	13	93	
KALGLLOTA	1736	9	12	86	

## HCV A03 Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0301
KDELAAK	1404	8	12	86	
KFGYGAKDVR	2553	10	12	86	
KGGRHLIF	1391	8	11	79	
KGGRHLFCH	1391	10	11	79	
KGGRKPAR	2604	8	11	79	
KLGVPLR	2944	8	12	86	
KSTKVPAA	1241	8	12	86	
KSTKVPAAAY	1241	9	12	86	0.0009
KSTKVPAAAYA	1241	10	12	86	
KSTKVPAAAYAA	1241	11	11	79	
KTKRNTNR	10	8	12	86	
KTKRNTNRR	10	9	12	86	0.0110
KTSSRSQPR	51	9	13	93	0.1600
KTSSRSQPRGR	51	11	12	86	
KVIDILTCGF	121	10	12	86	
KVIDILTCGFA	121	11	12	86	
KVLVLPNSVA	1255	10	14	100	
KVLVLPNSVAA	1255	11	14	100	
KVPAAYAA	1244	8	11	79	
LADGGCSGGA	1305	10	11	79	
LADGGCSGGAY	1305	11	11	79	
LAQFKK	1729	8	12	86	
LAQFKKA	1729	9	12	86	
LAGYGAGVA	1057	9	11	79	
LAGYGAGVAGA	1057	11	11	79	
LCECYDAGCA	1522	10	11	79	
LDOAETAGA	1338	9	12	86	
LDOAETAGAR	1338	10	12	86	
LFLLADA	727	8	14	100	
LFLLADAR	727	9	14	100	
LFNLLGGWA	1813	10	12	86	
LFNLLGGWAA	1813	11	12	86	
LFTFSRR	290	8	11	79	0.0810
LGFGAYMSK	1267	9	12	86	
LGFGAYMSKA	1267	10	12	86	
LGFGAYMSKAH	1267	11	12	86	
LGGARALA	144	9	11	79	
LGGARALAH	144	10	11	79	
LGGWVAOLA	1817	10	12	86	
LIGTVLDOA	1332	10	13	93	
LGVFATRK	44	8	12	86	
LGVFATRK	44	8	14	100	
LGVFCEK	2618	8	14	100	
LGVFCEKMA	2618	10	14	100	
LAFASRGNH	1924	10	14	100	
LIEANLLWR	2235	9	12	86	0.0008



## ICY A03 Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0301
NFISGIQY	1772	8	14	100	
NFISGIQYLA	1772	10	14	100	
NGVCWTYV	1080	8	11	79	
NGVCWTYVH	1080	9	11	79	
NGVCWTYVHGA	1080	11	11	79	
NILGGWVA	1815	8	12	86	
NILGGWVAA	1815	9	12	86	0.0010
NITRVESENIK	2249	10	12	86	0.0005
NIVOVYLY	700	9	12	86	
NILPAILSPGA	1886	11	13	93	
NLPGCSFSIF	160	10	13	93	
NTCVTQTVD	1460	10	12	86	
NTNRPPQDK	14	10	11	79	
NTNRPPQDKF	14	11	11	79	
NTPLPVQDCH	1549	11	13	93	
PAILSPGA	1889	8	13	93	
PALSTGLIH	688	9	12	86	
PALSTGLIHLH	688	11	12	86	
PCSGSWLR	1976	8	11	79	
PCTGSSDLY	1127	10	11	79	
PDLGVRVCEK	2616	10	13	93	
PGALVVGWCA	1894	11	11	79	
PCCSFIF	170	8	14	100	
PCCSFISILLA	170	11	14	100	
PGCVPCVR	224	0	12	86	
PGEAGVQWMAIR	1913	11	13	93	
PGEINRVA	2932	0	11	79	
PGERPGMF	1509	9	12	86	
PGGGONGGVY	25	11	14	100	
PGLPVQDCH	1551	9	13	93	
PGYWFY	79	8	14	100	
PITYSTYK	1295	9	11	79	
PITYSTYKFK	1295	10	11	79	
PLGGAARA	143	8	11	79	
PLGGAARALAH	143	10	11	79	
PLLYRLGA	1628	8	13	93	
PMGFSYDTR	2667	9	11	79	
PMGFSYDTRCF	2667	11	11	79	
PSPVWGTIDR	514	11	13	93	
PSVAATLGF	1261	9	14	100	
PSVAATLGFGA	1261	11	14	100	
PSWDQWVK	1607	8	11	79	
PTDCFRKH	507	0	13	93	
PTDFRRSR	109	9	12	86	0.0006

# IICV A03 Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0301
PTGSGKSTK	1236	9	13	93	0.0002
PTHVVPESDA	1936	10	12	86	
PTHVVPESDAA	1836	11	12	86	
PTLHGPTILLY	1621	11	11	79	
PTPLLYRLGA	1626	10	13	93	
PVOCHLEF	1554	9	12	86	
PWVGTTOR	516	8	13	93	0.0008
QATAGAR	1340	8	12	86	
QATVCARA	1595	8	13	93	
QATVCARAQA	1595	10	11	79	
QWGGVILPR	29	11	13	93	
QLFTFSR	289	8	12	86	
QLFTFSRPR	289	9	11	79	0.7500
QLLRPQA	336	8	12	86	
QLSAPSLK	2210	8	11	79	
QLSAPSLKA	2210	9	11	79	
QTYDFSLDPTF	1465	11	12	86	
RAAVCTRGVA	1186	10	11	79	
RAAVCTRGVAK	1186	11	11	78	
RALAHGVR	149	8	14	100	
RATKTSE	47	9	11	79	
RGNHVSPTI	1930	9	12	86	0.0003
RGNHVSPTIY	1930	10	12	86	0.0003
RGPRLGVR	40	8	13	93	
RGPRLGVRA	40	9	13	93	
RGPRLGVRATR	40	11	11	79	
RGRROPIPK	59	9	13	93	0.0120
RGSLLSR	1154	8	12	86	
RGVAKAVDF	1192	9	11	79	
RLGVRATR	43	8	11	79	0.9400
RLGVRATR	43	9	11	79	
RLHGLSAF	2918	8	12	86	
RLHGLSAFSLH	2918	11	11	79	
RLIAFASR	1923	8	14	100	
RLIAFASRGNH	1923	11	14	100	
RLVFPQLGVR	2611	11	11	79	
RLAPITA	1029	8	12	86	
RLAPITAY	1029	9	12	86	2.7000
RLVVLATA	1347	8	12	86	
RLMLTHF	2875	8	12	86	
RLMLTHFF	2875	9	12	86	
RLWVGVEH	635	9	14	100	
RLWVGVEHR	635	10	14	100	0.7200
RSOPRGR	55	8	13	93	
RVCEKMALY	2621	9	14	100	0.1800

## HCV A03 Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0301
RVLEDGVNY	156	9	12	86	0.0120
RVLEDGVNYA	156	10	12	86	
SAFSLHSY	2023	8	11	79	
SASOLSAPSLK	2207	11	11	79	
SCSSNVSA	2018	9	14	100	
SCSSNVSAH	2018	10	12	86	
SDLVLTTR	1133	8	12	86	
SOLVLTTRH	1133	9	12	86	
SOLVLTTRHA	1133	10	12	86	
SFSIFLLA	173	8	14	100	
SGKSTKVP	1239	9	12	86	
SGKSTKVPAA	1239	10	12	86	
SGKSTKVPAAH	1239	11	12	86	
SMLTOPSH	2170	8	14	100	
SMLTOPSHITA	2170	11	14	100	
SSASQLSA	2206	8	14	100	
SSDLVLTTR	1132	9	12	86	0.0003
SSDLVLTTRH	1132	10	12	86	0.0003
SSDLVLTTRHA	1132	11	12	86	
SSNVSAH	2820	8	12	86	
SSASQLSA	2205	9	14	100	
STGLIHLH	691	8	12	86	
STKVPAAY	1242	8	12	86	
STKVPAAYA	1242	9	12	86	
STKVPAAYAA	1242	10	11	79	
SILPGNPA	1704	8	14	100	
STNPKPOR	2	8	11	79	
STNPKPORIK	2	9	11	79	
STNPKPORIKTK	2	11	11	79	
STWMLVGGVLA	1653	11	12	86	
STYGFALA	1299	8	12	86	
SVAATLGF	1262	8	14	100	
SVAATLGFGA	1262	10	14	100	
SVAATLGFAGAY	1262	11	14	100	
TAGARLWLA	1343	10	12	86	
TCGFADLMGY	127	10	13	93	
TCGSSDLY	1129	8	11	79	
TCVTOTVDF	1461	9	12	86	
TDPRIRESR	110	8	12	86	
TDPSSHITA	2101	8	14	100	
TGEIPFYGK	1375	9	11	79	
TGEIPFYGKA	1375	10	11	79	
TGLTHIDAH	1568	8	13	93	0.0003
TGLTHIDAHF	1568	9	13	93	
		10	13	93	

## HCV Δ03 Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0301
TGMLPGCSF	166	9	13	93	
TGSOKSTK	1237	8	13	93	
TGSOKSTKPA	1237	11	12	86	
TIMAKNEVF	2590	9	11	79	
TLGFGAYMSK	1266	10	12	86	0.0810
TLGFGAYMSKA	1266	11	12	86	
TLHGPTLLY	1622	10	11	79	0.0890
TLHGPTLLYR	1622	11	11	79	
TLPALSTGLIH	686	11	11	79	
TLWARMILMTH	2871	11	11	79	
TSCSSNVSA	2017	10	14	100	
TSCSSNVSAH	2017	11	12	86	
TSERSOPR	52	8	13	93	0.0003
TSERSOPTGR	52	10	12	86	
TSERSOPTGRH	52	11	12	86	
TSLTGRDK	1050	8	12	86	
TSMITDP SH	2177	9	13	93	0.0003
TTIMAKNEVF	2589	10	11	79	
TTMRSPVF	1208	8	12	86	
TVCARACA	1597	8	11	79	
TVDFSLDPTF	1466	10	12	86	
TVLDOAETA	1336	9	14	100	
TVLDOAETAGA	1306	11	12	86	
VAATLGFQA	1203	9	14	100	
VAATLGFQAY	1203	10	14	100	
VAGALVAF	1864	8	12	86	
VAGALVAFK	1864	9	12	86	
VAYQATVCA	1592	9	12	86	0.2400
VAYQATVCAR	1592	10	11	79	
VAYQATVCARA	1592	11	11	79	0.0005
VCAAILRR	1902	8	11	79	
VCAAILRRH	1902	9	11	79	
VCEKMALY	2622	8	14	100	
VOGPVYCF	505	8	13	93	
VOOCHLEF	1555	8	12	86	
VCTRGVAK	1189	8	11	79	
VCTRGVAKA	1189	9	11	79	
VQWTVYHGA	1082	9	11	79	
VDFSLDPTF	1467	9	14	100	
VDILAGYGA	1854	9	11	79	
VDYPRYLWH	614	9	13	93	
VDYPRYLWHY	614	10	13	93	
VFOVPEK	2597	8	12	86	
VFOVPEKGR	2597	11	11	79	
VFPDLGVR	2614	8	11	79	

## HCV Δ03 Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0301
VFTGLTHDA	1566	10	13	93	
VFTGLTHDAH	1566	11	13	93	
VGDLGVSF	277	9	12	86	
VGGLAALA	1668	9	12	86	
VGGLAALAA	1668	10	12	86	
VGGLAALAAAY	1668	11	12	86	
VGGLYLLPR	31	9	13	93	0.0003
VGGLYLLPRR	31	10	13	93	
VGGLYLLPR	3036	9	11	79	0.0007
VGWCAAILR	1809	10	11	79	
VGWCAAILRR	1809	11	11	79	
VIDTLTCGF	122	9	12	86	
VIDTLTCGFA	122	10	12	86	
VLAALAAAY	1671	8	12	86	
VLECEYDA	1521	11	11	79	
VLECEYDAGCA	1521	11	11	79	
VLDQAETA	1337	8	14	100	
VLDQAETAGA	1337	10	12	86	
VLDQAETAGAR	1337	11	12	86	
VLEDGVNY	157	8	12	86	
VLEDGVNYA	157	9	12	86	
VLENGVAA	1259	8	14	100	
VLTSMLTDP SH	2175	11	13	93	
VLVDILAGY	1852	9	11	79	
VLVDILAGYGA	1852	11	11	79	
VLVGGVLA	1666	8	12	86	
VLVGGVLA	1666	9	12	86	0.0003
VLVGGVLAALA	1666	11	12	86	0.0003
VLVLPNSVA	1256	9	14	100	
VLVLPNSVAA	1256	10	14	100	
VMGSSYGF	2639	8	11	79	
VMGSSYGFQY	2639	10	11	79	
VTRHADVPVR	1138	11	11	79	
VVCAAILR	1901	8	11	79	
VVCAAILRR	1901	9	11	79	
VVCAAILRRH	1901	10	11	79	
VVGWVCAA	1898	8	11	79	
VVGWVCAAILR	1898	11	11	79	
VVGTIDR	517	8	13	93	
WAGWLLSPR	93	9	12	86	
WAKHMWNF	1766	8	12	86	
WAQPGYPWPLY	76	11	12	86	
WARMILMTH	2873	9	12	86	
WARMILMTHF	2873	10	12	86	
WARMILMTHFF	2873	11	12	86	



## HCV A03 Motif with Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*0301
WGPTDPRR	107	8	12	86	
WGPTDPRRR	107	9	12	86	
WGPTDPRRISR	107	11	12	86	
WLLSPRGR	96	9	12	86	0.0008
WMNRLIAF	1920	8	14	100	
WMNRLIAFA	1920	9	14	100	0.0003
WMNRLIAFASR	1920	11	14	100	
WMNSTGFTK	557	9	11	79	0.0530
WVLVGGVLA	1665	9	12	86	
WVLVGGVLA	1665	10	12	86	
YATGNLPGCSF	164	11	12	86	
YDAGCAWY	1526	8	11	79	
YDIIICDECH	1315	10	12	86	
YGAGVAGA	1860	8	12	86	
YGAGVAGALVA	1860	11	12	86	
YGFQYSPQCR	2644	10	11	79	
YLPFRGPR	35	9	13	83	0.0054
YLVAYQATVCA	1590	11	12	86	
YSPGENR	2930	8	11	79	
YSPGENRVA	2930	10	11	79	
YSPQQRVEF	2648	9	11	79	
YSTYKFLA	1298	9	12	86	
YVGLOGSVF	276	10	12	86	
YGGVBRH	637	8	14	100	
YVPESDAA	1939	8	12	86	
YVPESDAAA	1939	9	12	86	
YVPESDAAR	1939	10	12	86	0.0003

567

Table XVII  
IICV AII Motif With Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*1101
AACNWTGR	647	10	12	86	0.0140
AARALAHGVR	147	10	11	79	
AATLFGAY	1264	9	14	100	
AAVCTRGVAK	1107	10	11	79	
ADNWTGR	648	9	12	86	
ADGCGSGAY	1306	10	11	79	
ADVIPVR	1142	8	12	86	
ADVIPVRR	1142	9	11	79	
AFASRGNH	1926	8	14	100	
AGALVAFK	1865	8	12	86	
AGVAGALVAFK	1862	11	12	86	
AGWLLSPR	94	8	12	86	
AGWLLSPRGR	94	11	12	86	
ALSTGLIH	689	8	12	86	
ALSTGLIHLH	609	10	12	86	0.0027
ASQLSAPSLK	2208	10	11	79	
ASRGNHVSPH	1928	11	12	86	
ATLFGAY	1265	8	14	100	
ATLFGAYMSK	1265	11	12	86	
ATRKTSER	48	8	11	79	
AVCTRGVAK	1188	9	11	79	0.0250
CAILLRH	1903	8	13	93	
CGFADLMGY	128	9	13	93	
CGNLTCY	2742	8	11	79	
CGSSDLYLVTR	1130	11	11	79	
CLRIKLGVPRLR	2941	11	12	86	
CNCSYPGH	304	9	11	79	
CNWTGR	649	8	12	86	
CSSNWSVAH	2819	9	12	86	
CTCGSSDLY	1128	9	11	79	0.0063
CTWVNSTGFTK	555	11	11	79	0.7500
CVQPKGR	2599	9	11	79	0.0005
CVQPKGGK	2599	10	11	79	0.0008
DAHFLSQTK	1574	9	14	100	0.0005
DGCGCGGAY	1307	9	11	79	
DIICDEH	1316	9	12	86	
DLGVRVCEK	2617	9	13	93	0.0002
DLYLVTRH	1134	8	12	86	
DVIPVRR	1143	8	11	79	
ECYDAGCAWY	1524	10	11	79	
EGAVQNMNR	1815	9	14	100	0.0014
EMGNITR	2245	8	12	86	
EVFOQPEK	2596	9	12	86	0.0270
FOVQPKGGK	2598	10	11	79	
FOVQPKGGK	2598	11	11	79	

## HCV A11 Motif With Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*1101
FGAYMSKAH	1269	9	12	86	
FGYGAQVR	2554	9	12	86	0.0005
ELLADAR	728	8	14	100	
FTEAMTRY	2792	8	14	100	
FTGLTHDAH	1567	10	13	93	
GAARALAH	146	8	11	79	
GAARALAHGVR	146	11	11	79	
GAVQWNR	1916	8	14	100	
GAYMSKAH	1270	8	12	86	
GFAULMGY	129	8	13	93	
GFGAYMSK	1200	8	12	86	
GFGAYMSKAH	1260	10	12	86	
GFOYSPQRI	2645	9	11	79	
GGAARALAH	145	9	11	79	
GGCSGGY	1308	8	11	79	
GGCQNGWY	26	10	14	100	
GGONGWY	27	9	14	100	0.0001
GGRLHFOH	1382	9	14	100	
GGRLHFOH	1392	11	14	100	
GGVLAALAA	1669	10	12	86	
GGVLLPR	32	8	13	93	
GGVLLPRR	32	9	13	93	0.0010
GYLLPNR	3037	8	11	79	
GLPVCOOH	1552	8	13	93	
GLPVSAAR	1004	8	11	79	
GLSAFSLH	2921	8	11	79	
GLSAFSLHSY	2921	10	11	79	0.0005
GLTHIDAH	1569	8	13	93	
GNHVSPTIY	1931	8	12	86	
GNHVSPTIY	1931	9	12	86	
GNITRESENK	2248	11	12	86	
GSSDLVLTIR	1131	10	12	86	
GSSDLVLTIRH	1131	11	12	86	
GSSYGFQY	2641	8	11	79	
GTFPINAY	2063	8	11	79	
GVAGALVAFK	1863	10	12	86	1.4000
GVCAWVYH	1081	8	11	79	
GVGYLLPNR	3035	10	11	79	0.0140
GVLAALAA	1670	9	12	86	0.0110
GVRATRTSER	45	11	11	79	
GVRCEKMALY	2619	11	14	100	
GVRLEDGVNY	154	11	12	86	
GVVCAAILR	1900	9	11	79	
GVVCAAILR	1900	10	11	79	
GVVCAAILRRH	1900	11	11	79	

## HCV A11 Motif With Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*1101
GVYLPRR	33	8	13	93	
GVYLPRRGR	33	11	13	93	
HADVPR	1141	8	11	79	
HADVPRR	1141	9	11	79	
HADVPRRR	1141	10	11	79	
HAPTSKG	1234	8	14	100	
HAPTSKGSTK	1234	11	13	93	
HGLSAFSLH	2920	9	11	79	
HGLSAFSLHSY	2920	11	11	79	
HGTPILLY	1624	8	11	79	
HGTPILLYR	1624	9	11	79	
HIDAFSLSTK	1572	11	14	100	
HLHAPTSKG	1232	10	12	86	0.0024
HLHONVDVQY	686	11	11	79	
HLIFCHSK	1385	8	14	100	
HLIFCHSKK	1385	9	14	100	0.0006
HLIFCHSKKK	1385	10	14	100	0.0002
HMINFSGIQY	1789	11	13	93	
HSYSGEINR	2928	10	11	79	
HTPGCVPCVR	222	10	11	79	
IAPASRGNH	1925	9	14	100	0.0012
IDAFSLSTK	1573	10	14	100	0.0003
IFCHSKKK	1397	8	14	100	
IICDECH	1317	8	12	86	
INTNGSWH	415	8	11	79	
ITRVESENK	2250	9	12	86	
ITYSTYCK	1286	8	12	86	0.0079
IVOVYLY	701	8	12	86	
IVPDGVR	2613	9	11	79	
IVGGVYLLPR	30	10	13	93	0.0044
IVGGVYLLPRR	30	11	13	93	0.0056
KDELAOK	1404	8	12	86	
KFGYGAQVR	2553	10	12	86	
KGGHLLFCH	1391	10	11	79	
KGGKPAR	2804	8	11	79	
KLGVPLR	2944	8	12	86	
KNEFOVOPEK	2594	11	11	79	
KSTKVPAAV	1241	9	12	86	0.0001
KTKRNTNR	10	8	12	86	
KTKRNTNR	10	9	12	86	
KTSERSOPR	51	9	13	93	0.0100
KTSERSQPRR	51	11	12	86	0.0640
LADGGCSGAY	1305	11	11	79	
LAOFKCK	1728	8	12	86	
LQAEATGAR	1338	10	12	86	

# HCY A11 Motif With Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*1101
LFLLADAR	727	9	14	100	
LTFSRR	290	8	11	79	
LGFGAYNSK	1267	9	12	86	0.2903
LGFGAYMSKAH	1287	11	12	86	
LGGAARALAH	144	10	11	79	
LGVRATRK	44	8	12	86	
LGVRVCEK	2618	8	14	100	
LIASFSGNH	1924	10	14	100	
LIENLLWR	2235	9	12	86	
LIFCHSCK	1396	8	14	100	0.0005
LIFCHSKK	1396	9	14	100	0.1900
LINTGSMH	414	9	11	79	
LVPDLGVR	2612	10	11	79	0.0001
LLAPITAY	1030	8	14	100	
LLFLLADAR	726	10	14	100	
LUPRRGPR	36	8	13	93	
LSFRGSR	97	8	12	86	
LSAFSLHSY	2922	9	11	79	
LSNSLLRH	2479	8	12	86	0.0002
LSNSLLRH	2479	9	12	86	0.0001
LSTGLJHL	690	9	12	86	
LTGCFADLMGY	126	11	12	86	
LTSMLTDPFH	2176	10	13	93	
LVAYOATVCAR	1591	11	11	79	
LVDILAGY	1053	8	11	79	
MGFSYDTR	2660	0	11	79	
MGSSYGFQY	2640	0	11	79	
MNRLIAFASR	1921	9	11	79	
MNSTGFTK	630	10	14	100	
MSTNPKPOR	1	8	11	79	
MSTNPKPOR	1	9	11	79	
NOGYRRCR	2726	10	11	79	
NCSYFGH	305	8	11	79	
NFISGIQY	1772	8	11	79	
NGVQWTVY	1080	8	14	100	
NGVQWTVYH	1080	8	11	79	
NITRVESENK	2249	9	11	79	
NIVDOYLY	700	10	12	86	0.0062
NTNRPQDVK	14	9	12	86	0.0140
NTNRPQDVK	14	10	11	79	0.0007
NTRGUPVQDCH	1549	11	13	93	
PALSTGLIH	688	9	12	86	
PALSTGLJHL	688	11	12	86	
PCSGSWLR	1976	8	11	79	
PCTGSSDLY	1127	10	11	79	
PDLGVRVCEK	2616	10	13	93	

HCV A11 Motif With Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*1101
PGCVPCVR	224	8	12	86	
PGGAVQMMN	1913	11	13	93	
PGGGWGGVY	25	11	14	100	
PLPVOODH	1551	9	13	93	
PGYRWPLY	79	8	14	100	
PITYSTYK	1295	9	11	79	
PLGGAARALAH	143	11	11	79	
PMGFSYDTR	2667	9	11	79	
PNIRITGVR	1281	8	13	93	
PSPVVGTTDR	514	11	13	93	
PSWDQMMK	1607	8	11	79	
PTDCFRKH	507	8	13	93	
PTDPRRHSR	109	9	12	86	
PTSGSKSTK	1236	9	13	93	0.0005
PTLHGTPLLY	1621	11	11	79	0.0001
PVVGTTDR	518	9	13	93	0.0005
QMETAGAR	1340	8	12	86	
QVGGVLLPR	29	11	13	93	
QLTFSPR	289	8	12	86	
QLTFSPRR	289	9	11	79	0.0330
QLSAPSLK	2210	8	11	79	
QNVDOY	699	8	11	79	
QNVDOYLY	699	10	11	79	
RAAVCTRGVAK	1100	11	11	79	
RALAHGVR	149	8	14	100	
RATRKTSER	47	9	11	79	
RGNVMSPTI	1930	9	12	86	
RGNVSPTHY	1930	10	12	86	0.0001
RGPRLGVR	40	8	13	93	0.0001
RGPRLGVRATR	40	11	11	79	0.0017
RGRROPIK	59	9	13	93	
RGSLSPR	1154	8	12	86	
RLGVRAIR	43	8	11	79	
RLGLSAFSLH	43	9	11	79	0.0290
RLIAFASR	2918	11	11	79	
RLIAFASR	1923	8	14	100	
RLIAFASRGNH	1923	11	14	100	
RLNVFPLDGVH	2611	11	11	79	
RLAPITAY	1029	9	12	86	0.0270
RMVGGVEH	635	9	14	100	
RMVGGVEH	635	10	14	100	0.0200
RNTNRPQDVK	13	11	11	79	
RSCPTGRR	55	8	13	93	
RVCEKMALY	2621	9	14	100	0.5000
RVLEDGVNY	156	9	12	86	0.0068

## HCV Δ11 Motif With Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*1101
SAFSLHSY	2923	8	11	79	
SASLSAPSLK	2207	11	11	79	
SCSSNVSAH	2818	10	12	86	
SDLYLVTR	1133	8	12	86	
SDLYLVTRH	1133	9	12	86	
SGKSTKVPAA	1239	11	12	86	
SMLTDPH	2178	8	14	100	
SNSLRH	2400	8	12	86	
SSDLYLVTR	1132	9	12	86	
SSDLYLVTRH	1132	10	12	86	0.0044
SSNVSAH	2020	8	12	86	0.0013
STGLIH	691	8	12	86	
STKVPAAY	1242	8	12	86	
STNPKOR	2	8	11	79	
STNPKORK	2	9	11	79	
STNPKORKTK	2	11	11	79	
SVAATLGFAY	1262	11	14	100	
TCGFADLMGY	127	10	13	93	
TCGSSDLY	1129	8	11	79	
TDPRHRSR	110	8	12	86	
TGEPFYGK	1375	9	11	79	
TGLTHIDAH	1588	9	13	93	0.0001
TGSGKSTK	1237	8	13	93	
TLGFGAYMSK	1268	10	12	86	0.0610
TLHGPTLLY	1622	10	11	79	0.0007
TLHGPTLLYR	1622	11	11	79	
TLPALSTGLIH	888	11	11	79	
TLWARMILMTH	2871	11	11	79	
TNPKORK	3	8	11	79	
TNPKORKTK	3	10	11	79	
TNPKORKTKR	3	11	11	79	
TNPKPOVK	15	9	11	79	
TSCSSNVSAH	2017	11	12	86	
TSERSQPR	52	8	13	93	0.0001
TSERSQPRGR	52	10	12	86	
TSERSQPRGRH	52	11	12	86	
TSLTGRDK	1050	8	12	86	
TSMITDFSH	2177	9	13	93	0.0001
VAATLGFAY	1263	10	14	100	
VAGALVAFK	1864	9	12	86	0.8900
VAYOATVCAH	1592	10	11	79	0.0038
VCAAILRR	1802	8	11	79	
VCAAILRRH	1902	9	11	79	
VCEKMALY	2622	8	14	100	
VCTRGVAK	1189	8	11	79	

## HCV All Motif With Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*1101
VDYPRYLWH	614	9	13	93	
VDYPRYLWH	614	10	13	93	
VFCVPEK	2597	8	12	86	
VFCVPEKGR	2597	11	11	79	
VFDLGR	2614	8	11	79	
VTGLTHDAH	1566	11	13	93	
VGGVLAALAA	1668	11	12	86	
VGGVLP	31	9	13	93	0.0019
VGGVLP	31	10	13	93	
VGYLLPNR	3036	9	11	79	0.0100
VGYVCAILR	1099	10	11	79	
VGVCAILRR	1099	11	11	79	
VLAALAA	1671	8	12	86	
VLDQAEAGAR	1337	11	12	86	
VLEDGWY	157	8	12	86	
VLTSMLTOPSH	2175	11	13	93	
VLVDLAGY	1052	9	11	79	
VNGSSYGFQY	2639	10	11	79	
VTRHADVIPR	1138	11	11	79	
VVCAILR	1901	8	11	79	
VVCAILRR	1901	9	11	79	
VVCAILRRH	1901	10	11	79	
VGVVCAILR	1098	11	11	79	
VVGTTDR	517	8	13	93	
WAGWLLSPR	93	9	12	86	
WAPGYPWPL	76	11	12	86	
WARMILMTH	2073	9	12	86	
WGTPDRR	107	8	12	86	
WGTPDRRR	107	9	12	86	
WGTPDRRRSR	107	11	12	86	
WLLSPRGR	96	11	12	86	0.0005
WMNRLAFASR	1920	11	14	100	0.0810
WMNSTGFTK	557	9	11	79	
WNFSGIQY	1771	9	14	100	
YDAGCAWY	1526	8	11	79	
YDIICDECH	1315	10	12	86	
YGFQSPGQR	2644	10	11	79	
YLLPRGPR	35	9	13	93	0.0005
YSPGEINR	2930	8	11	79	
YGGVGR	637	8	14	100	
YVPESDAAR	1939	10	12	86	0.0001
311		3			



Table XVIII  
ICV A24 Motif With Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*2401
AWDMIMNW	319	8	12	86	
AYAAQGYKVL	1248	10	11	79	0.0009
AYRGLDVSVI	1421	11	14	100	
CYDAGCAW	1525	8	11	79	
CYDAGCAWTEL	1525	11	11	79	
DFSLDPTF	1468	8	14	100	
DFSLDPTFI	1468	10	14	100	
FWAKHMWNF	1765	9	12	86	6.9000
FWAKHMWNI	1765	10	12	86	
GFADLMGYI	129	9	13	93	
GFADLMGYPL	129	11	11	79	
GFSDYTRCF	2669	9	11	79	
GWLLAPI	1027	8	11	79	
GYGAGVAGAL	1059	10	12	86	
GYPLVGAPL	135	10	11	79	
GYPRCRASGVL	2728	11	12	86	
HMWNFSGI	1769	9	13	93	0.0003
IFLLALLSCL	176	10	12	86	0.0057
IMAKNEVF	2591	8	12	86	
KFPGGGQ	23	8	13	93	
LNILGGW	1813	8	12	86	
LWARMILMTHF	2872	11	12	86	
LWROBMGN	2241	10	12	86	
LYLVTRHADV	1135	11	11	79	
MWNFSGI	1770	8	14	100	
MWNFSGIOYL	1770	11	14	100	
MYGGVEHRL	636	10	13	93	0.0270
NFSGIOYL	1772	9	14	100	0.0170
PMGFSYDTRCF	2667	11	11	79	
QPKKALGL	1732	9	12	86	
QPKKALGL	1732	10	12	86	
QWNNRLAF	1919	9	14	100	
QYLAGLSTL	1778	9	14	100	
QYSPQORVEF	2647	10	11	79	0.0480
QYSPQORVEFL	2647	11	11	79	0.0180
QYSPQORVEFL	2647	10	12	86	
FMADWIMMIMNW	317	8	12	86	
RMILMTHF	2875	9	12	86	
RMILMTHF	2875	11	13	93	
RMVGGVEHRL	635	9	14	100	
SFSIFLLAL	173	10	14	100	0.0041
SFSIFLLAL	173	9	14	100	
SMLTDPISH	2178	9	14	100	
SWDMWKKL	1608	11	11	79	
SYLKGSSGGPL	1164	11	12	86	
TWNNSTGF	556	8	11	79	

JICV Δ24 Motif With Binding Information

Sequence	Position	No. of Amino Acids	Sequence Frequency	Conservancy (%)	A*2401
TWLVGGVL	1654	9	12	86	
TYSTYGKF	1297	8	13	93	
TYSTYGKFL	1297	9	12	86..	0.0230
VFTGLTH	1566	8	13	93	
VMGSSYGF	2639	8	11	79	
VYLLPRGPRL	34	11	13	93	0.0016
WMNRLAF	1920	8	14	100	
YYRGLDVSVI	1422	10	14	100	
53		2			

Table XIX a  
ICV DR-Super Motif

Core Sequence	Core Freq.	Core Consistency (%)	Exemplary Sequence	Position in ICV Poly-protein	Exemplary Sequence Frequency	Exemplary Sequence Consistency (%)
FGAIVSKAI	12	86	TILGIVNSKAIQVO	1266	6	36
FGCTWVNST	12	86	GANFECTWVNSTGT	559	11	79
FKOKALGL	12	86	AEQFKOKALGLLOTA	1730	12	88
FLALLSCL	12	86	FSIFLALLSCLTVP	174	6	43
FDUGVRVC	11	79	UMFDUGVRVCERKA	2612	11	79
FOVAHLHAP	12	86	POIFOVANLHAPLOS	1225	8	43
FRAVCTRG	12	86	VOIFRANVCTRGVAK	1182	7	60
FSIFLLAL	14	100	QCFSIFLLALSCL	171	12	86
FSDFPFI	14	100	TVDFSLDPFIETT	1466	11	79
FTEAMTRY	14	100	UNFTEAMTRYISQPP	2769	7	50
FIPSPVWG	13	93	VYCFIPSPVWOTD	509	13	93
FTLLPALST	11	79	PCSTFTLLPALSTGI	681	9	64
FWANWVNF	12	86	LEVFWANWVNTGQ	1702	3	21
IONIFLSOT	14	100	LTIIDIAIFLSOIKOA	1570	7	50
IONTCVTO	12	86	DSWDCNCTCVTOVD	1464	12	86
IDILTCGFA	12	86	GRWIDILTCGFAULM	120	12	86
IEANLWRC	12	86	AULIEANLWRCDEM	2233	7	50
IFLLALLSC	14	100	SFSIFLLALLSCLTV	173	6	43
ILGWAAQ	12	86	LFNILGWAAQOLAP	1813	8	57
ILGISTYLD	12	86	STILGISTYLDQAE	1328	8	57
IPFIMQPG	11	79	CWALIPFIMQREGA	1303	11	79
ILSPGALVW	13	93	LPALLSPGALVQGV	1088	11	79
INVTTPC	12	86	TFINVTTPGTCIPS	2064	8	57
IFLVGAPLB	11	79	MDVIFLVGAPLGCA	134	10	71
ITVSEBKM	12	86	GGMITVSEBKNWAI	2247	10	71
ITSCSNVS	11	100	LELITSCSNVSAII	2813	11	79
INFRLOVNI	11	79	AILINFRLOVNIQCE	2610	11	79
LAANAYCL	12	86	QDVALANAYCLTTO	1668	8	57
LAQECQXQ	11	70	GRVALAQECQXQND	1302	10	71
LAQLSTPG	14	100	LOTVALAQLSTPGYNA	1777	14	100
LAGTONGVA	11	79	VALAGTONGVAKAL	1834	10	71
LATATPPGS	12	86	LWLATATPPGSVTV	1348	9	64
LOPITFIET	12	86	DFSLPITFIETTV	1468	5	38
LOCAETAGA	12	86	QTYLQCAETAGARLV	1335	12	86
LEUTSCSS	13	93	EYOLEUTSCSSNVS	2810	13	93
LEWITSTWV	12	86	SAOLEWITSTWLVG	1655	11	79
LELLADAR	14	100	VALLLELLADARVCS	724	4	29
LGQWVAOL	12	86	PHILGQWVAOLAPP	1814	8	57
LGGTVLDO	13	93	THLGGTVLDOAET	1329	9	64
LOVRAIRKT	12	86	GPFLGVRAIRKTSER	41	10	71
LOVRVCEWM	14	100	FFDLGVRCVCEWALLY	2615	11	79
UHLSNFSL	11	79	IERUHLSNFSLISY	2916	8	43
UKOPTILLY	11	79	KOTUKOPTILLYLG	1620	11	79
LHONVDVO	12	86	LHILHONVDVOTLY	694	10	71
UHSYSGEI	11	79	AFHSYSGEINIV	2924	11	79
UAFASQBN	14	100	MAFLUAFASQBNNS	1921	12	86
UEANLWIR	12	86	DAQUEANLWIRDEM	2332	7	50
UFCHSKK	14	100	GRUFCHSKKODE	1393	14	100
UTSCSNV	14	100	ELUTSCSNVSA	2012	13	93
LUALLSCLT	12	86	SIFLUALLSCLTIFA	175	5	36
LUFLUADA	14	100	TYWLUFLUADARVC	723	5	36
LUFLILGGW	12	86	ONLUFLILGGWAA	1609	4	29
LLADARVC	13	93	UULLLADARVCACL	726	9	64
LUPNLSFG	13	93	LVNLLUPNLSFOALV	1864	10	71

## HCV DR-Super Match

Core Sequence	Core Seq	Core Consistency (%)	Exemplary Sequence	Position In HCV Poly-protein	Exemplary Sequence Frequency	Exemplary Sequence Consistency (%)
LMGYPLVG	11	79	FAQLAGYPLVGAPL	130	11	79
LNPSVAATL	14	100	VLVNPSVAATLGFQ	1256	14	100
LPALSRGA	13	93	VHLLPALSRGALVV	1885	11	79
LPALSTGLI	12	86	FTLLPALSTGLIHL	684	11	79
LPFGPTG	13	93	VLLPFGPTGLGRIA	34	13	93
LPDLAVAE	11	79	INQLDLAVAEPPV	966	4	29
LPKLOVPL	12	86	ASCLNKLOVPLRW	2939	7	50
LSAFSLHSY	11	79	LLGLSAFSLHSYSG	2919	11	79
LSAPSLKAT	11	79	ASOLSAPSLKATCT	2208	7	50
LSNSLPHH	12	86	PNALSNSLPHHNAV	2476	4	29
LSFGALVQ	12	93	PAISFGALVGVWC	1889	11	79
LSPLLSIT	11	79	RSELSPLLSITTEWD	664	7	50
LSYKSPS	11	79	QWLSYKSPSTENSWAP	95	11	79
LSLGLRLH	12	86	LPALSLGLRLHION	687	10	71
LTCGFADLM	12	86	LDLTCGFADLMGYI	123	12	86
LTHDALFL	13	93	FTQLLTHDALFLSOL	1567	13	93
LTSMLTDS	13	93	VVALTSMLTDSIIT	2173	9	64
LVAYQATVC	12	86	FPYLVAAYQATVCATA	1508	9	64
LVQILAGY	11	79	GHLVQILAGYAGYV	1850	9	64
LVGGYLAAL	12	86	TYVLVGGYLAALAY	1694	12	86
LVLNPSVAA	14	100	YKVLNPSVAAATLO	1254	14	100
LVNLPLAIL	11	79	TEDLVNLPLAILSPG	1881	10	71
LVTHIDAVI	11	79	OLVLVTHIDAVIPVN	1134	11	79
LVGVVCAAA	11	79	FGALVGVVCAALIN	1084	11	79
LVGLATATP	12	86	QALVGLATATPTPOS	1345	11	79
LVATMLMT	12	86	ACTLVATMLMTITFF	2069	11	79
LVNDEMQN	12	86	AAHLVNDEMQNTHI	2238	12	86
LVNGLAVON	11	79	HYLVNGLAVONENVT	1627	9	64
LVNNEVFCV	12	86	THLVNNEVFCVDE	2509	12	86
LVNDMMNNW	12	86	CHLVNDMMNNWNSPT	315	12	86
LVNCGNITVE	11	79	PCNCGNITVESEN	2243	11	79
LVGYPLVDA	11	79	ADLVGYPLVGPLO	131	9	57
LVLDPSIIT	14	100	LTGALDPSIITAEI	2176	14	100
LVNRLAFAS	14	100	VQVNLRLAFASIGN	1918	14	100
LVNRYSAFG	14	100	TEVNLRYSAFGDPP	2193	10	71
LVNRFSGQ	14	100	AKLVNRFSGQOLA	1767	12	86
LVNNGVEH	14	100	KVNLNNGVEHRLNA	633	5	36
VAGALVAPK	12	86	GAGVAGALVAPKRVMS	1861	7	50
VAGLHAPTQ	12	86	TFQVLAGLHAPTQSK	1227	6	43
VATDALMTG	12	86	VYVATDALMTQYTG	1437	6	43
VAYDATVCA	12	86	PYLVAAYDATVCAHMO	1589	11	79
VCAILRRH	11	79	VQVCAILRRHNGP	1899	10	71
VCEKALYD	14	100	GVNVCCKALYDWS	2619	11	79
VCOOALEFW	12	86	GLPVCOOALEFWESV	1552	6	43
VCTRGVAKA	11	79	PAVCTRGVAKAVDF	1165	11	79
VFOVDFNG	12	86	RAGVFOVDFNGGGR	2594	10	71
VFDNNSRP	11	79	RSPTFDNNSRPNP	1211	10	71
VFTGLTHD	13	93	WESVFTGLTHDHF	1563	6	43
VGGVLAALA	12	86	WYLVGGVLAALAAVC	1665	12	86
VGGVYLPH	13	93	GGVGGVYLPHTKGP	28	13	93
VGSQPCPE	12	86	ONLVGSQPCPEPFO	2158	6	43
VGVVCAAIL	11	79	ALVGVVCAAILNTH	1898	11	79
VDCNCTCT	12	86	FDSVDCNCTCTIV	1453	12	86
VIDLTCGF	12	86	LQVIDLTCGFADL	119	11	79



Table XIXb. IICV DR Super Motif With Binding Data

Core Sequence	Exemplary Sequence	DT1	DT2-w2 1	DT2-w2 2	DT0	DT4-w4	DT4-w5	DT5-w11	DT5-w12	DR5-w19	DT8-w2	DT7	DT9	DT-w53
FGVAVSWI	TIGGAVYVSKVLDVD													
PGTWNST	GNVPGTWNSTGRT	0.0150	0.0120	0.0013		0.4200	0.0750	0.0210		0.0001	0.0035	0.0250	0.0270	
FKQKGLL	AEQFKQKGLGLOIA	0.0490				0.0006						0.0058		
FLUALLSCL	FSFLUALLSCLTVP													
FLQGVVVC	UVFPLQGVVVCERN	0.2400												
FOVAHLHAP	POIFOVAHLHAPTGS					0.0053						-0.0003		
FRVAVCTIQ	VGIFRAVAVCTIQVAK					0.0015						0.0030		
FSFLUALL	GCSEFSFLUALLSCL	0.0060				0.1600						0.0005		
FSLPITFI	TVPFSLPITFIETI	0.0001												
FEANRHS	UVFTEANRHSAPP					0.0920								
FTSPVWG	VYCFTPSPVWDTTD	0.0180	-0.0001	-0.0003			0.0570	0.0096		-0.0001	0.0035	0.0740	0.1800	
FTLPALST	PCSTFTLPALSTGU													
FWAHAMNE	LEVFWAHAMNEGSG													
IDAHFSOT	LTHDAHFSOTKOA					-0.0009								
DCNTCTVQ	DSVDCNTCTVDTVD	0.0001												
IDRLTQGA	GRYDILTQGFADLM													
IEANLLWFO	AQIEANLLWFOEAG													
IFLLALLSC	SFSIFLLALLSCLTV													
ILGGWYVAG	LVNIGGWYVAGLAP													
ILQIOTVD	STILQIOTVLOAE	0.0034				-0.0003						0.0017		
ILFHWGPG	CAMILFHWGPGEGA													
ILSFGLVW	LPAILSFGLVWGVW													
INAVTTQPC	TFNNAVTTQPCPTPS													
IPLVGAPLG	MOYILPVGAPLGQAA													
ITVSESNK	GGMITVSESNKVM													
ITSCSNWS	LELITSCSNWSVAH	0.0245	0.0200	-0.0003		0.0070	0.0360	0.0008		0.0510	-0.0003	0.0350	0.0330	
INFPOLGR	AFILNFPOLGRVCE	0.0053				0.0017						0.0004		
LALLANTCL	GGVALLANTCLTIG													
LAGGCGSG	CHPLAGGCGSGVND													
LAGLSTLPQ	IOYLAGLSTLPQVFA													
LAGYQAGVA	VOILAGYQAGVAGAL													
LATATPPGS	LVVLATATPPGSVTV	3.0000	0.0430	0.0094		3.9000		1.7000		-0.0001		0.0021	0.0550	
LDTFTIET	DFSLDTFTIETTV													
LDQAEIAGA	QIVLDQAEIAGATLV													
LEUTSCSS	EYOLEUTSCSSNWS	0.0001				0.0170						-0.0005		
LEVVTSTWV	SADLEVVTSTWVLVG													
LFLLADAR	WLLLFLLADARVCS	0.0240				0.0120								
LGWVWAGL	FNILGWWVWAGLAPP													
LGIOIYLDQ	ITILGIOIYLDQAEI													
LGVRATRT	GPILGVRATRTSER													
LGVRCEEM	EPILGVRCEEMVLY	0.0001				-0.0003								
LIHLSAEFL	REPLIHLAEFLSUSY													
LIHPTFLY	RPLIHLPTFLYRLG	0.0300				0.0010						-0.0002		
LIHNVDOVQ	LIHLIHNVDVOTLY											0.0055		
LIHSPGEE	AFSLIHSYSGENTV	0.0042				-0.0003								
LIHASTON	NRLIHLASTONVNS	0.0760	1.9000	0.0130	0.0058	0.0079	0.0550	0.4400		0.4600		0.0024	0.2400	
LIHNLWR	DALILHNLWRDEM	0.0008				-0.0010						0.1100		
LIHNSWK	GRILHNSWKDEE	0.0001				-0.0009						0.0025		
LIHSCSNW	DEUTSCSNWSVA											-0.0005		
LIALLSCLT	SIFLALLSCLTTPA													
LIHLLADA	YVLLLIHLLADARVC													
LIHRLGGW	QNTLLIHLGGWVMA													
LIHLDARVC	LIFLILHLDARVCACL													
LIHLLSPG	LVLLLIHLLSPGALV													
LIHVIPLVG	FADLIHVIPLVGAPL													

ICV DR Super Motif With Binding Data

Core Sequence	Exemplary Sequence	DR1	DR2w2.1	DR2w2.2	DR3	DR4w4	DR4w5	DR5w11	DR5w12	DR5w19	DR6w2	DR7	DR8	DRw53
LNPVAATL	VLPNPSVAATLGG	1.8000	0.0120	0.0004		2.1000	0.0035	0.0140		0.3100	0.0012	1.5000	3.2000	
LPALSPGA	VHLLPALSPGALVV	4.3000	0.0036	0.0016		0.0071		0.0130		0.0002		0.0400	0.0310	
LPALSTGL	FTLPLALSTGLHLL	0.0140	0.4000	0.0350		-0.0014		0.0120		0.0001		-0.0003	0.0032	
LPDLAVAVE	VHLLDLAVAVEPVV													
LPKLVPL	ASCLRLKLVPLLRW	1.0000	0.5000	0.0920	0.0051	0.0080	0.4900	0.0330	1.9000	0.0014	0.0730	0.0280	0.0087	
LSAFSLHSY	UHLGSAFSLHSYSPG	1.8000				0.0095						0.0070	0.0070	
LSAPSLKAT	ASOLSAPSLKATCTT	0.0150				0.0056						0.0006		
LSNLSLPHH	INALSNLSLPHHNAV													
LSPOLVWQ	PALSPGALVQWVC													
LSPLLSIT	RSLSPLLSITTEWO													
LSPLGSPS	GWLLSPGSPSAGP													
LSGLLH	LPALSTGLHLLTON													
LTGCFALM	IDLTCGCFALMGM	0.0017												
LTHDILFL	FTGLTHDILFLSOT	0.7500	0.5200	0.1300		0.0024	0.0010	0.0003		0.0002	0.0500	-0.0003	0.0056	
LTSMLTDS	VAVLTSMLTDSHIT					0.0005						0.1400		
LVAOATVC	FPYLVAOATVCNRA													
LVDLAGYO	GKVLVDLAGYOGV													
LVGGLAAL	TVVLVGGLAALAAV	0.7700	0.0011	-0.0003		0.0015		0.0009		0.0001		0.0570	0.0058	
LVLNPSVAA	YKVLNPSVAAILG													
LVHLLPAIL	TEDLVHLLPAILSPQ													
LVTHHDMV	DLVLVTHHDMVPR	0.0081	0.0220	0.0011		0.0016		0.0076		0.0005		0.0810	0.0820	
LWGVVCAA	PGALWGVVCAALU													
LVLATATP	GAILVLATATPGRS	0.0300	0.0009	0.0004		0.8000		0.0094		0.0004		0.0440	0.0067	
LWATMLAT	APTLWATMLATIFF													
LWPCBAGCN	APLLWPCBAGCNTH	0.7000										0.0022		
LYFLGAVON	TPLLYFLGAVONEVT					0.0018								
MAXNEVCOV	TIIMAXNEVCOVDE	0.0014				0.0016						0.0025		
MANJANANW	QPMANJANANWSPIT	0.0280	0.0015	0.0044		0.1000		0.0078		0.0000		0.0017	0.0230	
MOCCATIRE	POCCMOCCATIREN	0.0001				-0.0003						-0.0002		
MOPLUGA	ADLMOPLUGAPLG	0.0006				0.0060						0.0018		
NALTOPSHIT	LTSNALTOPSHITAE	0.0004				0.0740						-0.0003		
NHLLNFAS	VOMNHLLNFASSTGN													
NIRTSAPFG	TEAMIRTSAPFGDPP	1.5000	0.0150	0.0570		0.0040	0.0500	0.0076		0.0004	0.0180	0.2300	0.2700	
MMVFESQ	AGMMVFESQDCLA													
MYAGVBR	KVMMYAGVBRUNA													
VAGALVAFK	GAGVAGALVAFKVAS													
VAILHAPTO	TEQVAILHAPTOGSK													
VATDALMTO	VYVATDALMTOITO	0.0048	0.0047	0.0014	1.1000			0.0006		0.0029	0.0029	0.0400		
VATDATVCA	PYLVAATDATVCAAAQ													
VCAMLRRH	VGVVCAMLRRHNGP													
VCEKALYD	GVRVCEKALYDQVS	0.0022			0.0053	0.0012						-0.0002		
VCDHLEFW	GLPVCDHLEFWESV													
VCTRGVAKA	RAVCTRGVAKAVDF	0.0100				0.0077						0.0024		
VDCQKQ	WGVDCQKQKQKQK													
VFTONSPP	RSPVFTONSPPAMP													
VFTGLTHD	WESVFTGLTHDAHF													
VGVVLNLA	WVLGVVLNLAALAYC	0.0310				0.0068						0.0005		
VGVVLLPR	GQVGVVLLPRFPG													
VGSQLECP	QYLVGSQLECPFPG													
VGVVCAAL	ALVGVVCAALRRH													
VDCNTCVT	FDSVDCNTCVTQTY													
WDLTQGF	LGRVWDLTQGFADL													
VLAALAYC	VGVVLAALAYCLIT	0.0015				0.0096								
VLATATPPG	RLVVLATATPPGVSIT											0.0079		

## HCV DR Super Motif With Binding Data

Core Sequence	Exemplary Sequence	DR1	DR2w2 1	DR2w2 2	DR3	DR4w4	DR4w15	DR5w11	DR5w12	DR6w19	DR6w2	DR7	DR9	DRw53
VLEDOVNYA	OVTVLEDOVNYATON	0.0007				0.0006						-0.0002		
VLPNSVMAAT	KVLVLPNSVMAATLOF													
VLTSMJDP	DVALTSMJDPH													
VLTSCGNT	ASQVLTSCGNTLIC													
VLDVLAGY	LQKVLVDVLAGYAG													
VLDGVVLA	STWVLDGVVLAALAA													
VVLVLPVA	GKVLVLPVSAATL	1.1000	0.0260	0.0004	0.0900	9.5000	0.0670	0.1400	0.0320	0.6900	0.1700	0.2800	1.4000	
VNLPAIS	EDLVNLLPALSPGA	0.3700				0.0110						0.0015		
VPESDAAR	THVPESDAARVTO													
VLESTWLVQ	LEWVSTWLVGGVH	0.0120	0.0078	-0.0003		0.0280		0.0008		0.0046		0.1600	0.0120	
VVATDALMT	DVVVATDALMTGTT	0.0110	0.0110	-0.0003		0.0180	0.0072	-0.0004		0.0140	-0.0003	0.0910	-0.0025	
VVCAMLTR	VGVVCMALTRRNG													
VVGWVWCA	GALVWVWCAARTR	0.0170				0.0087						0.0043		
VVLATATPP	ATLVLATATPPQSV													
VVCFTPSV	CGPVVCFIPSPVWQ	0.2700	0.0025	-0.0003		0.2600	0.4600	0.0005		-0.0001	0.0011	0.2700	0.4300	
WAGNLLSPH	CGWVWGNLLSPQSH													
WATMLMIII	PLTWATMLMIIIFFS	0.0064				0.0200						0.0190		
WQADTAACD	ITWQADTAACDQDI													
WQFTDFEER	FFSWQFTDFEERSTN													
WNNRLAFA	AVQWNNRLAFAASRO	2.2000				0.6035								
WRLAPITA	SKGWRLAPITAYAQ	14.0000	0.0730	0.8800	-0.0006	2.1000	0.2500	4.2000	0.0290	-0.0001	0.9000	0.0205	0.0630	
WTGALIPC	SYTWGALIPCQAE	0.0260	0.0007	0.0015		0.0660	0.0220	0.0031		-0.0001	0.0130	0.0260	0.0750	
WTELTPAET	CGWVTELTPAETTVR													
YATGALGCC	GWVYATGALGCCSFS	0.0011				0.0130						-0.0003		
YCFITPSW	GPVYCFITPSWVWGT													
YDAGCAWYE	CECYDAGCAWYELTP													
YDMDDEC	GGVYDMDDECST													
YDLEUTSC	DFEYDLEUTSCSN	0.0003				0.0004						-0.0002		
YDGVAGAL	LQGVYDGVAGALVAF	0.0410				-0.0003								
YDGVSTQD	GSSVYDGVSTQDQAE	0.4600	0.0001	0.0300	0.0007	0.1200	0.0510	0.0010	0.0003	0.1800	0.0007	0.0008	1.1000	
YDGLQGB	YSVYDGLQGBQGBQ											0.1600		
YVGLVWFS	AGGVYVGLVWFSVAA	0.8400	0.0140	0.0004	0.0045	6.3000	0.1700	0.2700	0.0370	0.3900	0.2800	0.0300	0.2000	
YVGLSTUP	GQVYVGLSTUPQNP													
YVGLSOPF	PVYVGLSOPFPLC													
YVGLTPTIP	RVYVGLTPTIPTPLAT													
YVGLVQARA	LVAYVGLVQARQAP													
YVGLVNSI	VAVYVGLVNSISPTS													
YVGLVONE	PLLYVGLVONEVTL													
YVGLVSGV	NOGVYVGLVSGVALIT													
YVGLVDP	GVYVGLVDPQI													
YVGLVNEV	LVSYVGLVNEVASC													
YVGLVQSV	SAVYVGLVQSVFLV													
YVGLVPR					-0.0017									



## HCV DR Super Motif With Binding Data

Core Sequence	Exemplary Sequence	DR1	DR2-w2 1	DR2-w2 2	DR3	DR4-w4	DR4-w5	DR5-w11	DR5-w12	DR6-w19	DR6-w2	DR7	DR8
VLEDGVNVA	GVNLEDGVNVA	0.0007	0.0260	0.0004	0.0900	0.0006	0.0570	0.1400	0.0520	0.6900	0.1700	-0.0002	
VLPNSVAAT	KVLVLPNSVAATLGF												1.4000
VLTSMALTD	DVAVLTSMALTDPSHI											0.2800	
VLTSCGNT	ASQVLTSCGNTLTC											0.0015	
VLDLAGY	LGRVLDLAGYDAG												
VLGGVLA	STWVLGGVLAALAA	1.1000	0.0260	0.0004	0.0900	0.0110	0.0570	0.1400	0.0520	0.6900	0.1700	0.2800	
VLPNSVA	GYKVLVLPNSVAATL	0.3700											
VNLLPALS	EDLVNLLPALSPOA												
VPESDAAR	THVPESDAARVTO												
VTSTWLVG	LEVVTSTWLVGGVL		0.0078	-0.0003		0.0200		0.0008		0.0046		0.1600	0.0120
WATDALMT	DVWVWATDALMTGYT	0.0110	0.0110	-0.0003		0.0180	0.0072	-0.0004		0.0140	-0.0003	0.0910	-0.0025
VVCANLRR	VGVVCANLRRIRG												
VGVWCAM	GALVGVWCAMLRV	0.0170				0.0067						0.0043	
VVLATAPP	ANLVVLATAPPDSY		0.0025	-0.0003		0.2600	0.4000	0.0005		-0.0001	0.0011	0.2700	0.4300
VYCTPSPV	QCPVYCTPSPVWVG	0.2700											
WAGWLLSPR	GGWAGWLLSPRSGI	0.0064				0.0200						0.0190	
WAWLMLH	PLWAWLMLHTFFS												
WGAOTMAG	ITWGAOTMAGGDI												
WGPTDFRR	FFSWGPTDFRRSTN												
WKNRLNFA	AVKNWKNRLNFAFRO	2.2000				0.0035						0.0205	
WELLAPITA	SKGWELLAPITAYAO	14.0000	0.0730	0.8800	-0.0006	2.1000	0.2500	4.2000	0.0290	-0.0001	0.9000	0.0260	0.0530
WTOALTPC	STWTOALTPCAME	0.0260	0.0007	0.0015		0.0680	0.0220	0.0031		-0.0001	0.0130	0.4900	0.0750
WYELTPAET	GCWYELTPAETTVR												
YATGNLPQC	GVATGNLPQCSEFS	0.0011				0.0130						-0.0003	
YCFTPSPW	GFVYCFTPSPWVGT												
YDAGCAWYE	CECYDAGCAWYELIP												
YDNCDEC	GVYDNCDECST												
YDELTSC	DFEYDELTSCSSN	0.0003				0.0004						-0.0002	
YDAGVAGAL	LAGYDAGVAGALVAF	0.0410				-0.0003						0.0008	
YGFQVSGQ	QSSYGFQVSGQFME	0.4600	0.0001	0.0300	0.0007	0.1200	0.0510	0.0010	0.0003	0.1800	0.0007	0.1600	1.1000
YGRFLAQG	YSTYGRFLAQGCSQ												
YKVLVLP'S	AOQYKVLVLPNSVAA	0.8400	0.0140	0.0004	0.0045	6.3000	0.1700	0.2700	0.0370	0.5900	0.2800	0.0300	0.2000
YLAGLSTUP	GQVYLAGLSTUPNP												
YUGSSQGP	PVSYUGSSQGPFLC												
YLTROPFTTP	RVYLTROPFTTPRAR												
YQATVCARA	LVAYQATVCARADAP												
YRGLDWSI	VAYYRGLDWSIPTIS												
YRLGAVONE	PLYRLGAVONEVTL												
YTRQVASSV	NQVYTRQVASSVLT												
YSIEPLDLP	QACYSIEPLDLPOL												
YSKSEINIV	LHSYSKSEINIVASC												
YVGLCGSV	SNAYVGLCGSVFLV				-0.0017								
YVTLTPMNI													

Table XXb HCY DR 3A Motif With Binding Information

Core Sequence	Exemplary Sequence	DR3	DR1	DR2w201	DR2w202	DR4w4	DR4w15	DR5w11	DR5w12	DR6w19	DR7	DR8w2	DR9	DRw53
FLVGGCSG	YKRLADGGCGGAY													
FSLDPTFI	TVDFSLDPTFTIET		0.0001			0.1600					0.0005			
LEGEGDPO	MPLEGEFGDQSD	-0.0017												
LPCEPEPV	GSQLPCEPEPDVAVL	-0.0017												
MANQMMNW	GHRAVQDMANNSPT		0.0200	0.0015	0.0044	0.1600	0.0079	0.0080		0.0080	0.0017		0.0230	
MLTDPSSH	LTSMLTDPSSHAEI		0.0004			0.0740					-0.0003			
MSADLEWV	MACHSADLEWVSTW													
VATDALMTG	VVVVATDALMTGYG	1.1000	0.0048	0.0047	0.0014		0.0006	0.0029		0.0029	0.0400	0.0029		
VCDHLEFW	GLPVCDHLEFWESV	0.0063												
VFPDLGVR	FLVFPDLGVRCVK													
VFTDNSSP	RSPVFTDNSSPPAVP													
VLCECYDAG	DSSVLCECYDAGCAW													
VLEDGVNYA	GVVLEDGVNYATGN	-0.0017	0.0007			0.0086					-0.0002			
VLVDILAGY	LKVLVDILAGYAGG													
VQPSGGRK	VFCVQPSGGTGPAT													
YDLKITSC	QPEYDLKITSCSSN		0.0003			0.0004					-0.0002			
YSIEPLDP	GACYSIEPLDLPQII													
YVGLOGSV	SAMYVGLOGSVFLV	-0.0017												
YVPESDAAA	PTIYVPESDAAARVT	0.0220												

Table XXc HCV3B Motif

Core Sequence	Core Freq.	Core Consistency (%)	Exemplary Sequence	Position In HCV Poly-protein	Exemplary Sequence Frequency	Exemplary Sequence Consistency (%)
FQISWQCD	14	100	11FCHSKWCELA	1395	14	100
PSYDIQCD	11	79	PMDFSTDTDFESTV	2667	11	79
LAQFQKKA	12	86	GMQLAEQFKALGL	1726	8	57
UKPTUQFT	11	79	URLKPTUIGPTLL	1616	10	71
VRATKISE	11	79	PLGVATKTSERSQ	43	10	71
YLVTRHADV	12	86	SOLYLVTRHADVPV	1130	11	78
ASTNKKQK	11	79		1		

Table XXd IICV 3B Motif Binding Data

Core Sequence	Exemplary Sequence	DR1	DR2v201	DR2v202	DR3	DR4w4	DR4w15	DR5w11	DR5w12	DR6w10	DR6w2	DR7	DR9	DRw63
FGSKKKCD	HJFC-HSKKKCELA													
PSYDTKCFD	PMGFSTDTKCFDSTV				0.0190									
LAEDPKOKA	GMCLAEOPKOKALGL													
LUKPTLHGPT	URUKPTLHGPTILL													
VRATKTKTSE	RLGWATKTKTSEPSO				0.0022									
YLVTRHADV	SOLVLTTRHADVIV													
MSTNRKFOR														

7

TABLE XXI. Population coverage with combined HLA Supertypes

<u>HLA-SUPERTYPES</u>	<u>PHENOTYPIC FREQUENCY</u>					
	Caucasian	North American Black	Japanese	Chinese	Hispanic	Average
<u>a. Individual Supertypes</u>						
A2	45.8	39.0	42.4	45.9	43.0	43.2
A3	37.5	42.1	45.8	52.7	43.1	44.2
B7	38.6	52.7	48.8	35.5	47.1	44.7
A1	47.1	16.1	21.8	14.7	26.3	25.2
A24	23.9	38.9	58.6	40.1	38.3	40.0
B44	43.0	21.2	42.9	39.1	39.0	37.0
B27	28.4	26.1	13.3	13.9	35.3	23.4
B62	12.6	4.8	36.5	25.4	11.1	18.1
B58	10.0	25.1	1.6	9.0	5.9	10.3
<u>b. Combined Supertypes</u>						
A2, A3, B7	83.0	86.1	87.5	88.4	86.3	86.2
A2, A3, B7, A24, B44, A1	99.5	98.1	100.0	99.5	99.4	99.3
A2, A3, B7, A24, B44, A1, B27, B62, B58	99.9	99.6	100.0	99.8	99.9	99.8

SF 184895 v1

Table XXII  
IICY ANALOGS

AA	Sequence	Fixed Nomen.	A1 Motif	A2 Super Motif	A3 Super Motif	A24 Motif	B7 Super Motif	1° Anchor Fixer
9	RVXKMAly		N	N	Y	N	N	
9	AVXTRGVAK		N	N	Y	N	N	
8	EVXVOPEK		N	N	Y	N	N	
9	HLFXHSKK		N	N	Y	N	N	
9	LPGXSFSIF		N	N	Y	N	N	
9	LFXHSKKK		N	N	Y	N	N	
10	VLAALAAyXL		N	Y	N	N	N	No
10	HLFXHSKKK		N	N	Y	N	N	
10	AAXNWTRGER		N	N	Y	N	N	
10	YLLPRRGPRV	L2.LV10	N	Y	N	N	N	1
9	FGCSFSIF		N	N	N	N	Y	
9	LPVCSFSIF		N	N	N	N	Y	
9	LPGCSFSYF		N	N	N	N	Y	
9	LPGCMFSIF		N	N	N	N	Y	
9	LPGCSFSIF		N	N	N	N	Y	
8	LPGCSFSIF		N	N	N	N	Y	
9	PPVHGCPI		N	N	N	N	Y	
9	KPTLHGPTPI		N	N	N	N	Y	
10	APTLWARMII		N	N	N	N	Y	
10	SPRGSAPSI		N	N	N	N	Y	
9	LPRRGPRLGI		N	N	N	N	Y	
10	SPGQRFET		N	N	N	N	Y	
9	LPGCSFSII		N	N	N	N	Y	
9	DPRRHSINI		N	N	N	N	Y	
10	SPGALVVGVI		N	N	N	N	Y	
10	TPLLYRLGAI		N	N	N	N	Y	
9	TISGVLWQV		N	N	N	N	Y	
9	SISGVLWQV		N	Y	N	N	N	No
9	SLMAFTASV		N	Y	N	N	N	No
9	GLRDCITMLV		N	Y	N	N	N	No
10	KLVALGVNAV		N	Y	N	N	N	No
10	YLLPSRGPKL		N	Y	N	N	N	No
10	KLGLGLNAV		N	Y	N	N	N	No
10	YVLPFRGPRL		N	Y	N	N	N	No
10	VFNILGGWV		N	Y	N	N	N	Rev
10	KLVSIGVNAV	LV2.L10	N	Y	N	N	N	No
10	CINGYCWTA	12.VA9	N	Y	N	N	N	No
9	CANGVCWTV	1A2.V9	N	Y	N	N	N	Rev
9			N	Y	N	N	N	Rev

HCY ANALOGS

AA	Sequence	Fixed Nomen.	A1 Mott	A2 Super Mott	A3 Super Mott	A24 Mott	B7 Super Mott	I <sup>a</sup> Anchor Fixer
9	CWNGVCWAV 40		N	Y	N	N	N	

Table XXIII. Immunogenicity of identified supermotif-bearing peptides

Supermotif	Peptide	Sequence	Protein	Position	Human <sup>a</sup>				Transgenic mice <sup>b</sup>		
					Barnaba;		Chisari	Pape	overall	Frequency	Response
					patients	contacts					
A2	1073.05	LLFNILGGWV	NS4	1812	1/6	7/17	2/21	0/6	10/50	6/6	6.4 (1.7)
	1090.18	FLLADARV	NS1/E2	728	2/6	7/17	1/21	0/6	10/50	5/6	9.5 (3.0)
	1013.02	YLVAYQATV	NS4	1590	1/6	4/17	1/21	0/6	6/50	5/6	8.5 (3.7)
	1090.22	RLIVFPDLGV	NS5	2578	2/6	5/17	0/21	0/6	7/50	0/6	-
	1013.1002	DLMGYIPLV	Core	132	2/6	7/17	1/21	1/6	11/50	5/6	8.8 (2.6)
	24.0073	WMNRLIAFA	NS4	1920	1/6	3/17	2/21	1/6	7/50	0/6	-
	24.0075	VLVGGVLA	NS4	1666	1/6	6/17	3/21	1/6	11/50	0/6	-
	1174.08	HMWNFISGI	NS4	1769	3/6	3/17	2/21	0/6	8/50	6/6	6.4 (1.7)
	1073.06	ILAGYGAGV	NS4	1851	2/6	3/17	0/21	0/6	5/50	3/6	54.7 (3.3)
	1073.07	YLLPRRGPR	CORE	35	2/6	5/17	7/21	1/6	17/50	4/6	59.1 (7.2)
	24.0071	LLFLLADA	NS1/E2	726	2/6	9/17	0/21	0/6	11/50	0/6	-
	1.0119	YLVTRHADV	NS3	1131	6/6	10/17	0/21	1/6	17/50	0/6	-
A3	1.0952	KTSESRQPR	CORE	51	2/16	1/4	3/12	0/6	6/38	3/6	23.4 (1.3)
	1073.11	RLGVRATRK	CORE	43	4/16	1/4	7/12	1/6	13/38	3/6	42.2 (1.2)
	1.0955	QLFTFSPPR	ENV	290	1/16	0/4	6/12	1/6	8/38		
	1073.13	RMVYGGVEHR	NS1/E2	632	5/16	1/4	4/12	1/6	11/38	2/6	2.8 (1.1)
	1.0123	LIFCHSKKK	NS3	1396	6/16	1/4	4/12	2/6	13/38	3/6	4.4 (1.1)
	1073.10	GVAGALVAFK	NS4	1863	3/16	0/4	6/12	2/6	11/38	6/6	56.5 (1.7)
	24.0090	VAGALVAFK	NS4	1864	4/16	1/4	6/12	0/4	11/38	1/6	7.1
	24.0086	TLGFGAYMSK	NS3	1262	6/16		2/12	2/5	10/33		
	1145.12	LPGCSFSIF	CORE	169			2	3/10	5		
	B7										



Table XXIV. Human and murine MHC-peptide binding assays established using purified MHC molecules and gel filtration chromatography

A. Class I binding assays

			Radiolabeled peptide		Notes
Species	Antigen	Allele	Cell line	Source	
Human	A1	A*0101	Steinlin	Hu. J chain 102-110	YTAVVPLVY
	A2	A*0201	JY	HBVc 18-27 F6->Y	FLPSDYFPSV
	A2	A*0202	P815 (transfected)	HBVc 18-27 F6->Y	FLPSDYFPSV
	A2	A*0203	FUN	HBVc 18-27 F6->Y	FLPSDYFPSV
	A2	A*0206	CLA	HBVc 18-27 F6->Y	FLPSDYFPSV
	A2	A*0207	721.221 (transfected)	HBVc 18-27 F6->Y	FLPSDYFPSV
	A3		GM3107	non-natural (A3CON1)	KVFPYALINK
	A11		BVR	non-natural (A3CON1)	KVFPYALINK
	A24	A*2402	KAS116	non-natural (A24CON1)	AYIDNYNKF
	A31	A*3101	SPACH	non-natural (A3CON1)	KVFPYALINK
	A33	A*3301	LWAGS	non-natural (A3CON1)	KVFPYALINK
	A28/68	A*6801	CIR	HBVc 141-151 T7->Y	STLPETYVVR
	A28/68	A*6802	AMAI	HBV pol 646-654 C4->A	FTQAGYPAL
	B7	B*0702	GM3107	A2 signal seq. 5-13 (L7->Y)	APRTLVIYLL
	B8	B*0801	Steinlin	IIVgp 586-593 Y1->F, Q5->R 60s	FLKDYQLL
	B27	B*2705	LG2		FRYNGLIHR
	B35	B*3501	CIR, BVR	non-natural (B35CON2)	FPEKYAAAF
	B35	B*3502	TISI	non-natural (B35CON2)	FPEKYAAAF
	B35	B*3503	EHM	non-natural (B35CON2)	FPEKYAAAF
	B44	B*4403	PITOUT	EF-1 G6->Y	AEMGKYSFY
	B51		KAS116	non-natural (B35CON2)	FPEKYAAAF
	B53	B*5301	AMAI	non-natural (B35CON2)	FPEKYAAAF
	B54	B*5401	KT3	non-natural (B35CON2)	FPEKYAAAF
	Cw4	Cw*0401	CIR	non-natural (C4CON1)	QYDDAVYKL
	Cw6	Cw*0602	721.221 transfected	non-natural (C6CON1)	YRHDGNNVL
	Cw7	Cw*0702	721.221 transfected	non-natural (C6CON1)	YRHDGNNVL
Mouse	D <sup>b</sup>		EL4	Adenovirus E1A P7->Y	SGPSNTYPEI
	K <sup>b</sup>		EL4	VSV NP 52-59	RGYVFQGL
	D <sup>d</sup>		P815	HIV-IIIIB ENV G4->Y	RGPYRAFVTI
	K <sup>d</sup>		P815	non-natural (KdCON1)	KFNPMKTYI
	L <sup>d</sup>		P815	HBVs 28-39	IPQSLDSYWTSL

Table XXIV. Human and murine MHC-peptide binding assays established using purified MHC molecules and gel filtration chromatography

B. Class II binding assays				Radiolabeled peptide		Notes
Species	Antigen	Allele	Cell line	Source	Sequence	
Human	DR1	DRB1*0101	LG2	HA Y307-319	YPKYVKQNTLKLAT	optimal assay pH is 4.5
	DR2	DRB1*1501	L466.1	MBP 88-102Y	VVHFFKNIVTPRPY	
	DR2	DRB1*1601	L242.5	non-natural (760.16)	YAAFAAAKTAFAA	
	DR3	DRB1*0301	MAT	MT 65kD Y3-13	YKTIADFDEARR	
	DR4w4	DRB1*0401	Preiss	non-natural (717.01)	YARFOSQTTLKQKT	
	DR4w10	DRB1*0402	YAR	non-natural (717.10)	YARFQRQTTLKAAA	
	DR4w14	DRB1*0404	BJN 40	non-natural (717.01)	YARFOSQTTLKQKT	
	DR4w15	DRB1*0405	KT3	non-natural (717.01)	YARFOSQTTLKQKT	
	DR7	DRB1*0701	Pitout	Tet. tox. 830-843	QYIKANSKFIGITE	
	DR8	DRB1*0802	OLL	Tet. tox. 830-843	QYIKANSKFIGITE	
	DR8	DRB1*0803	LUY	Tet. tox. 830-843	QYIKANSKFIGITE	
	DR9	DRB1*0901	HID	Tet. tox. 830-843	QYIKANSKFIGITE	
	DR11	DRB1*1101	Sweig	Tet. tox. 830-843	QYIKANSKFIGITE	
	DR12	DRB1*1201	Herhuf	unknown eluted peptide	EALHQLKINPYVLS	no NEM in PI mix
	DR13	DRB1*1302	H0301	Tet. tox. 830-843 S->A	QYIKANAKFIGITE	
	DR31	DRB5*0101	GM3107 or L416.3	Tet. tox. 830-843	QYIKANAKFIGITE	
	DR51	DRB5*0201	L255.1	HA 307-319	PKYVKQNTLKLAT	
	DR52	DRB3*0101	MAT	Tet. tox. 1272-1284	NGQIGNDPNRDIL	
	DR53	DRB4*0101	L257.6	non-natural (717.01)	YARFOSQTTLKQKT	
	DQ3.1	DQA1*0301/DQB1*0301	PF	non-natural (ROIV)	YAHAAHAAHAAHAAHAA	
Mouse	IA <sup>b</sup>		DB27.4	non-natural (ROIV)	YAHAAHAAHAAHAAHAA	optimal assay pH is 5.5
	IA <sup>d</sup>		A20	non-natural (ROIV)	YAHAAHAAHAAHAAHAA	optimal assay pH is 5.0
	IA <sup>k</sup>		CH-12	HEL 46-61	YNTDGSYDYGILQNSR	
	IA <sup>s</sup>		LS102.9	non-natural (ROIV)	YAHAAHAAHAAHAAHAA	optimal assay pH is 5.0
	IA <sup>u</sup>		91.7	non-natural (ROIV)	YAHAAHAAHAAHAAHAA	
	IE <sup>d</sup>		A20	Lambda repressor 12-26	YLEDARRKKAIYEKKK	
	IE <sup>k</sup>		CH-12	Lambda repressor 12-26	YLEDARRKKAIYEKKK	

**Table XXV. Monoclonal antibodies used in MHC purification.**

Monoclonal antibody	Specificity
W6/32	HLA-class I
B123.2	HLA-B and C
IVD12	HLA-DQ
LB3.1	HLA-DR
M1/42	H-2 class I
28-14-8S	H-2 D <sup>b</sup> and L <sup>d</sup>
34-5-8S	H-2 D <sup>d</sup>
B8-24-3	H-2 K <sup>b</sup>
SF1-1.1.1	H-2 K <sup>d</sup>
Y-3	H-2 K <sup>b</sup>
10.3.6	H-2 IA <sup>k</sup>
14.4.4	H-2 IE <sup>d</sup> , IE <sup>k</sup>
MKD6	H-2 IA <sup>d</sup>
Y3JP	H-2 IA <sup>b</sup> , IA <sup>s</sup> , IA <sup>u</sup>

Table XXVI: HCV-derived conserved high algorithm A\*0201-binding peptides

Peptide	Molecule	1st Position	Sequence	Consv.	A2-supertype binding capacity (IC50 nM)					
					A*0201	A*0202	A*0203	A*0206	A*6802	A2 XRN
1073.05	NS4	1812	LLFNILGGWV	85	4.2	113	3.2	19	33	5
1090.18	NS1/E2	728	FLLADARV	92	18	90	149	247	111	5
1013.02	NS4	1590	YLVAYQATV	85	20	39	16	82	33	5
1090.22	NS5	2611	RLIVFPDLGV	79	56	391	10	370	8000	4
1013.1002	CORE	132	DLMGYIPLV	79	80	4778	204	481	12	4
24.0073	NS4	1920	WMNRLIAFA	100	122	130	3.3	1609	400	4
24.0075	NS4	1666	VLVGGVLAA	85	185	331	32	308	3077	4
1174.08	NS4	1769	HMWNFISGI	92	15	10750	77	132	7547	3
1073.06	NS4	1851	ILAGYGAGV	79	116	143	5.0	755	889	3
1073.07	CORE	35	YLLPRRGPRLL	92	125	6143	455	416	10256	3
24.0071	NS1/E2	726	LLFLLADA	100	217	287	455	3364	3077	3
1.0119	LORF	1131	YLVTRHADV	85	455	2048	3.6	71	3077	3
24.0065	NS4	1891	ILSPGALVV	92	238	10750	27	1028	3077	2
1013.12	NS1/E2	686	ALSTGLIHL	85	313	7167	45	18500	10256	2
939.14	NS1/E2	696	HLHQNVIVDV	85	500	3071	19	1370	10811	2
1090.21	NS5	2918	RLHGLSAFSL	79	179	782	625	18500	12500	1

Table XXVII: HCV-derived conserved high algorithm A\*03 and/or A\*11 binding peptides

Peptide	Molecule	1st Position	Sequence	Consv.	A3-supertype binding capacity (IC50 nM)						A3 XRN
					A*03	A*11	A*3101	A*3301	A*6801	A3	
1.0952	CORE	51	KTSESRQPR	92	69	94	67	1813	145	4	4
1073.11	CORE	43	RLGVRATRK	79	12	207	429	-	-	3	3
1.0955	ENV1	290	QLFTFSRR	79	15	182	621	3766	3	3	3
1073.13	NS1/E2	632	RMVYGGVEHR	100	15	300	95	9667	1778	3	3
1.0123	NS3	1396	LIFCHSKK	100	20	32	2535	24167	333	3	3
1073.10	NS4	1863	GVAGALVAFK	85	28	4	3273	26364	118	3	3
24.0090	NS4	1864	VAGALVAFK	85	46	7	3750	11600	258	3	3
24.0086	NS3	1262	LGFGAYMSK	85	136	21	2950	22308	222	3	3
1174.16	NS1/E2	557	WMNSTGFTK	79	208	74	12857	690	1429	2	2
1073.14	NS3	1261	TLGFGAYMSK	85	136	98	-	22308	8889	2	2
1090.23	LORF	1183	AVCTRGVAK	79	423	240	16364	-	-	2	2
1090.24	NS5	2596	EVFCVQPEK	85	13750	222	-	-	18	2	2
24.0103	NS1/E2	647	AACNWTGRER	85	36667	429	400	5273	4444	2	2
1073.16	NS3	1232	HLHAPTSGSK	85	19	2500	-	-	2857	1	1
1073.12	NS3	1395	HLIFCHSKK	100	423	-	20000	-	-	1	1
1090.26	NS3	1395	HLIFCHSKK	100	440	10000	-	-	8000	1	1

\* A dash indicates IC50nM &gt;30,000

Table XXVIII: HCV derived conserved B\*0702 binding peptides

## A. High conservancy 9- and 10-mer peptides.

Peptide	Molecule	1st Position	Sequence	Consv.	B7-supertype binding capacity (IC50 nM)					
					B*0702	B*3501	B*51	B*5301	B*5401	B7 XRN
1145.12	Core	169	LPGCSFSIF	92	28	90	100	114	6667	4
15.0048	E2	681	LPALSTGLI	85	157	-	2.8	1500	20000	2
15.0234	NS3	1620	KPTLHGPTPL	79	3.9	-	27500	-	-	1
15.0247	NS5	2835	APTLWARMIL	79	6.3	-	5500	-	-	1
15.0042	CORE	99	SPRGRPSW	79	14	-	11000	-	-	1
15.0039	Core	57	QPRGRRQPI	92	24	-	-	-	-	1
15.0218	Core	37	LPRRGPRLG	92	29	-	6111	-	4000	1
15.0060	NS5	2615	SPGQVEFL	79	46	-	27500	-	-	1
15.0043	Core	111	DPRRRSRNL	85	324	-	-	-	-	1
15.0063	NS5	2835	APTLWARM	79	344	-	4583	-	-	1
1292.17	NS5	2317	PPVHGCPL	79	393	-	-	-	-	1
15.0239	NS4	1893	SPGALVGVV	79	423	-	3438	-	-	1
15.0235	NS3	1621	TPLLRLGAV	92	458	-	6875	-	909	1

Table XXVIII: HCV derived conserved B\*0702 binding peptides

## B. Additional HCV derived B7 supermotif peptides.

Peptide	Molecule	1st Position	Sequence	Consv.	B7-supertype binding capacity (IC50 nM)					B7 XRN
					B*0702	B*3501	B*51	B*5301	B*5401	
29.0035	NS3	1378	IPFYGKAI	92	458	-	46	-	50	3
29.0040	Core	37	LPRRGPR	92	0.85	-	306	-	5000	2
29.0036	Core	137	IPLVGAPL	79	13	2250	79	-	2857	2
16.0187	NS1/E2	680	LPCSFTTLPA	64	423	24000	9167	-	15	2
29.0039	Core	169	LPGCSFSI	92	500	200	932	620	6250	2
15.0219	Core	142	APLGGAARAL	71	9.5	-	-	-	12500	1
29.0031	NS5	2869	APTLWARM	79	13	-	4583	-	4348	1
15.0231	NS3	1512	RPSGMFDSSV	71	153	-	-	-	-	1
29.0085	NS5	2474	LPINALNSL	57	220	18000	1170	-	11111	1
29.0037	NS5	2608	KPARLIVF	85	367	-	3235	-	16667	1
15.0237	NS4	1789	NPAIASLMAF	71	393	9000	5000	-	-	1
29.0118	NS5	2869	APTLWARMILM	79	423	-	-	-	3030	1
29.0042	NS4	1720	LPYIEQGM	85	423	-	1375	-	7692	1

## C. Engineered analogs of B7 supermotif peptides.

Peptide	Molecule	1st Position	Sequence	Consv.	B7-supertype binding capacity (IC50 nM)					B7 XRN
					B*0702	B*3501	B*51	B*5301	B*5401	
1145.12	Core	169	LPGCSFSIF	92	28	90	100	114	6667	4
1292.24	Core	169	LPGCSFSII		37	4364	5.3	262	1056	3
1145.13	Core	169	FPGCSFSIF		19	1.6	132	3.2	6.7	5

\* A dash indicates IC50 nM &gt;30,000.

**Table XXIX: HCV-derived A1- and A24-motif containing peptides****A. A1-motif peptides**

Peptide	Molecule	Position	Sequence	Conserv.	HLA-A*0101 binding (IC50 nM)
13.0019	NS5	2922	LSAFSLHSY	79	31
1.0509	NS5	2921	GLSAFSLHSY	79	61
1069.62	NS3	1128	CTCGSSDLY	79	68
24.0093	NS5	2129	EVDGVRLLHRY	100	167
13.0016	NS3	1241	KSTKVPAAAY	85	1923
1.0125	NS3	1525	CYDAGCAWY	79	4032
24.0008	E1	206	DCSNSSIVY	85	16667
24.0094	NS5	2720	TNSKGQNCGY	100	-
24.0096	NS3	1240	GKSTKVPAAAY	85	-
24.0100	NS3	1292	TGAPITYSTY	85	-
	NS3	1263	VAATLGFGAY	100	-
	NS5	2639	VMGSSYGFQY	79	-
	NS5	2640	MGSSYGFQY	79	-

A dash indicates IC50 nM &gt;25000

**B. A24 -motif peptides**

Peptide	Molecule	Position	Sequence	Conserv.	HLA-A*2402 binding (IC50 nM)
24.0092	NS4	1765	FWAKHMWNF	85	1.7
13.0075	NS4	1778	QYLAGLSTL	100	250
1073.18	NS1/E2	636	MYVGGVEHRL	92	444
13.0074	NS3	1297	TYSTYGKFL	85	522
13.0134	NS5	2647	QYSPGQRFVEF	79	667
24.0091	NS4	1772	NFISGIQYL	100	706
13.0131	Core	135	GYIPLVGAPL	79	2105
24.0108	Core	173	SFSIFLLALL	100	2927
13.0132	NS3	1248	AYAAQGYKVL	79	13333
13.0133	NS4	1859	GYGAGVAGAL	85	-
1174.08	NS4	1769	HMWNFISGI	93	-
	E1	317	RMAWDMMMNW	85	-
	NS1/E2	635	RMVVGVEHRL	93	-
	NS3	1422	YYRGLDVSVI	100	-
	NS3	1468	DFSLDPTFTI	100	-
	NS3	1608	SWDQMWKCL	79	-
	NS3	1664	TWVLVGGVL	85	-
	NS4	1732	QFKQKALGL	85	-
	NS4	1732	QFKQKALGLL	85	-
	NS4	1765	FWAKHMWNFI	85	-
	NS4	1919	QWMNRLIAF	100	-
	NS5	2241	LWRQEMGGNI	85	-
	NS5	2669	GFSYDTRCF	79	-
	NS5	2875	RMILMTHFF	85	-

A dash indicates IC50 nM &gt;25000



Table XXX: Immunogenicity of A2-supertype cross-reactive binders

Peptide	Immunogenicity									
	Human <sup>a</sup>					Transgenic mice <sup>b</sup>				
	Sequence	Protein	Position	Barnaba; Barnaba; contacts	Chisari	Pape	overall	Frequency	Response	
1073.05	LLFNILGGWV	NS4	1812	1/6	7/17	2/21	0/6	10/50	6/6	6.4 (1.7)
1090.18	FLLLADARV	NS1/E2	728	2/6	7/17	1/21	0/6	10/50	5/6	9.5 (3.0)
1013.02	YLVAYQATV	NS4	1590	1/6	4/17	1/21	0/6	6/50	5/6	8.5 (3.7)
1090.22	RLIVFPDLGV	NS5	2578	2/6	5/17	0/21	0/6	7/50	0/6	-
1013.1002	DLMGYIPLV	Core	132	2/6	7/17	1/21	1/6	11/50	5/6	8.8 (2.6)
24.0073	WMNRLJAFV	NS4	1920	1/6	3/17	2/21	1/6	7/50	0/6	-
24.0075	VLVGGVLA	NS4	1666	1/6	6/17	3/21	1/6	11/50	0/6	-
1174.08	HMWNFISGI	NS4	1769	3/6	3/17	2/21	0/6	8/50	6/6	6.4 (1.7)
1073.06	ILAGYGAGV	NS4	1851	2/6	3/17	0/21	0/6	5/50	3/6	54.7 (3.3)
1073.07	YLLPRRGPRV	CORE	35	2/6	5/17	7/21	1/6	17/50	4/6	59.1 (7.2)
24.0071	LLFLLADA	NS1/E2	726	2/6	9/17	0/21	0/6	11/50	0/6	-
1.0119	YLVTRHADV	NS3	1131	6/6	10/17	0/21	1/6	17/50	0/6	-

a. Data shown represents the number of positive responses over the total number of patients or contacts examined.

b. Frequency represents the number of positive responses over the total number of mice examined. Response indicates the average magnitude (standard deviation) of the response in positive animals, measured in lytic units.

Table XXXI: Immunogenicity of A3-supertype cross-reactive binders

Peptide	Sequence	Protein	Position	Immunogenicity					
				Human <sup>a</sup>			Transgenic mice <sup>b</sup>		
				Barnaba patients	Barnaba contacts	Chisari	Pape	overall	Frequency Response
1.0952	KTSESRQPR	CORE	51	2/16	1/4	3/12	0/6	6/38	3/6 23.4 (1.3)
1073.11	RLGVRATRK	CORE	43	4/16	1/4	7/12	1/6	13/38	3/6 42.2 (1.2)
1.0955	QLFTSPRR	ENV	290	1/16	0/4	6/12	1/6	8/38	
1073.13	RMYYVGGVEHR	NS1/E2	632	5/16	1/4	4/12	1/6	11/38	2/6 2.8 (1.1)
1.0123	LIFCHSKKK	NS3	1396	6/16	1/4	4/12	2/6	13/38	3/6 4.4 (1.1)
1073.10	GVAGALVAFK	NS4	1863	3/16	0/4	6/12	2/6	11/38	6/6 56.5 (1.7)
24.0090	VAGALVAFK	NS4	1864	4/16	1/4	6/12	0/4	11/38	1/6 7.1
24.0086	TLGFGAYMSK	NS3	1262	6/16		2/12	2/5	10/33	

a. Data shown represents the number of positive responses over the total number of patients or contacts examined.

b. Frequency represents the number of positive responses over the total number of mice examined. Response indicates the average magnitude (standard deviation) of the response in positive animals, measured in lytic units.

Table XXXII. Candidate HCV-derived HTL epitopes

Selection criteria	Peptide	Sequence	Source	Conservancy	
				Total	Core
A. DR-supermotif conserved 15mers	1283.01	GQIVGGVYLLPRRGPR	HCV Core 28	93	93
	1283.02	VYLLPRRGPRLGVR	HCV Core 34	93	93
	1283.03	GWLLSPRGSRPSWGPT	HCV Core 95	79	79
	1283.04	LGKVIDTLTCGFADL	HCV Core 119	79	86
	1283.05	IDTLTCGFADLMGYI	HCV Core 123	86	86
	1283.06	ADLMGYIPLVGAPLG	HCV Core 131	79	79
	1283.07	GVRVLEDGVNYATGN	HCV Core 154	86	86
	1283.08	GVNYATGNLPGCSFS	HCV Core 161	79	86
	1283.09	GCSFSIFLLALLSCL	HCV Core 171	86	100
	1283.10	GHRMAWDMMMNWSPT	HCV E1 315	86	86
	1283.11	CGPVYCFTPSPVVVG	HCV NS1/E2 506	93	93
	1283.12	VYCFTSPVVGTTD	HCV NS1/E2 509	93	93
	1283.13	GNWFGCTWMNSTGFT	HCV NS1/E2 550	79	86
	1283.14	FTTLPALSTGLIHLH	HCV NS1/E2 684	79	86
	1283.17	DLYLVTTRHADVIPVR	HCV NS3 1134	79	79
	1283.18	RAAVCTRGVAKAVDF	HCV NS3 1186	79	79
	1283.20	AQGYKVLVLNPSVAA	HCV NS3 1251	79	100
	1283.21	GYKVLVLNPSVAATL	HCV NS3 1253	100	100
	1283.22	VLVLNPSVAATLGFG	HCV NS3 1256	100	100
	1283.23	GTVLDQAETAGARLV	HCV NS3 1335	86	86
	1283.24	GARLVVLATATPPGS	HCV NS3 1345	79	86
	1283.25	GRHLIFCHSKKCCDE	HCV NS3 1393	100	100
	1283.27	DSVIDCNTCVTQTVD	HCV NS3 1454	86	86
	1283.28	TVDFSLDPTFTIETT	HCV NS3 1466	79	100
	1283.30	FTGLTHIDAHFLSQT	HCV NS3 1567	93	93
	1283.31	YLVAQATVCARAQA	HCV NS3 1591	79	93
	1283.32	KPTLHGPTPLLYRLG	HCV NS4 1620	79	79
	1283.33	LEVVTSTWVLVGGVL	HCV NS4 1658	86	86
	1283.34	TWVLVGGVLAALAAY	HCV NS4 1664	86	86
	1283.35	AEQFKQKALGLLQTA	HCV NS4 1730	86	86
	1283.40	PAILSPGALVVGVVCA	HCV NS4 1889	79	93
	1283.41	GALVVGVVCAAILRR	HCV NS4 1895	79	79
	1283.42	CAAILRRHVGPGEA	HCV NS4 1903	79	79
	1283.43	AVQWMNRLIAFASRG	HCV NS4 1917	100	100
	1283.44	MNRLIAFASRGNHVS	HCV NS4 1921	86	100
	1283.48	ANLLWRQEMGGNITR	HCV NS5 2238	86	86
	1283.49	RQEMGGNITRVESEN	HCV NS5 2243	86	86
	1283.52	ARLIVFPDLGVRVCE	HCV NS5 2610	79	79
	1283.53	FPDLGVRVCEKMALY	HCV NS5 2615	79	100
	1283.54	GVRVCEKMALYDVVS	HCV NS5 2619	79	100
	1283.56	QPEYDLELITSCSSN	HCV NS5 2808	79	93
	1283.57	LELITSCSSNVSAH	HCV NS5 2813	79	100
	1283.58	PTLWARMILMTHFFS	HCV NS5 2870	79	86
	1283.59	LHGLSAFSLHSYSPG	HCV NS5 2919	79	79
	1283.60	AFSLHSYSPGEINRV	HCV NS5 2924	79	79

Table XXXII. Candidate HCV-derived HTL epitopes

Selection criteria	Peptide	Sequence	Source	Conservancy	
				Total	Core
B. High algorithm conserved core	1283.15	VVLLFLLADARVCS	HCV NS1/E2 724	29	100
	1283.16	SKGWRLAPITAYAQ	HCV NS3 1025	29	79
	1283.19	PQTFQVAHLHAPTGS	HCV NS3 1225	43	85
	1283.26	DVVVVATDALMTGYT	HCV NS3 1436	43	79
	1283.29	WESVFTGLTHDAHF	HCV NS3 1563	43	92
	1283.45	LTSMLTDPSSHITAET	HCV NS5 2176	57	100
	1283.46	ASQLSAPSLKATCTT	HCV NS5 2208	50	79
	1283.47	DADLIEANLLWRQEM	HCV NS5 2232	50	85
	1283.50	SYTWTGALITPCAAE	HCV NS5 2456	64	79
	1283.51	TTIMAKNEVFCVQPE	HCV NS5 2589	64	85
	1283.55	GSSYGFQYSPQRVE	HCV NS5 2641	71	79
	1283.61	ASCLRLGVPLRVW	HCV NS5 2939	50	85
C. Collaborator	F098.03	AAYAAQGYKVLVLPNSVAAT	HCV NS3 1242-1261	71	100
	F098.04	GYKVLVLPNSVAATLGFAY	HCV NS3 1248-1267	100	
	F098.05	GYKVLVLPNSVAAT	HCV NS3 1248-1261	100	
	F134.01	RRPDVKFPGGGQIVGGVY	HCV Core 17-35	86	
	F134.02	DVKFPGGGQIVGGVYLLPRR	HCV Core 21-40	86	
	F134.03	GYKVLVLPNSVAATLGFAY	HCV NS3 1253-1272	100	
	F134.04	TLHGPTPLLYRLGAVQNEIT	HCV NS4 1622-1641		79
	F134.05	NFISGIQYLAGLSTLPGNPA	HCV NS4 1772-1791	100	
	F134.06	LLFNILGGWVAAQLAAPGAA	HCV NS4 1812-1831		86
	F134.07	GPGEQAVQWMNRLIAFASRG	HCV NS4 1912-1931	86	100
	F134.08	GEGAVQWMNRLIAFASRGNHV	HCV NS4 1914-1934	100	
	Pape 21	AIPLEVIKGRHLIFCHSKR	HCV NS3 1379-1398	21	100
	Pape 22	GRHLIFCHSKRKDELATKL	HCV NS3 1388-1407		100
	Pape 29	SVIDCNTCVTQTVDFSLDPT	HCV NS3 1450-1469	86	
D. DR3 motif	35.0102	GVRVLEDGVNYATGN	HCV 154	86	86
	35.0103	SAMYVGDLGCSVFLV	HCV 273	57	86
	35.0104	GHRMAWDMMNWSPT	HCV 315	86	86
	35.0105	SDLYLVRHADVIPV	HCV 1133	79	86
	35.0106	VVVVATDALMTGYTG	HCV 1437	42	86
	35.0107	TVDFSLDPTFTIETT	HCV 1466	79	100
	35.0108	DSSVLCECYDAGCAW	HCV 1518	71	93
	35.0109	GLPVCQDHLEFWESV	HCV 1552	42	86
	35.0110	GMQLAEQFKQKALGL	HCV 1726	57	86
	35.0111	PTHYVPESDAAARVT	HCV 1936	86	86
	35.0112	GSQLPCEPEPDVAVL	HCV 2162	64	86
	35.0113	LTSMLTDPSSHITAET	HCV 2176	57	100
	35.0114	MPPLEGEFGDPDLSD	HCV 2401	79	100
	35.0115	QPEYDLELITSCSSN	HCV 2808	79	93
	1283.25	GRHLIFCHSKKKCDE	HCV NS3 1393-1407		

Table XXXIII. HLA-DR screening panels

Screening Panel	Antigen	Alleles	Representative Assay		Phenotypic Frequencies						
			Allele	Alias	Cauc.	Blk.	Jpn.	Chn.	Hisp.	Avg.	
Primary	DR1	DRB1*0101-03	DRB1*0101	(DR1)	18.5	8.4	10.7	4.5	10.1	10.4	
	DR4	DRB1*0401-12	DRB1*0401	(DR4w4)	23.6	6.1	40.4	21.9	29.8	24.4	
	DR7	DRB1*0701-02	DRB1*0701	(DR7)	26.2	11.1	1.0	15.0	16.6	14.0	
	Panel total				59.6	24.5	49.3	38.7	51.1	44.6	
Secondary	DR2	DRB1*1501-03	DRB1*1501	(DR2w2 β1)	19.9	14.8	30.9	22.0	15.0	20.5	
	DR2	DRB5*0101	DRB5*0101	(DR2w2 β2)	-	-	-	-	-	-	
	DR9	DRB1*09011,09012	DRB1*0901	(DR9)	3.6	4.7	24.5	19.9	6.7	11.9	
	DR13	DRB1*1301-06	DRB1*1302	(DR6w19)	21.7	16.5	14.6	12.2	10.5	15.1	
	Panel total				42.0	33.9	61.0	48.9	30.5	43.2	
Tertiary	DR4	DRB1*0405	DRB1*0405	(DR4w15)	-	-	-	-	-	-	
	DR8	DRB1*0801-5	DRB1*0802	(DR8w2)	5.5	10.9	25.0	10.7	23.3	15.1	
	DR11	DRB1*1101-05	DRB1*1101	(DR5w11)	17.0	18.0	4.9	19.4	18.1	15.5	
	Panel total				22.0	27.8	29.2	29.0	39.0	29.4	
Quaternary	DR3	DRB1*0301-2	DRB1*0301	(DR3w17)	17.7	19.5	0.4	7.3	14.4	11.9	
	DR12	DRB1*1201-02	DRB1*1201	(DR5w12)	2.8	5.5	13.1	17.6	5.7	8.9	
	Panel total				20.2	24.4	13.5	24.2	19.7	20.4	

**Table XXXIV. HLA-DR binding capacity of target derived peptides: DR-supermotif and algorithm positive peptides.**

Peptide	Sequence	Source	Binding capacity (IC50 nM)										DR alleles	
			DR1	DR2w2B1	DR2w2B2	DR4w4	DR4w15	DR5w11	DR6w19	DR7	DR8w2	DR9	Lab	bound
1283.21	AAAYAAQGYKVLVLPNSVAATLFGGAY	HCV NS3 1242-1267												
1283.20	GYKVLVLPNSVAATL	HCV NS3 1253	4.5	350		5.2	567	143	5.1	89	288	54	175	9
	AQGYKVLVLPNSVAA	HCV NS3 1251	6.0	650		7.9	224	74	5.9	833	175	375	298	9
F98.03	AAAYAAQGYKVLVLPNSVAAT	HCV NS3 1242	2.9	48	483	18	1234	103	11	96	60	240		9
F98.05	GYKVLVLPNSVAAT	HCV NS3 1248-1261	1.4	39	3695	7.8	141	75	3.5	126	21	266		9
F98.04	GYKVLVLPNSVAATLFGGAY	HCV NS3 1248-1267	3.5	42	8154	9.7	1500	240	4.1	23	80	20		8
	GEAVQWMNRLIAFASRGNHVS	HCV NS4 1914-1935												
1283.44	MNRLIAFASRGNHVS	HCV NS4 1921	66	4.8			585	45	7.3	227	102	313	147	8
F134.08	GEAVQWMNRLIAFASRGNHV	HCV NS4 1914	3.2		182	361		345		221	158	6818		6
1283.16	SKGWRLAPITAYAQ	HCV NS3 1025	0.36	125	23	24	152	4.8		962	54	1190	384	8
1283.55	GSSYGFQYSPGQRYE	HCV NS5 2641	11		667	417	745	20000	19	156		68	571	7
1283.61	ASCLRLKLVPLRVW	HCV NS5 2939	5.0	16	217	6250	78	645	2500	862	671	8621	-	7
F134.05	NFISGIQYLAGLSTLPNPA	HCV NS4 1772	10		606	84		29			70	441		6

Shading indicates IC50 > 1 µM.

A dash (-) indicates IC50 > 20 µM.

**Table XXXV. HLA-DR binding capacity of 3 DR3 motif-containing peptides**

Peptide	Sequence	Source	DR3 binding (IC50 nM)
35.0106	VVVVATDALMTGYTG	HCV 1437	427
35.0107	TVDFSLDPTFTIETT	HCV 1466	235
1283.25	GRHLIFCHSKKKCDE	HCV NS3 1393	ND

Table XXXVIa: HCV-derived CTL epitope candidates

Peptide	Molecule	1st Position	Sequence	Consv.	Selection criteria
1073.05	NS4	1812	LLFNILGGVV	85	A2-supertype
1090.18	NS1/E2	728	FLLLADARV	92	A2-supertype
1013.02	NS4	1590	YLVAYQATV	85	A2-supertype
1090.22	NS5	2611	RLVFPDLGV	79	A2-supertype
1013.1002	CORE	132	DLMGYIPLV	79	A2-supertype
24.0073	NS4	1920	WMNRLIAFA	100	A2-supertype
24.0075	NS4	1666	VLVGGVLA	85	A2-supertype
1174.08	NS4	1769	HMWNFISGI	92	A2-supertype
1073.06	NS4	1851	ILAGYGAGV	79	A2-supertype
1073.07	CORE	35	YLLPRRGPR	92	A2-supertype
24.0071	NS1/E2	726	LLFLLADA	100	A2-supertype
1.0119	LORF	1131	YLVTRHADV	85	A2-supertype
1.0952	CORE	51	KTSSRSQPR	92	A3-supertype
1073.11	CORE	43	RLGVRATRK	79	A3-supertype
1.0955	ENV1	290	QLFTFSRR	79	A3-supertype
1073.13	NS1/E2	632	RMVVGVEHR	100	A3-supertype
1.0123	NS3	1396	LIFCHSKKK	100	A3-supertype
1073.10	NS4	1863	GVAGALVAFK	85	A3-supertype
24.0090	NS4	1864	VAGALVAFK	85	A3-supertype
24.0086	NS3	1262	TLGFGAYMSK	85	A3-supertype
F104.01	NS5	3003	VGIYLLPNR	79	A31
1145.12	Core	169	LPGCSFSIF	92	B7-supertype
29.0035	NS3	1378	IPFYGKAI	92	B7-supertype
13.0019	NS5	2922	LSAFSLHSY	79	A1
1069.62	NS3	1128	CTCGSSDLY	79	A1
24.0092	NS4	1765	FWAKHWNF	85	A24



**Table XXXVlb: HCV-derived HTL epitope candidates**

Region	Peptide	Motif <sup>1</sup>	Sequence
HCV NS3 1025-1039	1283.16	DR	SKGWRL LAPITAYAQ
HCV NS3 1242-1267	F98.03	DR	AAYAAQGYKVLVLPNSVAAT
HCV NS3 1393-1407	1283.25	DR3	GRHLIFCHSKKKCDE
HCV NS3 1437-1451	35.0106	DR3	VVVVATDALMTGYTG
HCV NS3 1466-1480	35.0107	DR3	TVDFSLDPTFTIETT
HCV NS4 1772-1790	F134.05	DR	NFISGIQYLAGLSTLPGNPA
HCV NS4 1914-1935	F134.08	DR	GEGAVQWMNRLIAFASRGNHV
HCV NS5 2641-2655	1283.55	DR	GSSYGFQYSPGQRE
HCV NS5 2939-2953	1283.61	DR	ASCLRKLGVPPPLRVW

1. Peptides identified on the basis of either the DR P1-P6 supermotif or by use of the DR1-4-7 algorithms are indicated by 'DR'. Peptides identified using the DR3 motif are indicated by 'DR3'.

**Table XXXVII. Estimated population coverage by a panel of HCV derived HTL epitopes**

Antigen	Alleles	Representative assay	No. of epitopes <sup>2</sup>	Population coverage (phenotypic frequency)					
				Cauc.	Blk.	Jpn.	Chn.	Hisp.	Avg.
DR1	DRB1*0101-03	DR1	6	18.5	8.4	10.7	4.5	10.1	10.4
DR2	DRB1*1501-03	DR2w2 β1	3	19.9	14.8	30.9	22.0	15.0	20.5
DR2	DRB5*0101	DR2w2 β2	6	-	-	-	-	-	-
DR3	DRB1*0301-2	DR3	2	17.7	19.5	0.40	7.3	14.4	11.9
DR4	DRB1*0401-12	DR4w4	5	23.6	6.1	40.4	21.9	29.8	24.4
DR4	DRB1*0401-12	DR4w15	3	-	-	-	-	-	-
DR7	DRB1*0701-02	DR7	5	26.2	11.1	1.0	15.0	16.6	14.0
DR8	DRB1*0801-5	DR8w2	5	5.5	10.9	25.0	10.7	23.3	15.1
DR9	DRB1*09011,09012	DR9	3	3.6	4.7	24.5	19.9	6.7	11.9
DR11	DRB1*1101-05	DR5w11	5	17.0	18.0	4.9	19.4	18.1	15.5
DR13	DRB1*1301-06	DR6w19	2	21.7	16.5	14.6	12.2	10.5	15.1
Total <sup>1</sup>				98.5	95.1	97.1	91.3	94.3	95.1

1. Total population coverage has been adjusted to account for the presence of DRX in many ethnic populations. It has been assumed that the range of specificities represented by DRX alleles will mirror those of previously characterized HLA-DR alleles. The proportion of DRX incorporated under each motif is representative of the frequency of the motif in the remainder of the population. Total coverage has not been adjusted to account for unknown gene types.

2. Number of epitopes represents a minimal estimate, considering only the epitopes shown in Table 6. Additional alleles possibly bound by nested epitopes have not been accounted.

TABLE Ia

SUPERMOTIFS	POSITION	POSITION	POSITION
	2 (Primary Anchor)	3 (Primary Anchor)	C Terminus (Primary Anchor)
A1	<b>T, I, L, V, M, S</b>		<b>F, W, Y</b>
A2	<b>V, Q, A, T</b>		<b>I, V, L, M, A, T</b>
A3	<b>V, S, M, A, T, L, I</b>		<b>R, K</b>
A24	<b>Y, F, W, I, V, L, M, T</b>		<b>F, I, Y, W, L, M</b>
B7	<b>P</b>		<b>V, I, L, F, M, W, Y, A</b>
B27	<b>R, H, K</b>		<b>F, Y, L, W, M, I, V, A</b>
B58	<b>A, T, S</b>		<b>F, W, Y, L, I, V, M, A</b>
B62	<b>Q, L, I, V, M, P</b>		<b>F, W, Y, M, I, V, L, A</b>
MOTIFS			
A1	<b>T, S, M</b>		<b>Y</b>
A1		<b>D, E, A, S</b>	<b>Y</b>
A2.1	<b>V, Q, A, T*</b>		<b>V, L, I, M, A, T</b>
A3.2	<b>L, M, V, I, S, A, T, F, C, G, D</b>		<b>K, Y, R, H, F, A</b>
A11	<b>V, T, M, L, I, S, A, G, N, C, D, F</b>		<b>K, R, H, Y</b>
A24	<b>Y, F, W</b>		<b>F, L, I, W</b>

\*If 2 is V, or Q, the C-term is not L

Bolded residues are preferred, italicized residues are less preferred: A peptide is considered motif-bearing if it has primary anchors at each primary anchor position for a motif or supermotif as specified in the above table.

WHAT IS CLAIMED IS

1. A composition comprising a prepared hepatitis C virus (HCV) epitope consisting of an amino acid sequence selected from the group consisting of:

FLLADARV,	YLVAYQATV,	RLIVFPDLGV,
DLMGYIPLV,	WMNRLIAFA,	VLVGGVLAA,
HMWNFISGI,	ILAGYGAGV,	YLLPRRGPR,
LLFLLADA,	YLVTRHADV,	KTSESRQPR,
RLGVRATRK,	QLFTFSPRR,	RMVVGVEHR,
LIFCHSKKK,	GVAGALVAFK,	VAGALVAFK,
TLGFGAYMSK,	LPGCSFSIF,	LSAFSLHSY,
CTCGSSDLY,	FWAKHMWNF,	SKGWRLAPITAYAQ,
AAAYAAQGYKVLVLNPSVAAT,	GRHLIFCHSKKKCDE,	VVVVATDALMTGYTG,
TVDFSLDPTFTIETT,	NFISGIQYLAGLSTLPGNPA,	
GEGAVQWMNRLIAFASRGNHV,	GSSYGFQYSPGQORVE,	ASCLRKLGVPPLRVW,

and LTCGFADLMGY.

2. The composition of claim 1, further comprising two epitopes selected from the group in claim 1.

3. The composition of claim 2, further comprising three epitopes selected from the group in claim 1.

4. The composition of claim 1, wherein the composition further comprises a CTL epitope selected from the group consisting of LTDPSHITA, LADGGCSGGAY, RMILMTHFF, VMGSSYGF, FWAKHMWNFI, LLFNILGGWV, IPFYGKAI, and VGIYLLPNR.

5. The composition of claim 1, wherein the composition further comprises an HTL epitope.

6. The composition of claim 5, wherein the HTL epitope is a pan DR binding molecule.

7. The composition of claim 1, wherein the epitope is on or within a liposome.
8. The composition of claim 1, wherein the peptide is joined to a lipid.
9. The composition of claim 1, wherein the epitope is bound to an HLA heavy chain,  $\beta$ 2-microglobulin, and streptavidin complex, whereby a tetramer is formed.
10. The composition of claim 1, wherein the epitope is bound to an HLA molecule on an antigen presenting cell.
11. The composition of claim 10, wherein the antigen presenting cell is a dendritic cell.
12. The composition of claim 1, the composition further comprising a pharmaceutical excipient.
13. The composition of claim 1, further wherein the epitope is in a unit dose form.
14. A composition comprising a prepared peptide of less than 250 amino acid residues comprising at least two hepatitis C virus (HCV) peptide epitopes selected from the group consisting of:

FLLADARV,	YLVAYQATV,	RLIVFPDLGV,
DLMGYIPLV,	WMNRLIAFA,	VLVGGVLAA,
HMWNFISGI,	ILAGYGAGV,	YLLPRRGPR,
LLFLLADA,	YLVTRHADV,	KTSESRQPR,
RLGVRATRK,	QLFTFSPRR,	RMVVGGEHR,
LIFCHSKKK,	GVAGALVAFK,	VAGALVAFK,
TLGFGAYMSK,	LPGCSFSIF,	LSAFSLHSY,
CTCGSSDLY,	FWAKHMWNF,	SKGWRLAPITAYAQ,

AAAYAAQGYKVLVLNPSVAAT, GRHLIFCHSKKKCDE, VVVVATDALMTGYTG,  
TVDFSLDPTFTIETT, NFISGIQYLAGLSTLPGNPA,  
GEGAVQWMNRLIAFASRGNHV, GSSYGFQYSPGQRVE, ASCLRKLGVPLRVW,  
and LTCGFADLMGY.

15. The composition of claim 14, wherein at least two epitopes are linked via a spacer.
16. The composition of claim 14, further comprising a third epitope.
17. The composition of claim 16, wherein the third epitope is selected from the group consisting of LTDPSHITA, LADGGCSGGAY, RMILMTHFF, VMGSSYGF, FWAKHMWNFI, LLFNILGGWV, IPFYGKAI, and VGIYLLPNR.
18. The composition of claim 16, further comprising a third epitope that is an HTL epitope.
19. The composition of claim 18, wherein the HTL epitope is a panDR binding molecule.
20. The composition of claim 14, wherein the peptide is on or within a liposome.
21. The composition of claim 14, wherein the peptide is joined to a lipid.
22. The composition of claim 14, wherein the peptide further comprises at least three of the epitopes in the group of claim 14.
23. The composition of claim 14, wherein the peptide further comprises at least four of the epitopes in the group of claim 14.
24. The composition of claim 14, wherein the peptide further comprises at least five of the epitopes in the group of claim 14.

25. The composition of claim 14, wherein the peptide further comprises at least six of the epitopes in the group of claim 14.

26. The composition of claim 14, the composition further comprising a pharmaceutical excipient.

27. The composition of claim 14, further wherein the epitope is in a unit dose form.


28. A composition comprising at least six prepared HCV epitopes each consisting of an amino acid sequence selected from the group consisting of:

FLLADARV,	YLVAYQATV,	RLIVFPDLGV,
DLMGYIPLV,	WMNRLIAFA,	VLVGGVLAA,
HMWNFISGI,	ILAGYGAGV,	YLLPRRGPR,
LLFLLADA,	YLVTRHADV,	KTSERSQPR,
RLGVRATRK,	QLFTFSPRR,	RMVVGVEHR,
LIFCHSKKK,	GVAGALVAFK,	VAGALVAFK,
TLGFGAYMSK,	LPGCSFSIF,	LSAFSLHSY,
CTCGSSDLY,	FWAKHMWNF,	SKGWRLAPITAYAQ,
AAYAAQGYKVLNPSVAAT,	GRHLIFCHSKKKCDE,	VVVVATDALMTGYTG,
TVDFSLDPTFTIETT,	NFISGIQYLAGLSTLPGNPA,	
GEGAVQWMNRLIAFASRGNHV, GSSYGFQYSPGQORVE, ASCLRKLGVPLRVW,		
and LTCGFADLMGY.		

29. The composition of claim 28, further comprising at least one epitope selected from the group consisting of LTDPSHITA, LADGGCSGGAY, RMILMTHFF, VMGSSYGF, FWAKHMWNFI, LLFNILGGWV, IPFYGKAI, and VGIYLLPNR.

1 / 2

# Monte Carlo population coverage analysis for HCV candidate epitopes



Plot of total frequency of genotypes as a function of the number of HCV candidate epitopes bound by HLA-A and B alleles, in an average population. Genotype values were derived by averaging the gene frequencies in Caucasian, North American, Black, Japanese, Chinese, and Hispanic populations. Also shown is the cumulative frequency of genotypes.

Using currently available HLA typing data, a residual fraction (about 15%) of the genes, in an average population, are unspecified. To arrive at 100% accounting of genes, a fraction of the residual has been added for each hit population cluster in proportion to the relative frequency of the cluster within the HLA specified population. One peptide, 24.0086, was not incorporated into the present analysis.

FIG. 1



2 / 2

# HVC Minigene

## CTL Epitopes

Core 43	NS4 1590	NS3 1128	NS5 2611	Core 169	NS1/E2 632	NS4 1765	NS4 1863	Core 132
Kozak	1073.11	1013.02	1069.62	1090.02	1145.12	1073.13	24.0092	1073.10
	A3	A2	A1	A2	B7	A3	A24	A3
								A2

NS3 1253	NS4 1921	1437	NS5 2641	1466
1283.21	1283.44	35.0106	1283.55	35.0107
DR	DR	DR3	DR	DR3

## HTL Epitopes

FIG. 2

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/19774

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(7) : A61K 38/00, 38/04, 38/08, 38/10, 39/29, 39/295 US CL : 514/2, 12, 13, 14, 15, 885; 424/185.1, 189.1 According to International Patent Classification (IPC) or to both national classification and IPC														
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/2, 12, 13, 14, 15, 885; 424/185.1, 189.1  Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) MEDLINE, BIOSIS, EMBASE, DERWENT WPI, WEST 2.0, search terms: author names, hcv, peptid?, HLA, htl, ctl,														
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>														
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim N												
Y	WENTWORTH et al. Differences and similarities in the A2.1-restricted cytotoxic T cell repertoire in humans and human leukocyte antigen-transgenic mice. Eur. J. Immunol. 1996. Vol 26. pages 97-101, see entire document.	1-29												
Y	US 5,736,142 A (SETTE et al.) 07 April 1998, see entire document.	1-29												
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.														
<table border="0"><tr><td>* Special categories of cited documents:</td><td>*T* later document published after the international filing date or prior date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td></tr><tr><td>*A* document defining the general state of the art which is not considered to be of particular relevance</td><td>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td></tr><tr><td>*E* earlier document published on or after the international filing date</td><td>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document combined with one or more other such documents, such combination being obvious to a person skilled in the art</td></tr><tr><td>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td><td>*G* document member of the same patent family</td></tr><tr><td>*O* document referring to an oral disclosure, use, exhibition or other means</td><td></td></tr><tr><td>*P* document published prior to the international filing date but later than the priority date claimed</td><td></td></tr></table>			* Special categories of cited documents:	*T* later document published after the international filing date or prior date and not in conflict with the application but cited to understand the principle or theory underlying the invention	*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document combined with one or more other such documents, such combination being obvious to a person skilled in the art	*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family	*O* document referring to an oral disclosure, use, exhibition or other means		*P* document published prior to the international filing date but later than the priority date claimed	
* Special categories of cited documents:	*T* later document published after the international filing date or prior date and not in conflict with the application but cited to understand the principle or theory underlying the invention													
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone													
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document combined with one or more other such documents, such combination being obvious to a person skilled in the art													
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family													
*O* document referring to an oral disclosure, use, exhibition or other means														
*P* document published prior to the international filing date but later than the priority date claimed														
Date of the actual completion of the international search 19 SEPTEMBER 2000		Date of mailing of the international search report 17 OCT 2000												
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer RON SCHWADRON Telephone No. (703) 308-0196												